RESEARCH ARTICLE

Open Access



Predictive equation of metastasis in patients with malignant ovarian epithelial tumors with the Ca-125 marker

Juan Fernando Sánchez Vega¹, Magdali del Rocío Murillo Bacilio^{2,3}, Adrián Santiago Vintimilla Condoy¹, Araceli Miroslava Palta González^{2,4}, José Alfredo Crespo Astudillo⁵ and Franklin Geovany Mora-BravoMsC^{3*}

Abstract

Background: Cancer antigen (CA) 125 (CA-125) is used in ovarian cancer detection and monitoring, whose serum level has a positive correlation with tumor stage. The aim of this study was to obtain a prediction metastasis equation in a group of patients with ovarian cancer based on Ca-125.

Methods: A 2-group comparative observational study was conducted at a single oncologic institution (SOLCA) in Cuenca-Ecuador. All patients who were diagnosed with ovarian cancer between January 1996 and December 2016 were included in the current study. Group 1 (G1) patients with the I and II International Federation of Gynecology and Obstetrics (FIGO) stage and Metastasis Group (MG), with III and IV stage, were subdivided. A logistic regression equation was performed to predict metastasis based on Logarithm of serum Ca-125 levels.

Results: We included 85 cases in G1 and 64 patients in MG, with 47.8 ± 15 years (G1) and 57.5 ± 13.6 years (MG) of age (P < 0.001). Mortality in G1 was 2 cases (3.1%) and 53 cases (62.4%) in MG (P < 0.001). The CA-125 serum level was 163.5 ± 236 in G1 and 1220.9 ± 1940 u / ml in MG (P < 0.001). The equation to predict metastasis = (Age*0.053) + [(Logarithm Ca-125 value) * 1.078] - 8.163 with an OR 2.940 (Cl 95% 2.046-4.223) P < 0.001. The sensitivity of the equation was 82.4% and the specificity was 79.7%.

Conclusions: It is possible to predict the presence of metastasis in a group of patients with ovarian cancer based on Ca-125.

Keywords: Prediction of metastasis, Ovarian Cancer, Cancer antigen 125

Background

Ovarian cancer is one of the ten leading causes of death worldwide, this pathology occurs in all ages including childhood and adolescence, most cases are diagnosed in very advanced stages [1]. In the detection and monitoring of ovarian cancer, the tumor marker Ca-125 is used, but because of its low sensitivity (70%) and specificity (90%) [2], it is used in conjunction with other methods such as imaging such as transvaginal doppler ultrasound [3, 4], improving the diagnosis up to 90% of cases of ovarian cancer [4, 5]. On the other hand, it has been used as a prognostic factor for recurrence and survival,

BM

Full list of author information is available at the end of the article



© The Author(s). 2018 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

^{*} Correspondence: franklin.mora@ucuenca.edu.ec

³School of Medicine, University of Cuenca, Av. 12 de abril and Paraíso Street, 010201 Cuenca, Ecuador

clinical stage IV, all patients with serous subtype show abnormal Ca-125 values. The total proportion of abnormal Ca-125 values (stages I to IV) was 89% in the endometrioid subtype of ovarian carcinoma, while patients with FIGO stage I endometrioid carcinoma showed 19% of Ca levels. -125 normal. As in the serous subtype, endometrioid ovarian carcinoma shows 100% abnormal Ca-125 in clinical stage IV [4]. There is a relationship between the tumor stage and the increase in the value of Ca-125, with 50% of patients in stage I having an increase, 75% in stage II, 90% in stage III and 98% in Stage IV [7]. There is a need to establish an index to improve cancer care in low/middle income countries where there is limited/no availability of cross-sectional imaging - an accurate index would facilitate triage of patients to a facility with the necessary oncological expertise for optimal surgery and/or chemotherapy (including potentially neo-adjuvant chemo). With this background, the present study aims to perform an equation to predict patients with ovarian cancer who have metastasized on the basis of Ca-125 as a predictor variable.

Methods

The present in an observational study conducted at the Cancer Institute SOLCA in the city of Cuenca - Ecuador, province of Azuay. Clinical histories of patients diagnosed with Ovarian Cancer who were treated in the 1996–2016 period were reviewed. Cases with insufficient data were excluded for the analysis. We collected demographic variables, type of ovarian cancer, stage and serum levels of Ca-125. The classification of the stages was the one adopted by the FIGO [11] described below to form the study groups:

Group 1: patients with stage I and II.

Group 2: patients with stage III and IV.

Stage I: Stage in which the tumor is confined to the ovaries.

Stage II: Stage in which the tumor involves one or both ovaries, with extension adjacent to pelvic tissues.

Stage III: The tumor involves one or both ovaries, with cytology or confirmed histology of spreading out of the pelvis and with metastases to the retro peritoneum and / or lymph nodes.

Stage IV: Advanced stage of cancer in which there is metastasis confirmed at a distance that excludes peritoneal metastasis.

The analysis of post-harvest information was tabulated and processed using the IBM SPSS Statistics license-free software updated to version 15.0. For the analysis of qualitative variables such as sex, residence, histological type, stage, value of Ca-125 and associated factors, a distribution of relative and absolute frequencies was

Table 1	Demographic d	ata and background	of the study	groups
---------	---------------	--------------------	--------------	--------

	Group 1Stages I & II-FIGO	Group 2 Advanced ovarian cancer group Stages III & IV-FIGO	Р
	n = 85	n = 64	
Civil Status			
Single	20 (23.5%)	17 (26.6%)	0.829
Free union	3 (3.5%)	3 (4.7%)	
Married	42 (49.4%)	34 (53.1%)	
Divorcee	4 (4.7%)	2 (3.1%)	
Widow	16 (18.8%)	8 (12.5%)	
Place of residence			
Cuenca	39 (45.9%)	24 (37.5%)	0.399
Machala	9 (10.6%)	4 (6.3%)	
Gualaceo	3 (3.5%)	1 (1.6%)	
Paute	1 (1.2%)	3 (4.7%)	
Azogues	5 (5.9%)	7 (10.9%)	
Others	28 (32.9%)	25 (39.1%)	
Cancers Family Antecedents	20 (23.5%)	18 (28.1%)	0.524
Breast Cancer Family History	4 (4.7%)	6 (9.4%)	0.493
Hormonal Therapy History	3 (3.5%)	3 (4.7%)	0.722
Biological state			
Premenopausal vs Postmenopausal	20 (23.5%)	27 (42.2%)	0.015

performed. For analysis of the relationship between the value of Ca-125 versus age, histological type, stage and associated factors, we used chi-square and bivariate analysis. CA-125 value was transformed into a logarithm of base "e" for the regression eq. *P* values less than 0.05 were considered statistically significant. Logistic regression was performed between the groups of Stages I and II versus Stages III and IV to predict the presence of the last stages using Ca-125. The project did not imply any risk for the patients since we worked on clinical history only and this report ensures the confidentiality of

	Table 2 Histologic	al type of Ovary	^r Cancer in the	study groups
--	--------------------	------------------	----------------------------	--------------

Variable	Group 1 Stages I & II-FIGO n = 85	Group 2 Advanced ovarian cancer group Stages III & IV-FIGO n = 64	Р
Histological type			
Serous	75 (88.2%)	46 (71.9%)	0.024
Mucinous	9 (10.6%)	10 (15.6%)	
Endometroid	0	5 (7.8%)	
Clear cells	1 (1.2%)	1 (1.6%)	
Transitional cells	0	2 (3.1%)	

Table 3 Binary logistic

regression between Metastasis and Ca-125 logarithm								
3	S.E.	Wald	df	Sig.	Exp(B)	95% C.I.for EXP(B)		
						Lower	Upper	

	В	S.E.	Wald	df	Sig.	Exp(B)	95% C.I.for EXP(B)	
							Lower	Upper
Step 2ª								
Age	0.053	.015	11.847	1	0.001	1.054	1.023	1.087
LOG CA-125	1.078	.185	34.010	1	< 0.0001	2.940	2.046	4.223
Constant	-8.163	1.415	33.295	1	< 0.0001	0.000		

^aVariable(s) entered on step 1: Age, LOG CA.125

identity and the management of the database. The Bioethics Committee of the Cancer Institute SOLCA-Cuenca approved the present study as well as obtaining the permission of the director of the Hospital.

Results

In the study, 149 cases were enrolled, of which 85 patients were classified as having ovarian cancer without metastasis and 64 patients in the metastasis group. The mean age of the group without metastasis was $47.8 \pm$ 15 years and the group without metastasis was 57.5 \pm 13.6 years (P < 0.001). With a median of 4 pregnancies in group 1 with an interquartile range of 5 and with 4 pregnancies and an interquartile range of 5 in group 2 (P =0.796). There were no differences between marital status and place of residence between the groups (Table 1). Additionally, the family history of cancer was not different between the groups (Table 1). The histological type of ovarian cancer predominant in both groups was serous (Table 2), with a greater group of women in premenopause in group 1 (Table 1). In group 1 there were 52 patients (81.3%) in Stage I and 12 patients (18.8%) in stage II, in group 2 there were 49 patients (57.6%) in stage III and 36 patients (42.4%) in stage IV. Mortality in group 2 was higher than in group 1 (Table 2). The Ca-125 antigen was positive in 42 cases in group 1 (65.6%) and in 79 cases (92.9%) in group 2 (P < 0.001), with an average of 163.5 ± 236 u / ml in group 1 and of 1220.9 ± 1940 u / ml in group 2 (*P* < 0.001).

Metastasis probability

The metastasis probability (≥ 0.50) was established as:

$$P = \frac{1}{1 + exp^{-F}}$$

Where F is the logistic regression equation (Table 3 and Additional file 1):

 $F = (Age^{*}0.053) + [(logarithm of Ca-125) *1.078] - 8.163.$

Were excluded non-significant variables named: "Pregnancies number" and biological status "Menopause" (Table 4). An internal validation was performed with the bootstrap method (Table 5).

Diagnostic test of the metastasis prediction equation

A 2×2 contingency table was constructed. The data of groups 2 and 1 were placed in columns and the metastasis prediction equation in rows (Presence and Absence). Table values were (a) 70 cases (b) 13 cases, (c) 15 cases and (d) 51 cases. This table reported a sensitivity of 82.4%, and a specificity of 79.7%. The positive predictive value of 84.3% and the negative predictive value of 77.2%. The positive likelihood ratio was 4.059 and the negative likelihood ratio was 0.22. The ROC curve reported an Area under the curve of 0.870 (95% CI 0.812-0.928) with a standard error of 0.03 and P < 0.0001 (Fig. 1).

Discussion

The main finding of the present study was that high levels of Ca-125 are related to the presence of metastases in this group of patients diagnosed with Ovarian Cancer. The Odds Ratio of this relationship is 2.940 (CI 95% 2.046–4.223) *P* < 0.001 (Table 3), the logistic regression equation was statistically significant and establishes the serum level of Ca-125 as a statistically significant

Table 4 Binary logistic regression between Metastasis and Ca-125 logarithm, Age, Pregnancies number, Menopause

, 0			0					
	В	B S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for EXP(B)	
							Lower	Upper
Step 1 ^a								
Age	0.096	0.028	11.656	1	0.001	1.101	1.042	1.163
Pregnancies number	-0.098	0.083	1.406	1	0.236	.907	.771	1.066
LOG CA-125	1.034	0.185	31.236	1	< 0.0001	2.812	1.957	4.040
Premenopause/Menopause [1]	1.240	0.736	2.843	1	0.092	3.457	.818	14.619
Constant	-10.27	1.975	27.045	1	< 0.0001	.000		

^aVariable(s) entered on step 1: Age, Pregnancies Number, LOG_ca125 (logarithm), State of Premenopause

Table 5 Bootstrap for Variables in the Equation

	В	Bootstrap ^a						
		Bias	Bias Std.	Sig. (2-	95% Confidence Interva			
			Error tailed)		Lower	Upper		
Step 1								
LOG CA- 125	1.078	0.049	0.215	0.001	0.781	1.639		
Age	0.053	0.002	0.018	0.002	0.024	0.095		
Constant	-8.163	-0.364	1.777	0.001	-12.597 -5.754			
^a Pootstran results are based on 1000 bootstran samples								

^aBootstrap results are based on 1000 bootstrap samples

predictor of the presence of metastasis. The equation had a positive predictive value of 84.3% with a positive likelihood ratio of 4.059. The ROC Curve reported an Area under the curve of 0.87 which was statistically significant. The cut-off point of the Ca-125 value to predict the presence of metastases in this study group was 240 u / ml. In the equation, non-prognostic variables were excluded, such as the fact of dying, and spurious variables such as family history and origin were excluded. The significance of finding an association between the stage of metastasis (State II and III of the FIGO classification) and the serum levels of Ca-125 allows us to have a practical and applicable equation in daily clinical practice. It has been established that the serum levels of Ca-125 pretreatment are increased, showing a positivity of 55% in the serous types, and between 42 to 77% in stage III [4] and stage IV (15%), being these mostly serous epithelial type [12]. In the present study, serous epithelial



cancer was predominant (81.2%), and most of it was in stage III, which coincides with FIGO statistics [13] due to late diagnosis [14]. In the study it was found that the stage with the highest elevation of the marker was stage III (36.4%), this may be due to the fact that in advanced stages there is metastasis, while in the histological type it was the serous (85.27%), significant statistical relationship observed in other studies [14] however in no previous work an equation has been established to predict metastasis as in the present study. It has been shown that the association between Metastasis in patients with ovarian cancer and serum Ca-125 levels fits into the causality, since the tumor burden is intruded. The relationship is less associated with stages in which the load or tumor mass has been decreased in therapeutic form and may be an alternative explanation to the reason for decreasing the association with advanced stages already operated therapeutically. In daily practice, the equation helps to quickly predict the state of metastasis of patients and finally the mortality since the group with metastasis had higher mortality than the group without metastasis. This study does not include other tumor markers used in ovarian cancer. Future research should prospectively address the prediction equation to predict metastasis and compare it with a gold standard clinical.

Conclusions

There is a relationship between the clinical stage of metastasis and serum Ca-125 levels in this group of patients with ovarian cancer.

Additional files

Additional file 1: Excel sheet for calculating the probability of ovarian cancer metastasis. (XLSX 9 kb) Additional file 2: Database in Excel format. (XLSX 19 kb)

Abbreviations

CA-125: Cancer antigen 125; FIGO: International Federation of Gynecology and Obstetrics

Acknowledgements

Acknowledgment to the SOLCA-workers who participated indirectly in the study. Acknowledgment to Bernardo Vega, Dean of the Faculty of Medical Sciences of the School of Medicine for the economic management for financing.

Funding

The economic source of this research and publication was provided by School of Medicine at the University of Cuenca. I clarify that the funds allocated by the University of Cuenca did not participate in the design of the study, data collection, analysis, interpretation of the data or in the writing of the manuscript. The funds were allocated solely for the translation of the manuscript and to cover the costs of publishing the article.

Availability of data and materials

The data supporting the findings of this study are available upon request from the main author. The database is attached in Additional file 2.

Authors' contributions

Research Idea: JFSV, MRMB, FGMB. Study design: JFSV, ASVC, AMPG, FGMB.Bibliographic review: JFSV, ASVC, JACR. Data collection: JFSV, ASVC, JACRHealing of the data: MRMB. Data analysis: FGMB. Draft writing: JFSV. Critical analysis of the document: MRMB, AMPG, FGMB. All authors read and approved the final version of the article.

Authors' information

Juan Fernando Sánchez Vega: Rural Doctor Ministry of Public Health of Ecuador. Av May 1st 357 and Carlos Quinto. jfsanchezvega@icloud.com orcid.org/0000-0003-1582-4638

Magali del Rocío Murillo Bacilio. rocimurillo4@yahoo.com.mx Affiliation: Medical Pathologist from the Department of Pathology SOLCA-Cuenca. Professor of Pathology at the school of medicine, University of Cuenca. orcid.org/0000-0003-3183-7906

Adrián Santiago Vintimilla Condoy. a.dri_an@hotmail.com Rural Doctor Ministry of Public Health of Ecuador. orcid.org/0000-0002-7016-2262 Araceli Miroslava Palta González aracelimpg@hotmail.com Medical Pathologist from the Department of Pathology SOLCA-Cuenca. Head of Department. orcid.org/0000-0003-0516-1172

José Alfredo Crespo Astudillo. Resident Doctor Pablo Jaramillo Foundation. Franklin Mora-Bravo, MD, MsC. Main professor of nephrology at the school of medicine, University of Cuenca. Master's degree in health research.orcid.org/ 0000-0002-5978-3420

Ethics approval and consent to participate

The research protocol was approved by the bioethics committee of the University of Cuenca and the Bioethics Committee of the SOLCA-Cuenca Institute. A written consent to participate was requested to study patients. Only participants who signed the informed consent forms were included in the study.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Rural Doctor Ministry of Public Health of Ecuador, Zone 6, Cuenca, Ecuador.
²Department of Pathology of the Institute for the Fight against Cancer
Society -SOLCA-Cuenca, Cuenca, Ecuador. ³School of Medicine, University of
Cuenca, Av. 12 de abril and Paraíso Street, 010201 Cuenca, Ecuador. ⁴School of Medicine, University of Azuay, Cuenca, Ecuador. ⁵Pablo Jaramillo
Foundation, Complementary Health Network, Cuenca, Ecuador.

Received: 29 January 2018 Accepted: 10 May 2018 Published online: 24 May 2018

References

- Trétarre B, Molinié F, Woronoff AS, Bossard N, Bessaoud F, Marrer E, et al. Ovarian cancer in France: trends in incidence, mortality and survival, 1980-2012. Gynecol Oncol. 2015;139(2):324–9. https://doi.org/10.1016/j.ygyno. 2015.09.013.
- Coussy F, Chéreau E, Daraï E, Dhombres F, Lotz JP, Rouzier R, et al. Interest of CA 125 level in management of ovarian cancer. Gynecol Obstet Fertil. 2011;39(5):296–301. https://doi.org/10.1016/j.gyobfe.2010.10.018.22.
- Paul P, Talukder S, Sangma MA, Saha PL, Begum M, Razi AZ, et al. Detection of ovarian tumor by color Doppler ultrasonography and CA-125. Mymensingh Med J. 2017;26(4):705–9. PubMed PMID: 29208855.
- Morales-Vásquez F, Pedernera E, Reynaga-Obregón J, López-Basave HN, Gómora MJ, Carlón E, et al. High levels of pretreatment CA125 are associated to improved survival in high grade serous ovarian carcinoma. J Ovarian Res. 2016;9:41. Available from: http://www.ncbi.nlm.nih.gov/pmc/ articles/PMC4936203/.
- Varras M. Benefits and limitations of ultrasonographic evaluation of uterine adnexal lesions in early detection of ovarian cancer. Clin Exp Obstet Gynecol. 2004;31(2):85–98. Review. PubMed PMID: 15266758.
- Dolores-Velázquez R, Bustamante-Montes LP, Romero-Fierro JR, Gómez-Plata E, Ocampo-García KG, Dueñas-González A. Relación entre niveles de CA-125

y características clínicas e histopatológicas en el cáncer epitelial de ovario. Gac Mex Oncol. 2014;13(4):207–14.

- Tiwari RK, Saha K, Mukhopadhyay D, Datta C, Chatterjee U, Ghosh TK. Evaluation of preoperative serum levels of CA 125 and expression of p53 in ovarian neoplasms: a prospective Clinicopathological study in a tertiary care hospital. J Obstet Gynaecol India. 2016;66(2):107–14. https://doi.org/10.1007/ s13224-014-0611-7.
- Schmidt C. CA-125: a biomarker put to the test. J Natl Cancer Inst. 2011; 103(17):1290–1.
- Worzfeld T, Pogge von Strandmann E, Huber M, Adhikary T, Wagner U, Reinartz S, Müller R. The unique molecular and cellular microenvironment of ovarian Cancer. Front Oncol. 2017;7:24. https://doi.org/10.3389/fonc.2017.00024.
- Hauptmann S, Friedrich K, Redline R, Avril S. Ovarian borderline tumors in the 2014 WHO classification: evolving concepts and diagnostic criteria. Virchows Arch. 2017;470(2):125–42. https://doi.org/10.1007/s00428-016-2040-8.
- Prat J, Belhadj H, Berek J, Bermudez A, Bhatla N, Cain J, et al. Staging classification for cancer of the ovary, fallopian tube, and peritoneum. Int J Gynaecol Obstet. 2014;124(1):1–5. https://doi.org/10.1016/j.ijgo.2013.10.001.
- Willis S, Villalobos VM, Gevaert O, Abramovitz M, Williams C, Sikic BI, et al. Single gene prognostic biomarkers in ovarian Cancer: a meta-analysis. PLoS One. 2016;11(2):e0149183.
- Prat J. FIGO's staging classification for cancer of the ovary, fallopian tube, and peritoneum: abridged republication. J Gynecol Oncol. 2015;26(2):87–9. https://doi.org/10.3802/jgo.2015.26.2.87.
- Zivanovic O, Sima C, Iasonos A, Bell-McGuinn K, Sabbatini P, Leitao M, et al. Exploratory analysis of serum CA-125 response to surgery and the risk of relapse in patients with FIGO stage IIIC ovarian Cancer. Gynecol Oncol. 2009;115(2):2–17. https://doi.org/10.1016/j.ygyno.2009.06.038.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

