

## Global use of Ethnomedicinal Plants to Treat Toothache

José Luis Álvarez-Vásquez<sup>1,2,\*</sup>, Nathaly Fernanda Parra-Solano<sup>1,2</sup>,  
Gabriela Elizabeth Saavedra-Cornejo<sup>1,2</sup> and Ximena Elizabeth Espinosa-  
Vásquez<sup>1,2</sup>

<sup>1</sup>Especialización en Endodoncia, Facultad de Odontología, Universidad de Cuenca, Ecuador.

<sup>2</sup>Facultad de Odontología, Universidad de Cuenca, Cuenca, Ecuador.

\*Corresponding Author E-mail: jose.alvarezv@ucuenca.edu.ec

<https://dx.doi.org/10.13005/bpj/2421>

(Received: 27 April 2022; accepted: 20 June 2022)

Toothache is one of the most common global health problems, and medicinal plants are widely used to relieve the associated pain and inflammation. Several studies have been conducted on the use of plants to treat toothache, but no study has comprehensively assessed the types of plants and the mechanisms of action of the phytochemical compounds involved in their analgesic effect. This review aims to bridge this gap. This is the first review to collect a large volume of data on the global use of medicinal plants used in the treatment of toothache. It presents the relevant information for dentists, researchers, and academics on using medicinal plants to treat toothache. We found that preclinical studies and state-of-the-art technology hold promise for furthering our knowledge of this important topic. In total, 21 species of medicinal plants used to treat toothache were found in America, 29 in Europe, 192 in Africa, 112 in Asia, and 10 in Oceania. The most common species were *Allium sativum*, *Allium cepa*, *Acmella oleracea*, *Jatropha curcas*, *Jatropha gossypifolia*, and *Syzygium aromaticum*. The most commonly found family of medicinal plants was Asteraceae, followed by Solanaceae, Fabaceae, Lamiaceae, Euphorbiaceae, Rutaceae, and Myrtaceae. The most common phytochemicals found were flavonoids, terpenes, polyphenols, and alkaloids. The reported mechanisms of action involved in toothache analgesia were antioxidant effects, effects mediated by transient receptor potential channels, the  $\gamma$ -aminobutyric acid mechanism, and the cyclooxygenase/lipoxygenase anti-inflammatory mechanism.

**Keywords:** Dental Pain; Flavonoids; Medicinal Plants; Phytochemicals; Toothache.

Toothache is an unpleasant sensory and emotional experience<sup>1</sup>, originating in the tooth or adjacent structures and is caused by factors such as caries, periodontal disease, trauma, or dentoalveolar abscess<sup>2</sup>. It is one of the most common health problems worldwide<sup>3</sup>. Toothache has a higher prevalence in lower socioeconomic groups, in whom this disease is not always adequately treated<sup>4</sup>, and in developing countries, where access to healthcare is limited. This has

led many local communities to resort to using alternatives for toothache relief, such as medicinal plants<sup>3</sup>.

Medicinal plants are widely used in dental practices. The World Health Organization has reported that between 65% and 80% of the population in developing countries use them to reduce inflammation, inhibit oral pathogen growth, and trigger anti-inflammatory, antiseptic, antioxidant, and analgesic effects<sup>3, 4</sup>. Several

phytochemical studies conducted on these plants have identified compounds such as flavonoids, alkaloids, and terpenes, which reduce toothache through their mechanism of action<sup>3, 5-7</sup>.

Phytotherapy is the use of plants to treat diseases or as health-promoting agents. When used for this purpose, their original composition and integrity are generally preserved, so that an entire plant or a desired percentage of its components may be used for medicinal purposes, fulfilling a specific mechanism of action, generally, a specific pathway to relieve pain [8]. However, to our knowledge, no study has comprehensively tackled the mechanisms of action of the phytochemical compounds contained in medicinal plants used to treat toothache. This integrative review aimed to bridge this gap by compiling and analyzing the different studies available in the literature.

## MATERIALS AND METHODS

The available literature in PubMed, PMC, and Scopus databases was searched to identify relevant articles on medicinal plants used to relieve toothache, published in English until July 31, 2021, using the search terms toothache, dental pain, medicinal plants, medicinal herbs, and phytochemicals. Articles unrelated to the use of plants that relieved toothache or lacking data for at least one of the following characteristics were excluded: family, scientific name, plant parts used, and method of preparation.

Of a total of 300 articles, 80 met the inclusion criteria and were comprehensively analyzed for this review. In addition, we performed a manual search of the reference lists of the initially selected articles to complement the available information and found 294 additional articles. Ten books with relevant information were also included. Regional medicinal plant types retrieved from the articles and books were summarized by continents. Finally, owing to length restrictions, this review did not include information related to the possible adverse reactions and drug interactions resulting from the use of the plants included in this review.

### Medicinal plants for toothache treatment

For several millennia, plants have been used in traditional dentistry to treat toothache, periodontal disease, herpetic ulcers, stomatitis,

maxillary sinusitis, and other ailments<sup>6</sup>. In recent years, advances in science and technology have identified the phytochemical compounds in some of these plants and their mechanisms of action<sup>3</sup>. Phytochemicals are a large group of plant-derived chemical substances that have various biochemical and physiological effects that are beneficial for human health and nutrition<sup>6, 9</sup>.

Phytochemicals found in plants vary greatly in number, structural heterogeneity, and distribution, and they are classified into polyphenols, carotenoids, alkaloids, terpenes, and terpenoids<sup>10, 11</sup>. All the tables in this review outline the phytochemicals described in previous reports on medicinal plants used to treat toothache, focusing on their analgesic mechanisms of action.

### Plant parts and preparation method

As mentioned above, plants are used to treat diseases through phytotherapy, using either the entire plant or a desired percentage of its components [8]. The most commonly used parts of medicinal plants are the leaves, seeds, flowers, and roots. The roots, in particular, are highly important because they are higher in bioactive compound content than other plant parts<sup>3, 12-14</sup>.

Leaves contain high concentrations of secondary metabolites, phytochemicals, and essential oils that have various health benefits [14]. Hence, most of the research studies support the use of leaves instead of roots because root extraction threatens the conservation of several plant species, especially those that are widely used<sup>3, 14</sup>.

There is considerable variation in the preparation methods of plants used to treat toothache, and the most common methods of administration are: using the plant extract, chewing, crushing, and drinking a decoction<sup>3</sup>.

### Mechanisms of action of phytochemical compounds

Phytochemicals such as flavonoids, alkaloids, and terpenes<sup>3, 5</sup> are biologically active compounds found in plants that work through various mechanisms of action<sup>15, 16</sup>. Based on the information gathered in this review, the most salient mechanisms of action of phytochemicals used to treat toothache were antioxidant activity<sup>9, 17</sup>, action on transient receptor potential channels (TRP)<sup>18</sup>,  $\gamma$ -aminobutyric acid (GABA) mechanism<sup>19, 20</sup>, and anti-inflammatory mechanisms (cyclooxygenase (COX) and lipoxygenase (LOX) pathways)<sup>21</sup>.

### Antioxidant activity

In living organisms, reactive oxygen species (ROS) are generated during metabolism and do not generally cause oxidative damage to cellular components due to the action of antioxidants present in these organisms<sup>22</sup>.

Natural antioxidants are found in various plants and play a key role in stopping the generation of free radicals by preventing the oxidation of biomolecules in the body. Therefore, they are valuable therapeutic agents for preventing diseases caused by oxidative stress. The latter causes an imbalance that favors the production of prooxidants, represented by ROS, such as superoxide anions ( $O_2^-$ ), hydrogen peroxide ( $H_2O_2$ ), and hydroxyl radicals ( $OH^-$ )<sup>23</sup>, which damage key cellular components, such as DNA, proteins, and membrane lipids, and can even trigger cell death<sup>17, 24-27</sup>.

Conversely, during inflammatory processes, free radicals balance themselves by attacking the nearest stable molecule and “stealing” an electron. The attacked molecule then becomes a free radical by losing its electron and initiating a cascade of cell-damaging reactions<sup>24</sup>. Additionally, leukocytes present in damaged regions cause a “respiratory burst” from enhanced oxygen uptake, and inflammatory cells generate inflammatory mediators that act on the infection site to release more reactive species<sup>24, 28, 29</sup>.

Therefore, the role of antioxidants is to delay, prevent, or eliminate oxidative damage of target molecules by controlling the levels of free radicals and other reactive species<sup>30</sup>. Plants are responsible for our oxygenated environment, and because they are exposed to high intracellular levels of oxygen and ROS, they have developed specialized defense systems (antioxidants) to protect their structures and tissues. Antioxidant activity is inherent to all plants as they act to prevent, destroy, or neutralize free radicals<sup>17</sup>.

These antioxidant defense systems can be *enzymatic complexes and non-enzymatic systems*. Some enzymatic complexes are superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), and glutathione reductase (GR). *Non-enzymatic systems* consist of low-molecular-weight antioxidants such as ascorbic acid, glutathione, proline, carotenoids, phenolic

acids, and flavonoids; and high-molecular-weight secondary metabolites, such as tannins, which efficiently prevent the toxic effects of free radicals<sup>31, 32</sup>.

The phytochemicals in plants can act as antioxidants by directly eliminating ROS, chelating metals (Fe, Zn, Mg, and Mn), quenching the mitochondrial respiratory chain, and increasing the levels of endogenous antioxidant enzymes, such as SOD, CAT, and GPx<sup>9, 31</sup>.

ROS and reactive nitrogen species (RNS) are key players in various types of pain<sup>33</sup>. Evidence suggests that tissue injury induces the production of both ROS and RNS, which cause pain by promoting neuronal excitability in pain pathways through neural interactions and by triggering mitochondrial dysfunction and neuro-inflammation<sup>26, 34</sup>.

Peroxyntirite ( $ONOO^-$ ) (PN) and its precursor superoxide (SO) are critical in the development of chronic pain and in the transition from acute to chronic pain [35]. An increase in SO/PN production triggers thermal hyperalgesia associated with acute and chronic inflammation in response to the activation of the *N*-methyl-D-aspartate receptor (NMDAR), leading to the development of orofacial pain<sup>36</sup>.

PN improves protein kinase C (PKC) activity. This kinase is activated by peripheral and central sensitization and optimizes the translocation of regulatory subunits of NADPH oxidase to the cell membrane to increase the production of SO derived from NADPH oxidase. These two mechanisms together amplify the formation of SO-derived PN, leading to the development of central sensitization<sup>35</sup>.

Thus, antioxidants can be administered for pain management to prevent the negative impact of ROS and RNS on nociception, both of which play key roles in neuro-inflammatory processes both at the central and peripheral levels, leading to increased nociceptive and inflammatory responses<sup>26, 33, 37, 38</sup>.

In addition to their antioxidant activity, flavonoids and phenolic compounds exhibit effective anti-inflammatory biological properties by blocking two main signaling pathways, NF- $\kappa$ B and mitogen-activated protein kinase (MAPK)<sup>24</sup>. These pathways initiate a cascade of phosphorylation events and result in the production

of several pro-inflammatory mediators that mediate the transmission of extracellular signals from the membrane to the nucleus<sup>24, 27, 39</sup>.

#### Action on TRP channels

TRP channels are involved in various homeostatic and sensory functions, such as nociception and temperature sensation, and are expressed in both neuronal and non-neuronal cells. They are grouped into six subfamilies: TRP ankyrin (TRPA), TRP canonical (TRPC), TRP melastatin (TRPM), TRP mucolipin (TRPML), TRP polycystin (TRPP), and TRP vanilloid (TRPV). They are mostly non-selective cation channels expressed on the cell membrane, including the TRPA1 channel, a Ca<sup>2+</sup>-permeable channel expressed in sensory neurons and is activated by phytochemicals and multiple products of oxidative stress<sup>18</sup>.

The Ca<sup>2+</sup>-permeable TRP channels of presynaptic terminals can modulate synaptic transmission independent of action potentials. Thus, the TRP channels, TRPV1, and TRPA1 can cause the release of neurotransmitters at sensory nerve terminals where these channels are highly co-expressed and participate in inflammatory hyperalgesia<sup>18, 40</sup>. Capsaicin (hot pepper), allicin (garlic), camphor (*Cinnamomum camphora*), rosemary, and menthol (peppermint) are all analgesics that excite and desensitize nociceptive sensory neurons by acting on the TRPA1 and TRPV1 channels<sup>41-43</sup>.

Other phytochemicals also activate the TRP channels. For example, curcumin (*Curcuma longa*) activates TRPA1 channels; eugenol activates the TRPV1 and TRPV3 channels; menthol activates TRPM8 channels; ginger components activate the TRPV1 and TRPA1 channels; and the thymol and linalool compounds of thyme (*Thymus vulgaris*) activate the TRPV3 and TRPA1 channels<sup>18</sup>.

#### GABA mechanism

GABA is a major inhibitory neurotransmitter<sup>44</sup>, involved in most inhibitory actions in the central and peripheral nervous systems (CNS and PNS). GABA exerts its action through two types of receptors: ionotropic (GABAA and GABAC) and metabotropic (GABAB) receptors. GABAA and GABAC are ion channels found in CNS neurons that are permeable to chloride ions when activated by GABA. GABAB receptors belong to the superfamily of G protein-coupled

receptors and are present at different levels of the pain neuraxis where they regulate nociceptive transmission and pain<sup>19, 45, 46</sup>.

Some phytochemicals, including flavonoids and terpenes, modulate the function of ionotropic GABA receptors and can act as positive, negative, and neutralizing allosteric modulators. Thus, herbal preparations such as *Heliopsis longipes*, *Acmella caulirhiza*, *Ginkgo biloba*, *Panax ginseng*, and *Scutellaria lateriflora* may help modulate toothache by crossing the blood-brain barrier and influencing brain function. Past research has suggested that an increase in GABAergic activity in the rostral agranular insular cortex may induce analgesia by enhancing the descending inhibition of spinal cord nociceptive neurons<sup>19, 47</sup>.

*Spilanthes acmella* is a flowering herb species, also known as the “toothache plant”<sup>48-50</sup>. It has been used for centuries to treat oral pain owing to its analgesic, anti-inflammatory, and anesthetic properties attributed to its bioactive compounds, especially phytosterols, phenolic compounds, and *N*-alkylamides<sup>48, 50, 51</sup>. Spilanthol, which is mainly present in the flowers and shoots of *S. acmella*, is the most representative compound found in this genus. This plant species and other species such as *H. longipes* are used worldwide as traditional remedies for their analgesic, antinociceptive, antioxidant, and anti-inflammatory effects. The analgesic effect of this compound is attributed to GABA release in the temporal cerebral cortex, whereas the antinociceptive effect is caused by the activation of the opioid-adrenergic, serotonergic, and GABAergic systems<sup>52</sup>.

The flavonoid baicalein, which can be extracted from *S. lateriflora*, exerts sedative and anxiolytic effects by binding to GABAA receptors and, hence, could be used to manage orofacial pain. This flavonoid is also believed to modulate both intra- and extracellular calcium levels, which play key roles in pain signaling and transmission<sup>44</sup>.

GABA receptor systems are found in peripheral pathways and the spinal cord, which are both important sites for pain impulse formation and transmission; they are located in the marginal zone and substantia gelatinosa of the dorsal horn, which are essential for interpreting and responding to pain signals. These findings indicate that GABA plays a key role in nociceptive processing. Consequently,

agents that modify the function of this inhibitory neurotransmitter are used as analgesics<sup>46</sup>.

#### **Anti-inflammatory mechanism (COX and LOX pathways)**

Inflammation is mediated by several families of mediators such as eicosanoids, which are lipid mediators produced through arachidonic acid metabolism, primarily in the COX and LOX pathways<sup>53</sup>. The COX pathway leads to the formation of prostanoids (prostaglandins (PG), prostacyclin, and thromboxane), whereas the LOX pathway leads to the production of leukotrienes (LTs)<sup>54</sup>.

Nonsteroidal anti-inflammatory drugs (NSAIDs) inhibit the COX pathway, whereas other drugs such as licofelone are dual inhibitors that block both COX and LOX<sup>54,55</sup>. However, the selective inhibition of the two COX isoforms by NSAIDs has several reported side effects. This has encouraged the search for a dual inhibitor of both COX-2 and 5-LOX that possesses improved anti-inflammatory potency and fewer side effects<sup>53,56</sup>.

This anti-inflammatory effect leads to the elimination of harmful stimuli and the restoration of normal physiology through the complex molecular cascade mentioned above<sup>3,21</sup>. This is thought to be the mechanism by which herbal extracts act in the treatment of toothache<sup>21</sup>. Accordingly, medicinal plants, particularly herbs whose main component is curcumin, such as *C. longa*, seem to provide several advantages through their mediating action on the COX and LOX pathways. As a dual inhibitor, curcumin exhibits synergistic effects and optimal anti-inflammatory activity<sup>57</sup>. *Allium cepa* (onion), which also contains polyphenols and flavonoids, inhibits the COX and LOX pathways and prevents the formation of LTs, thromboxane B2 (TXB2), and prostaglandin E2 (PGE2)<sup>58,59</sup>.

Additionally, various ginger compounds, such as gingerols, shogaols, zingerones, gingerdiols, and paradols, exhibit antioxidant, analgesic, and anti-inflammatory activities. More specifically, they act through the inhibition of COX and LOX in addition to their antioxidant activity resulting in an analgesic effect<sup>60</sup>. *Allium sativum* also has antioxidant and anti-inflammatory properties, and its efficacy in reducing pro-inflammatory responses is based on its nature as a COX and LOX inhibitor<sup>59</sup>.

#### **Bioavailability of medicinal plants**

In humans, most phytochemicals exhibit low bioavailability after ingestion<sup>9</sup>. Hence, polyphenols have a rather low bioavailability because they exert most of their antioxidant activity in the gastrointestinal tract<sup>61</sup>. Additionally, a challenge with flavonoids is their low water solubility, which leads to decreased absorption and consequently decreased bioavailability following oral administration<sup>62</sup>.

Interindividual variability, which depends on several factors such as diet, genetic background, composition, and activity of the intestinal microbiota, must also be considered. For example, polyphenols are relatively poorly absorbed (0.3–43%), resulting in low circulating plasma concentrations of their metabolites<sup>63</sup>. Additionally, the quantity and composition of phytochemicals in plants are influenced by species, age, plant part, cultivation method, harvesting season, conservation method, and geographic distribution<sup>9,64</sup>.

To improve bioavailability, proper decoction practices and various plants combinations have been suggested<sup>65</sup>. due to their different phytochemical components and because they may provide different health benefits without requiring an increase in the dose<sup>9</sup>. For example, *Piper sarmentosum* combined with ginger is used to soothe toothache<sup>66</sup>. Medicinal plants containing hundreds of phytochemicals can produce many metabolites in the body, exerting more efficient beneficial effects than individual phytochemicals<sup>9</sup>. However, their combination can also directly affect their bioavailability in the body via mechanisms such as the first-pass effect<sup>18,67</sup>.

#### **Medicinal plants vs pharmaceutical drugs**

Comparisons between the analgesic effects of medicinal plants and pharmaceutical drugs have shown that the rhizome of *Zingiber officinale* (ginger) has long been used in traditional Chinese and Indian medicine to treat a wide range of ailments, including toothache<sup>68</sup>. Fresh ginger extracts have been subjected to chromatographic purification, and the resulting fractions were analyzed to assess their effect on PG synthesis. Through this method, plant extracts belonging to the Zingiberaceae family were found to inhibit PG synthesis *in vitro*<sup>69</sup>.

The rhizome of *Z. officinale* has pharmacological properties similar to those of dual-action NSAIDs [55]. It inhibits both COX and LOX and has significantly fewer side effects than conventional NSAIDs<sup>69, 70</sup>. Licofelone is an example of a dual-action NSAID (5-LOX/COX) that is currently in phase III clinical development<sup>55</sup>. Studies have shown that orally administered dry ginger or ginger extract can reduce acute inflammation<sup>68, 71</sup>, and *in vitro* and *in vivo* comparisons have confirmed the anti-inflammatory and analgesic actions of ginger extract<sup>69, 70</sup>.

However, most *in vitro* studies analyzed phytochemical profiles using indices such as the half-maximal inhibitory concentration (IC<sub>50</sub>), and the medicinal plant extracts have been tested for only a single biological target, COX or LOX. This is insufficient to validate their anti-inflammatory and analgesic properties and hinders direct comparisons between plants and dual-action NSAIDs<sup>72</sup>.

The anti-inflammatory properties of ginger extracts come from a mixture of biologically active components such as gingerols, shogaols, and paradols, which are phenolic compounds<sup>73</sup>. The inhibitory effects of ginger on PG synthesis can be attributed to the presence of hydroxymethoxyphenyl compounds in gingerols and shogaols, which in turn inhibit arachidonic acid metabolism via the COX pathway<sup>69, 74</sup>. Moreover, ginger components inhibit several genes encoding cytokines and chemokines involved in inflammatory responses<sup>69, 75</sup>.

Essential oil from the fruit of the plant *Dennettia tripetala* (DT), commonly known as pepper fruit, has analgesic effects like those induced by opioids morphine, aspirin, and indomethacin. The analgesic mechanism of DT has been inferred from studies showing that naloxone, which inhibits the analgesic effect of morphine, could also inhibit DT. These findings suggest that DT can also be used for toothache relief<sup>22</sup>.

#### **Plant combinations**

Mixtures of medicinal plants are a key field of research which accounts for a large volume of information because their polyvalent effects can be used to cure multicausal diseases<sup>76</sup>. In different regions and cultures, plants are used as the entire plant, a combination of plants, or a combination of a plant and a drug. When medicinal plants are mixed, side effects are more likely to happen

because interactions can occur between individual components. The most desirable interactions provide additional therapeutic benefits. However, natural extracts also contain multiple components. Therefore, the effects of interactions between two plants are often unpredictable and complex<sup>77</sup>.

Additionally, a combination of two or more phytochemicals does not always enhance a specific effect. Combining two or more active chemical substances can produce additive, synergistic, or antagonistic effects<sup>9, 78</sup>. An example of synergism in the use of medicinal plants is Iberogast®, a phyto-preparation used in European countries consisting of nine plant extracts. It is considered to have a multi-target effect (at the gastrointestinal level). Such a multi-target effect has advantages over that of synthetic single-target drugs<sup>79, 80</sup>.

Another example is the phytotherapeutic drug Lenidase®, which, when compared to ibuprofen, more efficiently and safely controls postoperative pain and discomfort following third molar extraction. Lenidase® contains a blend of herbal extracts, such as baicalin (190 mg), bromelain (50 mg), and escin (30 mg)<sup>81</sup>, which exhibit anti-inflammatory activities. Bromelain inhibits pain mediators such as PGE2 and substance P and exhibits anti-edematous activity. Baicalin regulates several genes associated with inflammation, such as *COX*, *LOX*, and the inducible nitric oxide synthase gene. Escin exerts anti-inflammatory and anti-edematous effects through antihistaminic and antiserotonergic activities<sup>81</sup>.

## **RESULTS AND DISCUSSION**

As discussed above, medicinal plants, their phytochemicals, and their mechanisms of action are key subjects of scientific research because they are used to treat and prevent various diseases. Further, plants are the basis of many drugs. Although they are highly complex compounds and are not always suitable substitutes for synthetic agents<sup>82</sup>, phytochemicals have been used to provide relief from toothache in various regions of the world, as outlined in the five tables included in this review.

The components of the medicinal plants used and their preparations vary by location and between species. For example, a study conducted

in America revealed that leaves were the most commonly used plant parts and that the most common preparations were pastes, extracts, and rinses<sup>83</sup>. Another study conducted in Africa found that *Datura stramonium* L. roots, leaves, stems,

and seeds were often used to provide relief from toothache<sup>3</sup>. These findings demonstrate the need for further phytochemical and pharmacological studies to identify the plant part that is most effective for toothache treatment and the optimal application




Fig. 1. Relative frequency of the global use of plant parts for the treatment of toothache




Fig. 2. Families of medicinal plants globally used for the treatment of toothache

Table 1. America

Family	Scientific name	Used part	Preparation	Phytochemicals	Analgasic effect	Anti-inflammatory effect	Mechanism of action	Ref.
Fabaceae	<i>Acacia farnesiana</i> (L.) Willd.	Stem, leaves	Filling, extract, rinse	Diterpenes, flavonoids	x	x	Antioxidant activity	[83]
Asclepiadaceae	<i>Asclepias curassavica</i> L.	Latex, leaves, stem	Filling, extract, rinse	Flavonoids, triterpenes, phenols, quinnins, tannins	x	x	N/A	[83, 100]
Malpighiaceae	<i>Byrsonima crassifolia</i> (L.)	Leaves, flowers	Filling, extract, rinse	Triterpenes, sterols, flavonoids	x	x	Antioxidant activity	[83, 101, 102]
Solanaceae	<i>Capiscum frutescens</i> L.	Leaves	Filling, extract, rinse	Alkaloids, tannins, saponins, flavonoids, carotenoids, sterols	x	x	Antioxidant activity	[83, 103]
Chenopodiaceae	<i>Chenopodium graveolens</i> (Willd.)	Leaves	Filling, extract, rinse	Sterols, flavonoids	x	x	N/A	[83]
Sterculiaceae	<i>Chromolaobolus pentadactylon</i> Lam.	Root	Filling, extract, rinse	Flavonoids, phenolic compounds	x	x	Antioxidant activity	[83]
Moraceae	<i>Dorstenia contrajerva</i> L.	Root	Filling, extract, rinse	Flavonoids, alkaloids	x	x	N/A	[83, 104]
Asteraceae	<i>Helipopsis longipes</i>	Root	Filling, extract, rinse	Alkarnides, affinin (spilanthol)	x	x	GABA mechanism, anti-inflammatory mechanism (COX and LOX)	[83, 105]
Campanulaceae	<i>Lobelia laxiflora</i> Kunth.	Whole plant	Filling, extract, rinse	N/A	x	x	N/A	[83]
Lauraceae	<i>Persea americana</i> Miller.	Fruit	Filling, extract, rinse	Phenolics, flavonoids	x	x	Antioxidant activity	[83, 106]
Malvaceae	<i>Sida rhombifolia</i> L.	Stem, leaves	Filling, extract, rinse	Flavonoids, terpenoids, alkaloids	x	x	N/A	[83, 107]
Sterculiaceae	<i>Theobroma cacao</i> L.	Grain	Filling, extract, rinse	Polyphenols	x	x	Antioxidant activity	[83]
Polygonacea	<i>Mexican Sanguinaria</i>	Root	Filling, extract, rinse	Flavonoids, lignans, lignins.)	x	x	N/A	[83]
Euphorbiaceae	<i>Jatropha gossypifolia</i>	Whole plant	Decoction	Alkaloids, terpenes, flavonoids, lignans, coumarins, phenolics	-	x	Antioxidant activity	[108]
Euphorbiaceae	<i>Jatropha curcas</i>	Sap	Topical	Alkaloids, terpenes, flavonoids, lignans, coumarins, phenolics	-	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[108, 109]
Amaryllidaceae	<i>Allium sativum</i> L.	Bulb	Crushed	Steroids, terpenoids, flavonoids, phenols	-	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[18, 59, 110, 111]
Araliaceae	<i>Panax ginseng</i>	Leaves, seeds	Extract	Flavonoids, terpenes	-	x	GABA mechanism	[110]
Zingiberaceae	<i>Zingiber officinale</i> roscoe	Leaves, seeds	Extract	Flavonoids, terpenes	-	x	Anti-inflammatory mechanism (COX and LOX)	[60, 110]
Asteraceae	<i>Tagetes lucida</i>	Aerial parts	Decoction, infusion, rinse, direct application	Carotenoids, flavonoids, thiophenes	-	x	Antioxidant activity	[112, 113]
Onagraceae	<i>Lopezia racemosa</i>	Leaves, seeds	Infusion	Tannins, flavonoids	-	x	Antioxidant activity	[114, 115]
Asteraceae	<i>Acmella oleracea</i>	Flowers	Chewed	Spilanthol, sequesterpenes.	x	x	GABA mechanism, antioxidant activity	[51, 116]



Table 2. Europe

Family	Scientific name	Used part	Preparation	Phytochemicals	Analgasic effect	Anti-inflammatory effect	Mechanism of action	Ref.
Amaryllidaceae	<i>Allium cepa</i> L.	Bulb	Bandage	Saponin, quercetin, anthocyanin	-	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[6, 59, 117]
Amaryllidaceae	<i>Allium sativum</i> L.	Bulb	Crushed	Alliin, allicin, flavonoids, terpenes	-	x	Antioxidant activity, TRP mechanism, anti-inflammatory mechanism (COX and LOX)	[6, 18, 59, 76, 111]
Asteraceae	<i>Antennaria dioica</i> (L.)	Leaves	Decoction	Phenolic, flavonoid	-	x	Antioxidant activity	[6, 118]
Brassicaceae	<i>Armoracia rusticana</i> P.	Root	Finely chopped, chew	Flavonoids	-	-	Antioxidant activity	[6, 119]
Brassicaceae	<i>Brassica oleracea</i> L.	Leaves	Rinse, gargle	Carotene, tocopherol, ascorbate	-	-	Antioxidant activity	[6, 120]
Lauraceae	<i>Cinnamomum verum</i> , <i>J. Presl</i>	Bark	Mix with cinnamon powder, honey, rub and spread	Monoterpenes, diterpenes, sesquiterpenes, polyphenols	-	x	Antioxidant activity	[6, 121]
Myrtaceae	<i>Eucalyptus globulus</i> Labill.	Leaves	Infusion mixed with lemon, inhalation	Flavonoids	-	x	Antioxidant activity	[6, 122]
Solanaceae	<i>Hyoscyamus niger</i> L.	Leaves	Infusion, gargle	Alkaloids	x	x	Antioxidant activity	[6, 123]
Asteraceae	<i>Matricaria chamomilla</i> L.	Flowers	Infusion, gargle	Flavonoids, coumarins, sesquiterpenes, and polyacetylenes	-	x	Antioxidant activity	[6, 124, 125]
Ranunculaceae	<i>Nigella arvensis</i> L.	Oil, seed	Dissolved in hot water	Phenols	-	x	Antioxidant activity	[6, 126]
Plantaginaceae	<i>Plantago lanceolata</i> L.	Leaves	Gargle	Phenolic compounds	-	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[6, 127, 128]
Lamiaceae	<i>Salvia officinalis</i> L.	Leaves	Decoction, gargle	Phenolic compounds	-	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[6, 129, 130]
Myrtaceae	<i>Syzygium aromaticum</i> L.	Flowers	Chewed	Phenolic compounds	-	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[6, 131, 132]
Asteraceae	<i>Taraxacum</i> sp.	Leaves, root	Boil, drop, or chew	Phenolic compounds	x	x	Antioxidant activity	[6, 133]
Vitaceae	<i>Vitis vinifera</i> L.	Fruit	Decoction, gargle	Phenols	-	x	Antioxidant activity	[6, 134]
Poaceae	<i>Zea mays</i> L.	Grain	Boil, gargle	Flavonoids, alkaloids, phenols, steroids, terpenoids, tannins	x	x	Antioxidant activity	[6]
Malvaceae	<i>Althaea officinalis</i> L.	Root	Tisane	N/A	-	-	N/A	[76]
Oleaceae	<i>Olea europaea</i> L.	Fruit	Juice, aerosol	Phenolic compounds	x	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[76, 135]
Asteraceae	<i>Calendula arvensis</i> L.	Flowers	Poultice	N/A	-	-	N/A	[76]
Apiaceae	<i>Eryngium campestre</i> L.	Root	Tisane	N/A	-	x	N/A	[76]
Anacardiaceae	<i>Pistacia lentiscus</i> L.	Leaves	Mouthwash	N/A	x	x	N/A	[76]
Lamiaceae	<i>Origanum vulgare</i> L.	Flowers	Tisane	N/A	x	x	N/A	[76]
Rutaceae	<i>Ruta chalepensis</i> L.	Aerial parts	Tisane	Alkaloids, flavonoids, phenols, saponins	x	x	N/A	[76, 136]
Rosaceae	<i>Crataegus monogyna</i> Jacq.	Flowers	Poultice	Flavonoids, phenolic compounds	-	x	N/A	[76]
Apiaceae	<i>Foeniculum vulgare</i> Mill.	Seed	Poultice	Flavonoids, phenolic compounds	-	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[76, 137, 138]
Fabaceae	<i>Glycyrrhiza glabra</i> L.	Root	Poultice	N/A	-	-	N/A	[76]
Papaveraceae	<i>Papaver rhoeas</i> L.	Flowers	Poultice	N/A	-	-	N/A	[76]
Lamiaceae	<i>Thymus vulgaris</i> L.	Aerial parts	Fumigation	N/A	-	-	TRP mechanism	[18, 76]
Rutaceae	<i>Citrus limon</i> L.	Fruit	Juice, mouthwash	N/A	-	-	N/A	[76]

Table 3. Africa.

Family	Scientific name	Used part	Preparation	Phytochemicals	Analgesic effect	Anti-inflammatory effect	Mechanism of action	Ref.
Ebenaceae	<i>Euclea undulata</i> Thunb.	Root, bark	Powder rubbed or root chewed	Flavonoids, saponins, diterpenes, alkaloids	-	x	Antioxidant activity	[139]
Fabaceae	<i>Milletia ferruginea</i> (Hochst.)	Bark	Grind, mix with water, chew	Flavonoids	-	-	N/A	[140, 141]
Lamiaceae	<i>Salvia nilotica</i> Juss.	Leaves, root, bark	Chewing, grinding, rubbing	N/A	-	x	N/A	[140, 142]
Lamiaceae	<i>Robeya myricoides</i> (Hochst.)	Leaves, root, bark	Rubbing, grinding, eating, boiling	N/A	-	x	N/A	[140]
Malvaceae	<i>Ceiba pentandra</i> L. Gaertn	Fruit	Grinding, hold on teeth	Flavonoids	x	x	N/A	[140, 143]
Acanthaceae	<i>Blepharis maritima</i> Pers.	Aerial part	Filling with powder	Flavonoids, phenolic acids	-	x	N/A	[144, 145]
Acanthaceae	<i>Barleria homotricha</i> C. B. Clarke	Bark	Drink	N/A	-	-	N/A	[3]
Acanthaceae	<i>Dyschoriste radicans</i> (Hochst. Ex. Rich.) Nees	Whole plant	N/A	N/A	-	-	N/A	[3]
Acanthaceae	<i>Justicia Schimperiana</i> (Hochst. ex Nees) T. Anderson	Twigs	Chewed	Saponins, alkaloids, terpenoids and flavonoids	-	-	Antioxidant activity	[3, 146]
Amaryllidaceae	<i>Allium sativum</i> L.	Bulb	Crushed	Alliin, alliein, flavonoids, terpenes	-	x	Antioxidant activity, TRP mechanism, anti-inflammatory mechanism (COX and LOX)	[3, 18, 111]
Amaranthaceae	<i>Amaranthus caudatus</i> L.	Bark, leaves	Chewed	Flavonoids, phenols	-	-	Antioxidant activity (COX and LOX)	[140, 147]
Amaryllidaceae	<i>Allium cepa</i> L.	Leaves	Paste	Flavonoids, saponin	-	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[59, 148, 149]
Anacardiaceae	<i>Mangifera indica</i> L.	Stem, root, leaves	Solution	N/A	-	x	N/A	[148, 150]
Anacardiaceae	<i>Rhus natalensis</i> Benth. Ex. Krauss	Leaves	Chewed	Flavonoids	-	x	Antioxidant activity	[3, 151-153]
Anacardiaceae	<i>Schinus molle</i> L.	Stem	Brushing	Monoterpenes, Naphthylisoquinoline alkaloids	x	-	Antioxidant activity	[3, 154]
Ancistrocladaceae	<i>Ancistrocladus abbreviatus</i>	Bark	Boil crust	Tannins, alkaloids, flavonoids, terpenoids, saponins	x	-	N/A	[148, 155]
Ammonaceae	<i>Denonnetia tripetala</i>	Fruits, leaves, seeds, roots	Extract	Flavonoids	x	x	N/A	[22]
Apiaceae	<i>Anmi visnaga</i>	Fruit	Raw, decoction, direct application, rinse	Tannins, coumarins, flavonoids, phenolic acids	x	x	Antioxidant activity	[156, 157]
Apiaceae	<i>Coriandrum sativum</i> L.	Fruit	Raw, direct application	Polypheols; flavonoids	x	x	Antioxidant activity	[156, 158]
Apiaceae	<i>Foeniculum vulgare</i> Mill.	Roots	Decoction	Flavonoids, phenolic compounds	-	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[3, 137, 138]
Apiaceae	<i>Oenanthe pilustris</i> (Hiov.) C. Norman	Leaves	Chewed	N/A	-	-	N/A	[3]
Apocynaceae	<i>Carissa edulis</i> (Forsk.) Vahl	Root, leaves	Pound, boil and press on tooth	Lignans, sesquiterpenes, phenols	-	x	Antioxidant activity	[159, 160]
Apocynaceae	<i>Nerium oleander</i>	Leaves, root	Raw, direct application	Terpene, flavonoids	-	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[156, 161, 162]
Apocynaceae	<i>Carissa spinarum</i> L.	Bark	Chew or hold on teeth for 5-10 min.	Coumarin, lignans	-	x	Antioxidant activity	[163, 164]
Apocynaceae	<i>Calotropis procera</i> (ait.) Dryand.	Bark	Pounded	Triterpenoids	x	-	Antioxidant activity	[3, 165]
Aquifoliaceae	<i>Ilex mitis</i> (L.) Radlk.	Twigs	N/A	N/A	-	-	N/A	[3]
Anfiaceae	<i>Schefflera abyssinica</i> (Hochst. Ex. a. Rich.) Harms	Bark	Chewed	Saponins	-	-	N/A	[3]
Areaceae	<i>Cocos nucifera</i> L.	Root	Powder and whole root	Phenols, tannins, flavonoids, terpenes, steroids, alkaloid	-	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[148, 166-168]
Asclepiadaceae	<i>Gomphocarpus purpurascens</i> a. Rich.	Root	Mix with honey and chew	N/A	-	-	N/A	[169, 170]
Asparagaceae	<i>Asparagus africanus</i> Lam.	Root	Drink	Saponins, flavonoids, tannins.	x	x	N/A	[3, 171]
Asteraceae	<i>Echinops kebericho</i> , mesfin	Root	Pounded dry root is mixed with coffee	N/A	x	x	N/A	[172, 173]
Asteraceae	<i>Eriangen tomentosa</i>	Leaves	Crush and press on the tooth	Alkaloids, saponins, coumarins, tannins	-	-	N/A	[159]
Asteraceae	<i>Chamaemelum nobile</i> L.	Flowers	Decoction, mouthwash	Flavonoids	-	x	Antioxidant activity	[156, 174]
Asteraceae	<i>Artemisia absinthium</i> L.	Leaves	Decoction, mouthwash	Phenolic compounds	-	x	Antioxidant activity	[156, 175]
Asteraceae	<i>Arctocylis gummifera</i>	Root	Raw, direct application	Polypheols, flavonoids, tannins	-	x	Antioxidant activity	[156, 176]
Asteraceae	<i>Sphlanthes africana</i>	Flowers	All parts	Alkaloids, saponins and glycosides	x	-	N/A	[148, 177]
Asteraceae	<i>Echinops purpurea</i>	Leaves, stem	Paste	N/A	-	x	Antioxidant activity	[148]
Asteraceae	<i>Ageratum conyzoides</i>	Whole plant	Dust	Alkaloids, flavonoids, terpenoids	x	x	N/A	[148, 178]

Asteraceae	Dichrocephala integrifolia	Whole plant	Paste						N/A	[148, 179]
Asteraceae	Acmella cauliflora Del.	Flowers	Chewed, topical application						N/A	[3, 152]
Asteraceae	Acmella oleracea	Flowers	Chewed						GABA mechanism, antioxidant activity	[51, 51, 180]
Asteraceae	Artemisia abyssinica sch.bip. Ex a. Rich.	Stem	Chewed						Antioxidant activity	[3, 181]
Asteraceae	Artemisia afra Jack. Ex wild.	Leaves	Chewed						N/A	[3]
Asteraceae	Echinops kebericho mes'h	Roots	Powdering						Antioxidant activity	[3, 182]
Asteraceae	Echinops macrochaetus Fresen.	Roots	Hold in mouth						N/A	[3]
Asteraceae	Galinsooga parvi?ora Cav.	Flowers	Rubbing						Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[3, 183, 184]
Asteraceae	Imula conferti?ora a. Rich.	Leaves	Chewed						N/A	[3]
Asteraceae	Kleimia squarrosa Cufod.	Stem	Brushing						N/A	[3]
Asteraceae	Laggeria intermedia c. B. Clarke	Leaves	Cushed						N/A	[3]
Asteraceae	Parthenium hysterophorus L.	Roots	Chewed						N/A	[3]
Asteraceae	Vernonia aureoaliffen Hiem	Roots	Chewed						N/A	[3, 185]
Balanitaceae	Balanites aegyptiaca (L.) Del.	Bark	Chewed						Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[3, 186, 187]
Bignoniaceae	Sterrospermum kunthianum Cham	Stem	Chewed						Antioxidant activity	[3, 188]
Bomniaceae	Ehretia cymosa Thonn.	Bark, leaves	Chew, grmd, boil, squeeze						Antioxidant activity	[140, 189]
Bomniaceae	Cordia africana Lam.	Bark	Chewed						Antioxidant activity	[3, 190]
Bomniaceae	Cynoglossum coeruleum Hochst. Ex a. De	Leaves	Hold in mouth						N/A	[3]
Brassicaceae	Leptidium sativum L.	Seeds	Chewed						Antioxidant activity	[3, 191]
Burseraeae	Commiphora hodia Sprague	Roots	Inhaling						N/A	[3]
Capparidaceae	Boscia salicifolia Oliv.	Leaves	Chew						Antioxidant activity	[169, 192]
Capparidaceae	Capparis tomentosa Lam.	Roots	Chewed						Antioxidant activity	[3, 193]
Capparidaceae	Capparis fascicularis DC.	Roots	Chewed						N/A	[3]
Capparidaceae	Cadaba rotundifolia Forsk.	Leaves	Chewed						Antioxidant activity	[3, 194]
Capparidaceae	Crateva adansonii DC.	Leaves	Heating						Antioxidant activity	[3, 195]
Caricaceae	Carica papaya	Leaves	Solution						N/A	[148, 150, 196]
Caryophyllaceae	Drymaria cordata	Leaves	N/A						N/A	[3, 197]
Celastraceae	Catha edulis (Vahl) Endl.	Leaves	Eat, boil, chew, mix with water.						N/A	[140]
Chenopodiaceae	Chenopodium opulifolium	Leaves	Drink						Antioxidant activity	[3, 159, 198]
Chenopodiaceae	Chenopodium ambrosioides	Whole plant, seeds	Paste						Antioxidant activity	[148, 199]
Clusiaceae	Clusia lanceolata Cambess.	Leaves	N/A						N/A	[3]
Clusiaceae	Garcinia livingstonei T. Anderson	Stem	Chewed						Antioxidant activity	[3, 200, 201]
Colchicaceae	Gloriosa superba L.	Leaves	Crushed						Antioxidant activity	[3, 202]
Combretaceae	Angecossus leucarpus (DC.)	Bark	Filling with powder						Antioxidant activity	[144, 203]
Asteraceae	African Asphila	Whole plant	Paste						N/A	[148, 204]
Asteraceae	Vernonia amygdalina	Leaves	Solution, chewed						Antioxidant activity	[148, 205]
Asteraceae	Arnica montana	Leaves	Solution						Antioxidant activity	[148, 206]
Comvolvaceae	Ipomoea batatas	Leaves	Paste						N/A	[148]
Crossulaceae	Kalanche laciniata (L.) DC	Roots	Chewed						Antioxidant activity	[3, 207]
Cucurbitaceae	Cucumis ficifolius	Roots	Chewed						N/A	[169]
Cucurbitaceae	Momordica foetida Schumacher	Roots	Chewed						Antioxidant activity	[3, 208]
Cupressaceae	Cupressus bethanis	Seed, bark	Solution / rinse						N/A	[148]
Cupressaceae	Cupressus lusitana Mill.	Leaves	Decoction						Antioxidant activity	[3, 209]
Cupressaceae	Juniperus procera Hochst. Ex endl.	Bark	Hold						Antioxidant activity	[3, 210]

Dniacraenaceae	Dniacraea fragrans L.	Bark	Chewed	-	-	-	N/A	[159, 211]
Dniacraenaceae	Dniacraea distelliana	Root	Solution	-	-	-	N/A	[148]
Ebenaceae	Euclea racemosa	Whole plant	Chewed	-	-	-	Antioxidant activity	[3, 169]
Ebenaceae	Euclea divaricata Hiern	Roots	Drink	x	x	x	Antioxidant activity	[3, 212]
Euphorbiaceae	Jatropha gossypifolia	Whole plant	Decoction	-	-	-	Antioxidant activity	[108]
Euphorbiaceae	Jatropha curcas	Sap	Topical	-	-	x	Antioxidant activity, anti-inflammatory mechanism (COX-LOX)	[108, 109]
Euphorbiaceae	Ricinus communis	Leaves	Solution	x	-	x	Antioxidant activity	[148, 213]
Euphorbiaceae	Alchornea cordifolia	Stem, bark	Boiled	-	-	-	N/A	[148, 214]
Euphorbiaceae	Ricinodendron heudelotii	Seeds	Paste	-	x	x	Antioxidant activity	[148, 215]
Euphorbiaceae	Cordia alliodora (L.) Jacq.	Leaves	Hold in mouth	-	-	x	Anti-inflammatory mechanism (COX-LOX)	[163, 216]
Euphorbiaceae	Phyllanthus sepioides	Roots	Chewed	-	-	-	N/A	[3]
Euphorbiaceae	Acalypha sp.	Leaves	Boil	x	-	x	Antioxidant activity	[148, 217]
Fabaceae	Calpania aurea (aiton) benth.	Leaves, root	Chewing, rubbing, powdering, grinding	-	-	-	N/A	[140]
Fabaceae	Acacia albidia delile	Stem	Chewing, slow drying	-	-	x	Antioxidant activity	[140, 218]
Fabaceae	Albizia gummifera	Leaves	Crushed, rubbed	-	-	-	N/A	[163]
Fabaceae	Acacia mifera	Stem, bark	Decoction	-	-	-	Antioxidant activity	[3, 219-221]
Fabaceae	Alysicarpus orientalis Rich.	Bark	Chewed	-	-	-	N/A	[222]
Fabaceae	Erythrina brucei Schweinf.	Bark	Chewed	-	-	-	N/A	[3, 223]
Fabaceae	Indigofera spicata Forsk.	Roots	Chewed	-	-	-	N/A	[3]
Fabaceae	Dorysila abyssinica (L. Rich.) Wats.	Seeds	Chewed	-	-	-	N/A	[3]
Ficoutriaceae	Miconia parvifolia Schinz	Leaves	Heated	-	-	-	N/A	[3]
Geraniaceae	Geranium sp.	Leaves	Tea	-	-	-	Antioxidant activity	[3, 224]
Clusiaceae	Coccoloba kola	Bark and seeds	Tea	-	-	x	Antioxidant activity	[148, 225]
Lamiaceae	Coleus blumei	Leaves	Paste	-	-	x	Antioxidant activity	[148, 226]
Lamiaceae	Thymus schimperii Rominger	Whole plant	Chew	-	-	-	N/A	[148, 226]
Lamiaceae	Cleodendrum myricoides (Hook.)	Roots, seeds	Crushed	-	-	-	N/A	[3, 227]
Lamiaceae	Isodon junosissimus (Hook.)	Roots, leaves	Chewed	-	-	-	N/A	[3]
Lamiaceae	Mentha pulgiana L.	Leaves	Chewed, infusion	-	-	-	TRP mechanism	[3, 18, 156]
Lamiaceae	Ocimum urticifolium Roth	Leaves	Chewed	-	-	-	N/A	[3, 228]
Lamiaceae	Ocimum schimperii Rominger	Whole plant	Chewed	-	-	-	N/A	[3]
Lauraceae	Persea americana	Seeds, stem	Decoction / hot solution	-	-	x	Antioxidant activity	[106, 148, 229]
Fabaceae	Acacia oerfota (forsk.) Schweinf.	Root	Paste	-	-	-	N/A	[144]
Fabaceae	Acacia senegal (L.) Willd., Schweinf.	Leaves, thorns	Filling with powder and boiling	-	-	-	Antioxidant activity	[144]
Fabaceae	Cassia arethifolia	Root	Filling with powder	-	-	-	Antioxidant activity	[144, 230]
Loranthaceae	Phoradendron robustum	Leaves	Pulverized	-	-	-	N/A	[3]
Loranthaceae	Tapinanthus globiferus (L. Rich.) Tiegh.	Leaves	Rub	x	-	-	Antioxidant activity	[3, 231]
Malvaceae	Celastrus scandens	Bark and fruit	Solution and paste	-	-	x	Antioxidant activity	[148, 232]
Malvaceae	Pavonia urens Cav.	Root	Decoction	-	-	-	N/A	[3]
Malvaceae	Sida emuicarpa Vahl	Root	Crushed	-	-	-	N/A	[3]
Malvaceae	Sida emuicarpa Vahl	Leaves, root	Liquid form	-	-	-	N/A	[140]
Malvaceae	Stellaria media Forsk.	Leaves, root, bark	Grinding, chewing, boiling, liquid form	-	-	x	Antioxidant activity	[140, 233-235]
Meliaceae	Melia azadirachta L.	Leaves	Chewed	-	-	-	Antioxidant activity	[3, 236]

Menispermaceae	<i>Stephania abyssinica</i>	Root	Brushed	N/A	-	N/A	[3]
Lamiaceae	<i>Marrubium vulgare</i> L.	Whole plant	Decoction, mouthwash	Phenolic compounds	x	Antioxidant activity, anti-inflammatory mechanisms (COX-1/LOX)	[156, 237, 238]
Lamiaceae	<i>Origanum majorana</i>	Whole plant	Mouthwash	Terpenoids, flavonoids, phenolic acids, Phenolic compounds, flavonoid	x	Antioxidant activity	[156, 239]
Monaceae	<i>Ficus palmata</i> forssk.	Root	Chewed	N/A	-	Antioxidant activity	[3, 240]
Monaceae	<i>Ficus sur</i> forssk.	Bark	Chewed	Tannins, saponins, steroids, tannins	-	N/A	[3, 241]
Monaceae	<i>Ficus vasta</i> forssk.	Bark	Eating, chewing	Phenolic compounds, flavonoids	-	Antioxidant activity	[140, 242]
Moringaceae	<i>Moringa oleifera</i>	Root	Decoction	Alkaloids, protein, quinine, saponins, flavonoids, tannin, steroids, glycosides	x	Antioxidant activity	[148, 243]
Myristicaceae	<i>Pycnanthus angolensis</i>	Stem, bark, leaves	Decoction / mouthwash	N/A	-	N/A	[148]
Myristicaceae	<i>Eswalypus globulus</i> Labill.	Leaves	Infusion mixed with lemon, inhalation	Flavonoids	-	Antioxidant activity	[6, 122, 156]
Myristicaceae	<i>Myrtus communis</i> L.	Leaves	Decoction, mouthwash	Phenolic compounds, flavonoids	-	Antioxidant activity	[156, 244]
Myristicaceae	<i>Syzygium aromaticum</i> L.	Flowers	Chewed	Phenolic compounds	-	Antioxidant activity, anti-inflammatory mechanism	[6, 132, 156, 245]
Myrtaceae	<i>Eucalyptus saligna</i>	Leaves	Solution, paste	Flavonoids, derived from phenylpropene	-	N/A	[148]
Myrtaceae	<i>Psidium guajava</i> L.	Leaves	Hot rinse, chewed	Flavonoids, terpenes, tannins	x	Antioxidant activity	[93, 148, 219, 246]
Myrtaceae	<i>Eucalyptus sp.</i>	Roots	Rob	N/A	x	Antioxidant activity	[3, 247]
Myrtaceae	<i>Eucalyptus camaldulensis</i>	Leaves	Rubbing, hold on teeth, filling with powder	N/A	x	Antioxidant activity	[140, 144, 248]
Nitriaceae	<i>Peganum harmala</i> L.	Seeds	Decoction, mouthwash	Flavonoids, alkaloids, saponins, tannins, glycosides, terpenoids and steroids	x	N/A	[156, 249-251]
Oleaceae	<i>Ximenesia americana</i> L.	Bark	Polyverized	Flavonoids, polyphenols	-	Antioxidant activity	[3, 252]
Oleaceae	<i>Olea europaea</i> L.	Leaves	Chewed	Phenolic compounds	-	Antioxidant activity, anti-inflammatory mechanism	[130, 135, 169]
Oleaceae	<i>Jasminum abyssinicum</i> hochst	Roots	Chewed	Alkaloids, flavonoids, terpenes, phenol	-	N/A	[3, 253]
Oleaceae	<i>Jasminum grandiflorum</i> L.	Leaves, stem	Crushed	Flavonoids, alkaloids, glycosides	-	Antioxidant activity	[3, 254]
Oliniaceae	<i>Olinia rocheiana</i> a. Juss.	Leaves, bark	N/A	N/A	-	N/A	[3]
Opiliaceae	<i>Zarziphus mauritiana</i> lam	Stem	Boiled	Flavonoids, glycoside, phenol, lignin, saponins, sterols and tannins	-	Antioxidant activity	[3, 255]
Ombonchaceae	<i>Orobanchae ramosa</i> L.	Roots	Chewed	N/A	-	N/A	[3]
Oxalidaceae	<i>Oxalis corniculata</i> L.	Leaves	Chewed	Flavonoids, alkaloids, tannins, phenols	-	Antioxidant activity	[3, 256]
Oxalidaceae	<i>Oxalis radicata</i> A. Rich.	Leaves, stem	Chewed	N/A	-	N/A	[3, 152]
Phytolobaceae	<i>Phytolobaea dodecandra</i> L'her	Stem	Chewed	Triterpenoids, saponins, alkaloid, phenolics, steroids, terpenoids	x	N/A	[3, 257, 258]
Plumbaginaceae	<i>Plumbago zeylanica</i> L.	Roots, bark	Crushed root, direct application	Alkaloids, glycosides, steroids, terpenoids, tannins, phenolic compounds, flavonoids, saponins, coumarins	-	N/A	[152, 259, 260]
Polygalaceae	<i>Securidaca longepedunculata</i> Fresen.	Leaves	Chewed	Flavonoids, tannins, triterpenoids	x	N/A	[3, 261]
Polygonaceae	<i>Rumex abyssinicus</i> Jacq	Roots	Crushed, drink	Phenolic compounds, tannins, saponins, flavonoids, terpenoids, steroids, alkaloids	x	N/A	[3, 169, 262]
Polygonaceae	<i>Rumex nepalensis</i> Spreng.	Roots	Chewed	Flavonoids, terpenoids, alkaloids, saponins, tannins	-	Antioxidant activity	[3, 263]
Polyodiaceae	<i>Drynaria volkensii</i> Heiron	Roots	Chewed, heat	N/A	-	N/A	[3, 163]
Protocaeae	<i>Fuaria speciosa</i> Welw	Roots, leaves	Chewed	N/A	-	N/A	[3, 152]
Ranunculaceae	<i>Clematis longicauda</i> Steud	Leaves	Crushed	Tannins, saponins, flavonoids, and steroids	-	N/A	[3, 264]
Ranunculaceae	<i>Ranunculus sinesis</i> Fresen	Bark, seed, roots	Chewed	Triterpenoids, saponins, alkaloids, polyphenol	x	N/A	[3, 265]
Ranunculaceae	<i>Ranunculus multifidus</i> forssk.	Roots	Chewed	N/A	-	N/A	[3]
Ranunculaceae	<i>Thalictrum thymocarpum</i> dill	Roots	Chewed	N/A	-	N/A	[3]
Rosaceae	<i>Prunus africana</i> (hook & f)	Bark	Chewed	Phenolic, flavonoid	-	Antioxidant activity	[3, 266]
Rosaceae	<i>Rosa persica</i> (L.) Batsch	Bark	Chewed	N/A	-	N/A	[3, 267]
Rubiaceae	<i>Galium boreae-aethiopicum</i>	Roots	Chewed	Phenolic compounds, carotenoids	x	N/A	[169]
Rubiaceae	<i>Gardenia ternifolia</i> Schumacher	Roots	Chewed	N/A	-	N/A	[3]
Rubiaceae	<i>Pavetta gardeniifolia</i> hochst.	Roots	Crushed	Saponins, steroids, triterpenes, tannins, flavonoids	-	Antioxidant activity	[3]
Rubiaceae	<i>Pentas lanceolata</i> (forssk.)	Roots	Chewed	N/A	-	N/A	[3]
Rutaceae	<i>Clausena anisata</i> (willd.)	Roots, stem, seeds, leaves	Chewed	Saponins, alkaloids	-	N/A	[3, 152, 268]
Rutaceae				Phenolic, alkaloid, flavonoid, saponins, tannin	-	Antioxidant activity	[3, 269]

Rutaceae	<i>Ruta chalepensis</i> L.	Leaves	Chewed	Alkaloids, flavonoids, phenols, saponins.	X	N/A	[3, 136]
Rutaceae	<i>Vapris dainelli</i> (pichi-serm.) Kokwaro	Bark	Chewed	N/A	-	N/A	[3]
Rutaceae	<i>Zanthoxylum chalybeum</i> engl.	Bark	Hold	Alkaloids, phenolic	-	N/A	[3, 270]
Salvadoraceae	<i>Salvadora persica</i> L.	Stem	Brushed	Saponins, alkaloids	-	N/A	[3, 271]
Sapindaceae	<i>Dodonaea angustifolia</i>	Roots, leaves	Brushed	Di and terpenes, saponins, flavonoids, phenolic compound.	X	Antioxidant activity	[3, 272, 273]
Scrophulariaceae	<i>Verbascum sinaiticum</i> benth.	Roots	Chewed	Alkaloids, phenolic compounds, anthraquinones, flavonoids, saponins, tannins, steroids, terpenes.	-	Antioxidant activity	[169, 274]
Simaroubaceae	<i>Brucea antisynterrea</i> J.F. Mill.	Roots, bark	Chewed	Alkaloids, flavonoids, glycosides, phenols, quinones, saponins, steroids, tannins, terpenoids	X	N/A	[3, 275]
Solanaceae	<i>Solanum americanum</i> mill.	Leaves, fruits	Grind, boiled, chewed	Alkaloids, flavonoids, tannins, saponins	-	N/A	[140]
Solanaceae	<i>Datura stramonium</i> L.	Fruits, leaves, stem, seeds	Grind, rub, liquid form, boiled, squeezed	Alkaloids, tannins, flavonoids, phenols	X	Antioxidant activity	[152, 276, 277]
Solanaceae	<i>Nicotiana tabacum</i> L.	Leaves	Boiled, chewed	Flavonoids, alkaloids, tannins, phenolic compounds, terpenoids	-	N/A	[140, 278]
Solanaceae	<i>Solanum hastifolium</i> hochst.	Roots	Chewed	N/A	-	N/A	[169]
Solanaceae	<i>Solanum incanum</i> L.	Fruits, roots	Chewed	Alkaloids, flavonoids, saponins, steroids	-	Antioxidant activity	[3, 279]
Solanaceae	<i>Solanum marginatum</i> L. F.	Roots	Chewed	Alkaloids, saponins, flavonoids, glycosides, terpenoids, steroid	-	N/A	[3, 280]
Solanaceae	<i>Capsicum frutescens</i> L.	Fruits	Paste	Alkaloids, tannins, saponins, flavonoids, carotenoids, steroids	X	N/A	[103, 148]
Tiliaceae	<i>Grewia bicolor</i> juss	Stem	Brushed	Alkaloids, terpenes	-	N/A	[3, 281]
Tiliaceae	<i>Grewia ferruginea</i> hochst.	Roots	Shredded	N/A	-	N/A	[3]
Verbenaceae	<i>Premna schimpert</i> engl.	Leaves, root, stem	Grind, chewed, rub, boiled	Flavonoids, diterpenoids, terpenes, steroids, alkaloids	X	Antioxidant activity	[140, 282]
Verbenaceae	<i>Stachyurpelta angustifolia</i>	Leaves	Decoction	Phenolic, tannin compounds	X	Antioxidant activity	[148, 283]
Verbenaceae	<i>Premna oligorticha</i>	Leaves	Chewed	Flavonoids, alkaloids, terpenoids, resins, tannins, saponins and steroids,	-	N/A	[3, 169, 284]
Verbenaceae	<i>Premna resinosa</i> (hochst.)	Roots	Chewed	N/A	-	Antioxidant activity	[3, 285]
Vitaceae	<i>Cyphostemma junceum</i> (webb)	Whole plant	Chewed	Alkaloids, phenols, terpenoids, flavonoids	-	N/A	[169]
Vitaceae	<i>Cissus quadrangularis</i> L.	Roots	Chewed	Sterols, tannins, alkaloids	-	Antioxidant activity,	[3, 286]
Xanthorrhoeaceae	<i>Albizia vera</i> L.	Leaves	Chewed	Phenolic compounds, alkaloids	X	anti-inflammatory mechanism (COX and LOX)	[287-290]
Zingiberaceae	<i>Aframomum corrorima</i>	Seeds, rhizome	Chewed	Flavonoids, phenoles, terpenoids	-	Antioxidant activity	[3, 291]
Zingiberaceae	<i>Zingiber officinale</i> roseoe	Rhizome	Chewed	Flavonoids, alkaloids, saponins, steroids, terpenoids, tannin.	X	Antioxidant activity, Anti-inflammatory mechanism (COX and LOX)	[3, 60, 70, 292]

Table 4. Asia

Family	Scientific name	Used part	Preparation	Phytochemicals	Analgasic effect	Anti-inflammatory effect	Mechanism of action	Ref.
Acanthaceae	<i>Barleria Cristata</i>	Leaves, roots	Chewed, infusion	Terpenoids, flavonoids, phenolic compounds	-	x	Antioxidant activity	[293]
Acanthaceae	<i>Hygrophila auriculata</i>	Roots, leaves	N/A	N/A	-	-	N/A	[294]
Actinidiaceae	<i>Saurauia tristylia</i> DC.	Roots	Decoction	Phenolic compounds, flavonoids, alkaloids, steroids, saponins	-	-	N/A	[295, 296]
Amaryllidaceae	<i>Allium cepa</i> L.	Bulb	Raw, cooked	Phenolic acids, flavonoids, anthocyanins	X	x	Antioxidant activity, Anti-inflammatory mechanism	[6, 59, 297, 298]
Amaranthaceae	<i>Beta vulgaris</i> L.	Leaves	Juice	N/A	-	-	(COX and LOX)	[298, 299]
Amaranthaceae	<i>Achyranthes aspera</i>	Leaves, roots	Decoction	Saponins, alkaloids	X	x	Antioxidant activity	[14, 219, 294]
Anacardiaceae	<i>Mangifera indica</i> L.	Bark	Powdered inner bark, keep in mouth	N/A	-	-	N/A	[219]
Apocynaceae	<i>Wrightia tinctoria</i>	Leaves	Chewed	Flavonoids, glycoflavonoids, phenolic acids	-	x	Antioxidant activity	[300]
Apocynaceae	<i>Plumeria rubra</i> L.	Stem, bark	N/A	N/A	-	-	N/A	[294]
Apiaceae	<i>Panax ginseng</i>	Leaves, seeds	Extract	Flavonoids, terpenes	-	x	GABA mechanism	[110]
Asteraceae	<i>Asarum longirhizomatosum</i> C.	Whole plant	Ground, decoction	Flavonoids	-	x	N/A	[296, 301]
Asteraceae	<i>Asarum caudigerum</i>	Whole plant	Direct application	N/A	-	-	N/A	[302]
Asclepiadaceae	<i>Pergularia daemia</i>	Latex	Direct application	Alkaloids, saponins, tannins, flavonoids, terpenoids, phenols	-	x	Antioxidant activity	[219, 303]
Asteraceae	<i>Achillea heberstemii</i>	Leaves	N/A	N/A	-	-	N/A	[250, 298, 304]
Asteraceae	<i>Bidens biternate</i>	Roots	Paste	Glycosides, flavonoids, alkaloids, tannins, steroids, terpenoids, coumarins	-	-	N/A	[14, 305]
Asteraceae	<i>Parthenium hysterophorus</i> L.	Flowers	Chewed	Phenols, flavonoids	-	-	N/A	[3, 14, 306]
Asteraceae	<i>Achillea heberstemii</i>	Flowers, leaves	N/A	N/A	-	-	Antioxidant activity	[250, 304]
Asteraceae	<i>Arnica montana</i>	Flowers	Chewed	Terpenes, flavonoids, phenolic acids	X	x	GABA mechanism, antioxidant activity	[51, 51, 180]
Berberidaceae	<i>Berberis wuliangshanensis</i>	Roots	N/A	N/A	-	-	N/A	[307]
Berberidaceae	<i>Berberis multifida</i>	Leaves, stem	Extract	Alkaloids	X	x	N/A	[298, 308]
Bignoniaceae	<i>Incarvillea sinensis</i> Lam.	Roots (fresh or dried)	Gargle	Alkaloids	X	x	N/A	[309, 310]
Boraginaceae	<i>Cynoglossum lanceolatum</i>	Fruits, leaves, seeds	Chewed	Alkaloids	X	x	Antioxidant activity	[14, 311, 312]
Caesalpiniaceae	<i>Caesalpinia coriaria</i>	Fruits	Powder	Tannins	-	-	Antioxidant activity	[219, 313]
Capparidaceae	<i>Capparis spinosa</i>	Stem, fruits	N/A	Alkaloids, flavonoids, steroids, terpenoids, tocopherol, phenolic compounds	X	x	Antioxidant activity	[250, 298, 314, 315]
Caryophyllaceae	<i>Dianthus crinitus</i>	Seeds	N/A	Flavonoids, phenols	-	-	Antioxidant activity	[298]
Caryophyllaceae	<i>Dianthus orientalis</i>	Seed, leaves, flowers, fruits	Extract	N/A	-	-	Antioxidant activity	[250, 298]
Cannaceae	<i>Canna indica</i> L.	Roots	Direct application	N/A	-	-	N/A	[294]
Clemnaceae	<i>Cleome gynandra</i> L.	Leaves	Extract	Polyphenols, oleic acid, linoleic acid	X	-	Antioxidant activity	[219, 316]
Convolvulaceae	<i>Merremia chrysoides</i>	Whole plant	Decoction	N/A	-	-	N/A	[219]
Ebenaceae	<i>Diospyros lotus</i>	Branch, roots	Decoction	Phenolic compounds	X	x	Antioxidant activity	[14, 317]
Elaeagnaceae	<i>Elaeagnus umbellata</i>	Branch	Rub	Alkaloids, steroids, terpenoids, saponins	X	x	Antioxidant activity	[14, 318]
Erocaeaceae	<i>Erocaulon baergerianum</i>	Flowers	Decoction	Flavonoids, phenolic compounds	-	x	N/A	[302, 319]
Euphorbiaceae	<i>Jatropha gossypifolia</i>	Whole plant	Decoction	alkaloids, terpenes, flavonoids, phenolics	-	x	Antioxidant activity	[108]
Euphorbiaceae	<i>Jatropha caracas</i>	Sap	Topical	alkaloids, terpenes, flavonoids, phenolics	-	x	Antioxidant activity, anti-inflammatory mechanism (COX-LOX)	[108, 109, 294]
Fabaceae	<i>Acacia nilotica</i>	Bark	Decoction	Alkaloids, flavonoid phenols, tannins, terpenes	-	-	Antioxidant activity	[219-221]
Fabaceae	<i>Lespedeza juncea</i>	Whole plant	Decoction	Flavonoids, alkaloids	-	-	Antioxidant activity	[14, 320]
Fabaceae	<i>Senna tora</i> (L.)	Leaves, Stem, Fruit, Root, Seed	N/A	N/A	-	-	N/A	[294]
Geraniaceae	<i>Erodium cicutarium</i>	Aerial parts	Boiled	Polyphenols	-	-	Antioxidant activity	[298, 321]
Ginkgoaceae	<i>Ginkgo biloba</i>	Leaves, seeds	Extract	flavonoids, terpenes	-	x	N/A	[110, 322]
Hypericaceae	<i>Hypericum perforatum</i>	Flowers	N/A	Flavonoids	-	x	N/A	[298, 323]
Iridaceae	<i>Iris dichotoma</i> Pall.	Roots, bark	Chop and chew	Phenolic compound, flavonoids or isoflavones	-	x	N/A	[310, 324]
Lamiaceae	<i>Leucas aspera</i> (Willd.)	Leaves	Extract	Terpenoids, diterpenoids, phenolic compounds, alkaloids	-	-	Antioxidant activity	[219, 325]
Lamiaceae	<i>Teucrium polium</i> L.	Aerial parts	Extract	Flavonoids, terpenoids	-	x	Antioxidant activity	[298, 326]
Lamiaceae	<i>Stachys pilifera</i>	Leaves	Extract	Flavonoids, diterpenes, saponins, terpenoids	-	x	Antioxidant activity	[298, 327]
Lamiaceae	<i>Micromeria biflora</i>	Roots	Paste	N/A	-	-	N/A	[14]

Lamiaceae	Plectranthus	Bud, flowers	Chewed	Monoterpene, sesquiterpenoids, diterpenoids, phenolics	-	x	N/A	[14, 328]
Lauraceae	Lissea cubeba (Lour.) Pers.	Fruits	Decoction	Flavonoids, Terpenoids, Alkaloids	-	x	Antioxidant activity	[296, 329]
Liliaceae	Allium condensatum Turcz.	Aerial parts	N/A	N/A	-	-	N/A	[310]
Liliaceae	Fritillaria imperialis L.	Fruits	N/A	Alkaloids, terpenoids, saponins	x	x	Antioxidant activity	[298, 330]
Liliaceae	Allium tuberosum Rottl. ex	Whole plant	Ground, decoction	Alkaloids, flavonoids, terpenoids, tannins, saponins	-	-	Antioxidant activity	[296, 331]
Loganiaceae	Strychnos nux-vomica L.	Seeds	Chewed	Phenolic compounds	x	x	N/A	[302, 332]
Meliaceae	Azadirachta indica A.	Bark, leaves, branches	Powdered inner bark keep in mouth, extract	Alkaloids, flavonoids, terpenoids, phenolic compounds	-	-	Antioxidant activity	[219, 233, 235]
Menispermaceae	Cyclea hypoglauca (Schauer)	Roots	Infusion	N/A	-	-	N/A	[296]
Moraceae	Streblus asper, Lour.	Bark	Powdered inner bark keep in mouth	N/A	-	-	N/A	[219]
Myrtaceae	Syzygium aromaticum L.	Flowers	Decoction, powder and hydrodistillate	Monoterpene, sesquiterpenes, phenolics, hydrocarbon compounds.	x	x	Antioxidant activity, anti-inflammatory mechanism	[132, 288, 333]
Myrtaceae	Psidium guajava L.	Leaves, bark	Chewed	Flavonoids, terpenoids, tannins	x	x	Antioxidant activity (COX and LOX)	[93, 219, 246]
Fabaceae	Mimosa pudica L.	Roots	Boiled	N/A	-	-	N/A	[294]
Fabaceae	Hiptage benghalensis	Stem	N/A	N/A	-	-	N/A	[294]
Malpighiaceae	Mirabilis jalapa L. var.	Whole plant	N/A	Alkaloids, tannins, flavonoids, glycosides, phenols, steroids, terpenoids, saponins	-	X	Antioxidant activity	[307, 334]
Nyctaginaceae								
Oleaceae	Olea ferruginea (Sol.)	Fruits, leaves, seeds and bark	Decoction	Phenolic compounds	-	-	N/A	[335, 335, 336]
Oxalidaceae	Oxalis corniculata L.	Flowers	Chewed	Flavonoids, phyosterols, phenolic compounds, tannins	-	-	N/A	[14, 337]
Fabaceae	Astragalus verus Oliv.	Resin	N/A	N/A	-	-	N/A	[298]
Fabaceae	Clitoria terrata L.	Roots	Paste	N/A	-	-	N/A	[294]
Fabaceae	Pongamia pinnata L.	Seeds	Oil	N/A	-	-	N/A	[294]
Pinaceae	Pinus roxburghii Sarg.	Aerial parts	Extract, juice	Flavonoids	x	x	Antioxidant activity	[335, 338]
Plantaginaceae	Plantago lanceolata L.	Leaves	Grind and keep in mouth	Phenols, flavonoids	-	X	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[127, 335, 339]
Polypodiaceae	Drynaria roosei Nakaike	Rhizome	Cooked with water, sliced, dried, medicinal liquor	Flavonoids	-	-	N/A	[302]
Poaceae	Cynodon dactylon (L.)	Roots, leaves	N/A	N/A	-	-	N/A	[294]
Poaceae	Desmostachya bipinnata	Roots	N/A	N/A	-	-	N/A	[294]
Poaceae	Vetiveria zizanioides (L.)	Roots	N/A	N/A	-	-	N/A	[294]
Rhamnaceae	Rhamnus persica	Fruits, leaves	Decoction, infusion, powder	Monoterpene, sesquiterpenes, aldehydes, phenols	-	-	Antioxidant activity	[298]
Rosaceae	Sanguisorba minor	Stem, aerial parts, roots	N/A	Flavonoids, phenols, terpenoids	X	x	Antioxidant activity	[298, 340, 341]
Rosaceae	Prunus persica	Branch	Chewed	N/A	-	-	N/A	[14]
Rosaceae	Rubus parvifolius L.	Whole plant	Ground, decoction	Phenolic compounds	-	-	Antioxidant activity	[296, 342]
Rosaceae	Geum japonicum Thunb.	Whole plant	Decoction	Triterpenoids, sterols, tannins	-	x	Antioxidant activity	[302, 343]
Rosaceae	Amygdalus davidiana	Bark, leaves	N/A	N/A	-	-	N/A	[307]
Rosaceae	Prunus humilis Bunge	Roots (fresh or dried)	Chop and chew, decoction	Flavonoids and phenolic compounds	-	-	Antioxidant activity	[310, 344]
Rubiaceae	Gakona Aucheri	Leaves, flowers	Decoction	N/A	-	-	N/A	[298]
Rubiaceae	Zanthoxylum nitidum	Stem, branches, leaves, roots	Decoction	Alkaloids, coumarins, lignans, flavonoids, terpenes, steroids, alkylamides	x	x	Antioxidant activity	[345]
Rubiaceae	Zanthoxylum bungeanum	Stem, leaves, seeds, roots, bark	Liquid, powder	Alkaloids, flavonoids, terpenoids	x	x	Antioxidant activity	[307, 346]
Rubiaceae	Zanthoxylum alatum	Branch	Rub. dust	Linolol	-	x	Antioxidant activity	[14, 347]
Rubiaceae	Murraya exotica L.	Roots, leaves	Ground, decoction	Alkaloids, tannins, alkalis	-	x	Antioxidant activity	[296, 348]
Solanaceae	Solanum melongena L.	Roots, stem	Direct application	Flavonoids, terpenes, glycoalkaloids	x	x	Antioxidant activity	[302, 349]
Solanaceae	Solanum verbascifolium L.	Whole plant	Decoction	Flavonoids	-	-	N/A	[302, 350]
Solanaceae	Solanum nigrum, L.	Leaves	Extract	N/A	-	-	Antioxidant activity	[219, 351]
Solanaceae	Solanum surattense.	Seeds	Stem from the seeds keep in the mouth	Alkaloids	-	-	N/A	[219, 352]
Solanaceae	Hyoscyamus niger L.	Leaves, seeds	Infusion, gargle	Alkaloids, atropine, tropane and scopolamine.	x	x	Antioxidant activity	[123, 298, 353]
Solanaceae	Lycopersicon esculentum	Fruits, leaves	N/A	Flavonoids, phenols	-	-	Antioxidant activity	[298, 354]



Solanaceae	Hyoisyanum muticus	and flowers	N/A	Alkaloids	-	N/A	[298]
Thymelaeaceae	Daphne mucronata	Seeds, roots	N/A	Coumarins, flavonoids, triterpenoids	-	N/A	[298, 355]
Verbenaceae	Vriex negundo, L.	Leaves, stem	Chewed	Flavonoids	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[219, 356]
Vitaceae	Cayratia albifolia C. L. Li	Root, leaves	Chewed	N/A	-	N/A	[302]
Xanthorrhoeaceae	Albe vera L.	Leaves	Decoction	Lignin, anthraquinones, saponins	x	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[287, 288, 290]
Zygophyllaceae	Tribulus terrestris, L.	Fruit	Powder, massage	Alkaloids, tannins, saponins	x	N/A	[219, 357]
Zygophyllaceae	Peganum harmala L.	Seeds, fruit	Infuse	Flavonoids, alkaloids, saponins, tannins, glycosides, terpenoids and steroids	x	N/A	[249-251, 298]
Zygophyllaceae	Fagonia olivieri	Roots, fruits, stem, flowers	N/A	Saponins, terpenoids	-	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[358-360]
Asteraceae	Emilia sonchifolia (L.)	Flowers	Paste	Alkaloids	x	N/A	[93, 361]
Asteraceae	Galinsega parviflora Cav.	Leaves	Decoction	Polypheonols and flavonoids, phenolic acids	-	Antioxidant activity, anti-inflammatory mechanism (COX and LOX)	[93, 183]
Asteraceae	Xanthium strumarium L.	Leaves, roots	Juice	Sesquiterpenic lactones, glycosides, phenols, polyesters	x	Antioxidant activity	[93, 362]
Fabaceae	Mimosa pudica	Seeds	Powder	Alkaloids, flavonoids, glycosides, sterols, terpenoids, tannins and fatty acids	x	Antioxidant activity	[93, 363]
Myrtaceae	Myrcia adenophora	Bark	Decoction	Flavonoids, triterpenoids, tannins, monoterpenoids, and benzenoids	x	Antioxidant activity	[93, 364]
Plantaginaceae	Plantago major L.	Flowers	Paste	Polysaccharides, flavonoids, glycosides, terpenoids, alkaloids, organic acids	x	Antioxidant activity	[93, 341]
Rubiaceae	Spermacoe neohispida Govaerts	Flowers	Paste	N/A	-	N/A	[93]
Rutaceae	Zanthoxylum oxyphyllum	Leaves, fruits	Juice	N/A	-	N/A	[93]
Sapindaceae	Sapindus mukorossi Gaert	Seeds	Powder	Saponins	-	N/A	[93, 365]
Solanaceae	Nicotiana plumbaginifolia Viv	Leaves	Paste	Alkaloids, saponin, tannin, flavonoids, phenolic compounds, steroids, terpenoids and carbohydrates.	-	N/A	[93, 366]
Solanaceae	Solanum aculeatissimum Jacq	Leaves, Fruits	Decoction	Steroidal saponins	-	N/A	[93, 294, 367]
Solanaceae	Solanum virginianum Dunal	Leaves, seeds	N/A	N/A	-	N/A	[294]
Urticaceae	Urtica parviflora Roxb	Fruits	Boiled juice	Alkaloids, polysaccharides, saponins, flavonoids, phenolic compounds, glycosides and tannins	-	Antioxidant activity	[93, 368]
Arecaceae	Borrassia flabellifer	Roots	Boiled, gargling	N/A	-	N/A	[294]
Verbenaceae	Vriex negundo L.	Leaves, bark	N/A	N/A	-	N/A	[294]

Table 5. Ocenmia

Family	Scientific name	Used part	Preparation	Phytochemicals	Analgesic effect	Anti-inflammatory effect	Mechanism of action	Ref.
Euphorbiaceae	<i>Petalostigma pubescens</i>	Fruits, bark	Hold in the mouth	Saponins, phenolic compounds, flavonoids, triterpenoids, tannins and alkaloids	-	-	Antioxidant activity	[90, 369]
Euphorbiaceae	<i>Petalostigma quadrioculare</i>	Fruits, bark	N/A	Saponins, phenolic compounds, flavonoids, triterpenoids, tannins and alkaloids	-	-	Antioxidant activity	[90, 369]
Myoporaceae	<i>Eremophila fraseri</i>	Leaves	N/A	Diterpenes, triterpenoids, phenylpropanoids, lignans, flavonoids	-	x	Antioxidant activity	[90, 370]
Myrtaceae	<i>Melaleuca cajuputi</i>	Bark, leaves	Extract	Flavonoids, phenolic compounds	-	-	Antioxidant activity	[90, 371]
Rhamnaceae	<i>Ventilago viminalis</i>	Bark, roots	N/A	N/A	-	-	N/A	[90]
Rutaceae	<i>Geijera parviflora</i>	Leaves	N/A	Alkaloids, flavonoids	-	x	N/A	[90, 372]
Rutaceae	<i>Melicope vitiflora</i>	Bark	N/A	N/A	-	-	N/A	[90]
Sapindaceae	<i>Dodonaea viscosa</i>	Roots, leaves	N/A	Alkaloids, flavonoids, steroids, phenolics, saponins, tannins	x	x	Antioxidant activity	[90, 373]
Tiliaceae	<i>Grewia retusifolia</i>	Fruit, roots, leaves	N/A	N/A	-	-	N/A	[90]
Asteraceae	<i>Acmella grandiflora</i>	Roots, flowers	Crushed	Spilanthol, acemollonate, tannins, flavonoids, and phenolic compounds	x	X	Antioxidant activity, GABA mechanism	[90, 374]

method to increase its action.

The parts used may vary with the medicinal plant; however, we found that the most commonly used plant parts were leaves and roots, followed by the bark, stem, and seeds (Fig. 1). Preferentially using the leaves instead of roots can also prevent detrimental effects on the plants.

The most common phytochemicals involved in the mechanism of action of medicinal plants for toothache treatment are polyphenols, more specifically, flavonoids and terpenes, which are the most abundant secondary metabolites and antioxidants in the human diet<sup>3,5-7</sup>. Flavonoids are the most ubiquitous group of all plant phenolics, which could explain the implicit antioxidant capacity of all medicinal plants<sup>84</sup>. Furthermore, flavonoids can modulate the function of ionotropic GABA receptors, suggesting that these phytochemicals can exert different mechanisms of action to relieve pain<sup>20</sup>.

Polyphenols are strong antioxidants that neutralize free radicals by donating an electron or a hydrogen atom<sup>61</sup>, thus exerting antioxidant effects in plants and organisms that consume them. However, polyphenols decrease the concentrations of ROS and RNS far from the site of the primary response because the local concentrations of these radicals around the inflammatory site are substantially high (> 1 mM). Therefore, polyphenols are highly unlikely to be effective where these free radicals are produced but could be quantitatively more effective as antioxidants in the surrounding unaffected tissues<sup>85</sup>.

Substances such as capsaicin, allicin, camphor, and menthol cause a state of activation and desensitization in the TRP receptor pathway through which pain may be reduced<sup>18, 43</sup>. Of these phytochemicals, only allicin was analyzed in the studies included in this review. Moreover, allicin and spilanthol, compounds present in *A. caulirhiza*, have various biological and pharmacological effects, which may cause analgesia<sup>3, 52</sup>. Further studies should be performed to better understand the roles of these phytochemicals.

The mechanisms of action of phytochemicals in toothache relief are only partly understood. This review shows that these mechanisms involve antioxidant activity, action on TRP receptors, GABA mechanism, and COX/LOX inhibitory activity. The tables in this review outline

163 medicinal plants with antioxidant mechanisms of action, 20 with an anti-inflammatory mechanism (COX/LOX), four with GABA mechanism, and two with TRP mechanism. Some plants have two reported mechanisms of action. However, in general, there is insufficient literature addressing each mechanism responsible for toothache relief.

Several reports cite the use of various plants for toothache treatment; however, in many of these reports, the mechanism of action underlying pain relief is not specified, as indicated in the tables in this review. Moreover, in several studies, the phytochemicals potentially responsible for the analgesic effect were not reported. Accordingly, future studies should focus on identifying the exact mechanisms that contribute to dental analgesia and the phytochemicals involved. Additionally, the “common names” of medicinal plants were not included in this research, considering the extensive information involved in the preparation of this manuscript.

As mentioned above, we found only one study comparing the pharmacological properties of medicinal plants with those of conventional pharmaceutical drugs<sup>81</sup>. However, 30 plants with dual anti-inflammatory mechanisms (COX/LOX) were identified, as outlined in the tables for each continent. This information could be useful in future comparative studies of conventional or dual NSAIDs.

Although dual inhibition of microsomal PGE2 synthetase (mPGES-1) and 5-LOX has not been described as a mechanism of action in the reports included in this review, several plants (some of which are indicated in our tables) contain acylphloroglucinols, phenolic compounds, and non-phenolic acidic structures that exhibit such dual action<sup>86</sup>. mPGES-1 is an inducible enzyme at inflammatory sites that preferentially receives its substrate from co-induced COX-2 and is responsible for the excessive formation of PGE2 during acute and chronic inflammation [87, 88.; thus, its inhibition could be a promising strategy for toothache treatment with medicinal plants. Furthermore, natural mPGES-1 inhibitors have advantages over NSAIDs since they are non-synthetic and safer because they do not inhibit COX-derived homeostatic eicosanoids<sup>86</sup>.

However, in several *in vitro* and *in vivo* studies (utilizing indices such as IC<sub>50</sub>), the vast

majority of medicinal plant extracts have been tested only against one biological target (COX or LOX), which is insufficient to validate their anti-inflammatory and analgesic properties and hinders direct comparisons between plants and conventional or dual-action NSAIDs<sup>72</sup>. Therefore, further studies should be conducted to address this gap in research and gather more relevant information.

Multiple experimental studies of COX-1/COX-2 inhibition have used IC<sub>50</sub> (the concentration at which an NSAID produces 50% inhibition of both COX enzymes) to rank the relative inhibitory activity of NSAIDs on these enzymes, and consequently, establish their selectivity over COX, correlating this *in vitro* inhibition with clinical efficacy and toxicity levels<sup>89</sup>. However, IC<sub>50</sub> values do not indicate the mechanism of enzyme inhibition and vary with substrate concentration. Furthermore, these values are not directly comparable unless identical experimental conditions are used, and they must be analyzed carefully when inhibition is time-dependent<sup>89</sup>. These drawbacks also (91) hinder direct comparison between medicinal plants and NSAIDs.

Only two studies on plant combinations were found for this review. The study on phytotherapeutic Lenidase®<sup>81</sup>. was already described above. The other reported on the use of seven popular medicinal plant mixtures for toothache in Europe (Catalonia), including several species<sup>76</sup>. whose use was not found in other continents.

Although medicinal plants are distributed throughout the world<sup>90</sup>, biodiversity could affect how intensely such plants are used for toothache, and thus the discovery of new drugs<sup>91</sup>. The 17 most megadiverse countries in the world are Brazil, Colombia, Mexico, Peru, Ecuador, Venezuela, the United States of America, Indonesia, Australia, Madagascar, China, the Philippines, India, New Guinea, Malaysia, South Africa, and the Democratic Republic of Congo; most of these are in the American continent<sup>49</sup>. However, in this present review, most of the information on medicinal plants was gathered from Asia (Table 4) and Africa (Table 5), possibly because herbal medicines remain a key component of healthcare systems in the developing cultures of these continents<sup>90</sup>. Oceania has only two of the 17 megadiverse countries, which may

explain the scarcity of plants in this continent (Table 5). Nevertheless, in some regions, much of the traditional knowledge about medicinal plants is only spread verbally and, thus, remains unexplored and unreported<sup>93</sup>.

In terms of plants used in different continents, *A. sativum* was found in America (Table 1), Europe (Table 2), and Africa (Table 3); *A. cepa* and *Syzygium aromaticum* were found in Europe, Africa, and Asia; and *Acmella oleracea*, *Jatropha curcas*, and *Jatropha gossypifolia* were all found in America (Table 1), Africa (Table 3), and Asia (Table 4). Conversely, some species are found only in one continent, such as *Thymus schimperii* Ronniger in Africa, perhaps because this species is a rare plant highly localized in and endemic to Ethiopia<sup>94</sup>.

Among the families of medicinal plants used worldwide, Asteraceae was the most common, followed by Solanaceae, Fabaceae, Lamiaceae, Euphorbiaceae, Rutaceae, and Myrtaceae (Fig. 2). The first three of these families have been widely reported as those most commonly used to treat inflammation and various types of pain<sup>44, 44, 76</sup>.

Notably, the present review discusses several medicinal plants for toothache treatment, which have been globally classified by continents, unlike all the referenced studies, which were approached separately. Moreover, the five tables provide details about the parts used, preparation, phytochemicals, analgesic/anti-inflammatory effect, and mechanisms of action, contrary to most studies that do not include such details. Additionally, this review analyzes all the mechanisms of action of the medicinal plants that have been ascribed until now for toothache treatment, unlike many studies that only cite these mechanisms. Furthermore, we have included a section comparing medicinal plants and pharmaceutical drugs, unlike all the referenced studies that do not provide such a comparison. Finally, this review discusses the use of medicinal plants for the treatment of dental pain, while most articles deal with this topic on a general basis.

#### **Future perspectives**

Although phytotherapy has a long history, natural medicines are considered a hidden source of drugs because many medicinal plants have not been studied in depth<sup>79</sup>. Accordingly, further studies should be conducted to better understand the role and benefits of phytotherapeutic drugs<sup>81</sup>. for

toothache treatment, particularly when combined based on the multi-objective therapeutic principle of phytotherapy<sup>79, 95</sup>. This principle would be analogous to the multimodal analgesic approach used for NSAIDs<sup>96-98</sup>.

Aromatherapy also has non-pharmacological therapeutic potential for reducing toothache by combining highly complex mixtures of essential oils to produce a therapeutic effect<sup>95</sup>. Therefore, further research should be conducted in this field of alternative medicine.

Although polyphenols in organic food extracts (extractable polyphenols) have already been analyzed, significant amounts of potentially bioactive polyphenols that remain in the residues (non-extractable polyphenols) have been overlooked. Additionally, significant amounts of non-extractable polyphenols are found in foods and vegetables<sup>61, 99</sup>. Therefore, these compounds should be considered for future studies.

A promising therapeutic option for the administration of flavonoids that may increase their bioavailability is to develop protective systems, such as microcapsules, nanoparticles, and nano-formulations, which improve water solubility, dissolution, absorption, and thermal stability. Accordingly, in the near future, such systems should be developed and administered for pain management<sup>62</sup>.

Finally, human clinical trials are essential to confirm the effectiveness of traditional phytotherapy for toothache and to investigate the pharmacodynamic and pharmacokinetic interactions between medicinal plants and other synthetic drugs. Similarly, predictive (*in silico*) models, phytochemical analyses, and ethnopharmacological studies could be milestones for drug discovery in traditional medicinal plants for toothache treatment, because many of them lack information or have not been studied.

## **CONCLUSION**

This is the first review to compile a large volume of data on the global use of medicinal plants for the treatment of toothache. A total of 21 species of medicinal plants were found in America (Table 1), 29 in Europe (Table 2), 192 in Africa (Table 3), 112 in Asia (Table 4), and 10 in Oceania (Table 5). Asia and Africa are the continents where

the most research has been done on this topic. Asteraceae was the most commonly found plant family in this review, followed by Solanaceae, Fabaceae, Lamiaceae, Euphorbiaceae, Rutaceae, and Myrtaceae.

In total, 364 medicinal plants used for toothache treatment were identified, of which 139 have not yet been scientifically studied, highlighting opportunities for ethnopharmacological research on toothache treatments. The most common species were *A. sativum*, *A. cepa*, *A. oleracea*, *J. curcas*, *J. gossypifolia*, and *S. aromaticum*. These families and species were more commonly found in Africa and Asia, corroborating our previously reported findings. As determined in this review, the most commonly used plant parts were the leaves and roots, followed by the bark, stems, and seeds.

We identified four mechanisms of action of medicinal plants implied in toothache treatment, namely, the antioxidant effect, effects mediated through TRP receptors, GABA mechanism, and the anti-inflammatory mechanism (COX/LOX). Flavonoids, terpenes, polyphenols, and alkaloids are the phytochemicals most commonly associated with toothache treatment. Many of the plants analyzed in this review have the potential to be used as agents for toothache treatment. Therefore, future studies must prioritize the analysis of their pharmacodynamic and pharmacokinetic interactions.

Finally, to more precisely clarify the usefulness of medicinal plants as a valid option for toothache treatment, comparative studies between medicinal plants and commonly used pharmaceutical drugs should be conducted. In addition, studies published in Spanish should be included in future reviews since we only analyzed studies published in English, and this may have limited our ability to gather additional information.

#### ACKNOWLEDGMENTS

None.

#### Ethical Statement

Ethical approval was not required for this review article.

#### Conflicts of Interest

The authors declare no conflicts of interest.

#### REFERENCES

1. Heir GM. The Emerging Specialty of Orofacial Pain. *J Indian Prosthodont Soc* 2013; 13: 140–141.
2. Cohen LA, Bonito AJ, Akin DR, et al. Toothache pain: behavioral impact and self-care strategies. *Spec Care Dent* 2009; 29: 85–95.
3. Megersa M, Jima TT, Goro KK. The Use of Medicinal Plants for the Treatment of Toothache in Ethiopia. *Evid Based Complement Alternat Med* 2019; 2019: 2645174.
4. Palombo EA. Traditional Medicinal Plant Extracts and Natural Products with Activity against Oral Bacteria: Potential Application in the Prevention and Treatment of Oral Diseases. *Evid Based Complement Alternat Med* 2011; 2011: 680354.
5. Sharifi Rad M, Yýlmaz YB, Antika G, et al. Phytochemical constituents, biological activities, and health-promoting effects of the genus *Origanum*. *Phytother Res* 2021; 35: 95–121.
6. Ilic DV, Radicevic BA, Nedelcheva A, et al. Traditional dentistry knowledge among Serbs in several Balkan countries. *J Intercult Ethnopharmacol* 2017; 6: 223–233.
7. Yadav R, Agarwala M. Phytochemical analysis of some medicinal plants. *J Phytol* 2011; 3: 10–14.
8. Falzon CC, Balabanova A. Phytotherapy: An Introduction to Herbal Medicine. *Prim Care* 2017; 44: 217–227.
9. Zhang L, Virgous C, Si H. Synergistic anti-inflammatory effects and mechanisms of combined phytochemicals. *J Nutr Biochem* 2019; 69: 19–30.
10. B.K. Tiwari, Nigel P. Bruton, Charles S. Brennan. *Handbook of Plant Food Phytochemicals: Sources, Stability and Extraction*. 2013.
11. Prakash Bhanu. *Functional and Preservative Properties of Phytochemicals*. 1st ed. Elsevier, 2020.
12. Mahmood A, Mahmood A, Malik RN, et al. Indigenous knowledge of medicinal plants from Gujranwala district, Pakistan. *J Ethnopharmacol* 2013; 148: 714–723.
13. Kunwar RM, Nepal BK, Kshhetri HB, et al. Ethnomedicine in Himalaya: a case study from Dolpa, Humla, Jumla and Mustang districts of Nepal. *J Ethnobiol Ethnomed* 2006; 2: 27.
14. Farooq A, Amjad MS, Ahmad K, et al. Ethnomedicinal knowledge of the rural communities of Dhirkot, Azad Jammu and Kashmir, Pakistan. *J Ethnobiol Ethnomed* 2019; 15: 45.
15. Luna-Vázquez FJ, Ibarra-Alvarado C, Rojas-Molina A, et al. Vasodilator compounds derived

- from plants and their mechanisms of action. *Molecules* 2013; 18: 5814–5857.
16. Liu RH. Potential synergy of phytochemicals in cancer prevention: mechanism of action. *J Nutr* 2004; 134: 3479S-3485S.
  17. Benzie IF, Choi SW. Antioxidants in food: content, measurement, significance, action, cautions, caveats, and research needs. *Adv Food Nutr Res* 2014; 71: 1–53.
  18. Premkumar LS. Transient receptor potential channels as targets for phytochemicals. *ACS Chem Neurosci* 2014; 5: 1117–1130.
  19. Johnston GAR, Hanrahan JR, Chebib M, et al. Modulation of ionotropic GABA receptors by natural products of plant origin. *Adv Pharmacol* 2006; 54: 285–316.
  20. Hanrahan JR, Chebib M, Johnston GAR. Interactions of flavonoids with ionotropic GABA receptors. *Adv Pharmacol* 2015; 72: 189–200.
  21. Tasneem S, Liu B, Li B, et al. Molecular pharmacology of inflammation: Medicinal plants as anti-inflammatory agents. *Pharmacol Res* 2019; 139: 126–140.
  22. Iseghohi SO. A Review of the Uses and Medicinal Properties of *Dennettia tripetala* (Pepperfruit). *Med Sci Basel* 2015; 3: 104–111.
  23. Nilius B, Flockerzi V. *Mammalian Transient Receptor Potential (TRP) Cation Channels*. Handb Exp Pharmacol, 2014. Epub ahead of print 2014. DOI: 10.1007/978-3-319-05161-1.
  24. Arulselvan P, Fard MT, Tan WS, et al. Role of Antioxidants and Natural Products in Inflammation. *Oxid Med Cell Longev* 2016; 2016: 5276130.
  25. Becker EM, Nissen LR, Skibsted LH. Antioxidant evaluation protocols: Food quality or health effects. *Eur Food Res Technol* 2004; 219: 561–571.
  26. Tobore TO. Towards a Comprehensive Theory of Non-Cancer Acute and Chronic Pain Management: The Critical Role of Reactive Oxygen and Nitrogen Species in Pain, and Opioid Dependence, Addiction, Hyperalgesia, and Tolerance. *Adv Redox Res* 2021; 2: 100003.
  27. Thannickal VJ, Fanburg BL. Reactive oxygen species in cell signaling. *Am J Physiol Lung Cell Mol Physiol* 2000; 279: L1005–L1028.
  28. Federico A, Morgillo F, Tuccillo C, et al. Chronic inflammation and oxidative stress in human carcinogenesis. *Int J Cancer* 2007; 121: 2381–2386.
  29. Hussain SP, Hofseth LJ, Harris CC. Radical causes of cancer. *Nat Rev Cancer* 2003; 3: 276–285.
  30. Tang SY, Halliwell B. Medicinal plants and antioxidants: what do we learn from cell culture and *Caenorhabditis elegans* studies? *Biochem Biophys Res Commun* 2010; 394: 1–5.
  31. Chanda S, Dave R. In vitro models for antioxidant activity evaluation and some medicinal plants possessing antioxidant properties: An overview. *Afr J Microbiol Res* 2009; 3: 981–996.
  32. Kasote DM, Katyare SS, Hegde MV, et al. Significance of Antioxidant Potential of Plants and its Relevance to Therapeutic Applications. *Int J Biol Sci* 2015; 11: 982–991.
  33. Little JW, Doyle T, Salvemini D. Reactive nitroxidative species and nociceptive processing: determining the roles for nitric oxide, superoxide, and peroxynitrite in pain. *Amino Acids* 2012; 42: 75–94.
  34. Grace PM, Gaudet AD, Staikopoulos V, et al. Nitroxidative Signaling Mechanisms in Pathological Pain. *Trends Neurosci* 2016; 39: 862–879.
  35. Salvemini D, Little JW, Doyle T, et al. Roles of reactive oxygen and nitrogen species in pain. *Free Radic Biol Med* 2011; 51: 951–966.
  36. Yeo JF, Ling SF, Tang N, et al. Antinociceptive effect of CNS peroxynitrite scavenger in a mouse model of orofacial pain. *Exp Brain Res* 2008; 184: 435–438.
  37. Bruno R, Ghiadoni L. Polyphenols, Antioxidants and the Sympathetic Nervous System. *Curr Pharm Des* 2018; 24: 130–139.
  38. Rokyta R, Holecek V, Pekárkova I, et al. Free radicals after painful stimulation are influenced by antioxidants and analgesics. *Neuro Endocrinol Lett* 2003; 24: 304–309.
  39. Boutros T, Chevet E, Metrakos P. Mitogen-Activated Protein (MAP) Kinase/MAP Kinase Phosphatase Regulation: Roles in Cell Growth, Death, and Cancer. *Pharmacol Rev* 2008; 60: 261–310.
  40. Tominaga M, Wada M, Masu M. Potentiation of capsaicin receptor activity by metabotropic ATP receptors as a possible mechanism for ATP-evoked pain and hyperalgesia. *Proc Natl Acad Sci U S A* 2001; 98: 6951–6956.
  41. Ruparel NB, Patwardhan AM, Akopian AN, et al. Homologous and heterologous desensitization of capsaicin and mustard oil responses utilize different cellular pathways in nociceptors. *Pain* 2008; 135: 271–279.
  42. Bautista DM, Movahed P, Hinman A, et al. Pungent products from garlic activate the sensory ion channel TRPA1. *Proc Natl Acad Sci U S A* 2005; 102: 12248–12252.
  43. Hossain MZ, Bakri MM, Yahya F, et al. The Role of Transient Receptor Potential (TRP) Channels in the Transduction of Dental Pain. *Int J Mol Sci* 2019; 20: 526.

44. Uritu CM, Mihai CT, Stanciu GD, et al. Medicinal Plants of the Family Lamiaceae in Pain Therapy: A Review. *Pain Res Manag* 2018; 2018: 7801543.
45. Goudet C, Magnaghi V, Landry M, et al. Metabotropic receptors for glutamate and GABA in pain. *Brain Res Rev* 2009; 60: 43–56.
46. Enna SJ, McCarson KE. The Role of GABA in the Mediation and Perception of Pain. *Adv Pharmacol* 2006; 54: 1–27.
47. Jasmin L, Rabkin SD, Granato A, et al. Analgesia and hyperalgesia from GABA-mediated modulation of the cerebral cortex. *Nature* 2003; 424: 316–320.
48. Dubey S, Maity S, Singh M, et al. Phytochemistry, Pharmacology and Toxicology of *Spilanthes acmella*: A Review. *Adv Pharmacol Sci* 2013; 2013: 423750.
49. Prachayasittikul V, Prachayasittikul S, Ruchirawat S, et al. High therapeutic potential of *Spilanthes acmella*: A review. *EXCLI J* 2013; 12: 291–312.
50. Abdul Rahim R, Jayusman PA, Muhammad N, et al. Potential Antioxidant and Anti-Inflammatory Effects of *Spilanthes acmella* and Its Health Beneficial Effects: A Review. *Int J Environ Res Public Health* 2021; 18: 3532.
51. Rondanelli M, Fossari F, Vecchio V, et al. *Acmella oleracea* for pain management. *Fitoterapia* 2020; 140: 104419.
52. Barbosa AF, Carvalho MG de, Smith RE, et al. *Spilanthes*: occurrence, extraction, chemistry and biological activities. *Rev Bras Farmacogn* 2016; 26: 128–133.
53. Julémont F, Dogné JM, Pirotte B, et al. Recent development in the field of dual COX / 5-LOX inhibitors. *Mini Rev Med Chem* 2004; 4: 633–638.
54. Yahfoufi N, Alsadi N, Jambi M, et al. The Immunomodulatory and Anti-Inflammatory Role of Polyphenols. *Nutrients* 2018; 10: 1618.
55. Martel-Pelletier J, Lajeunesse D, Reboul P, et al. Therapeutic role of dual inhibitors of 5-LOX and COX, selective and non-selective non-steroidal anti-inflammatory drugs. *Ann Rheum Dis* 2003; 62: 501–509.
56. Leval X d, Julemont F, Delarge J, et al. New trends in dual 5-LOX/COX inhibition. *Curr Med Chem* 2002; 9: 941–962.
57. Rao CV. Regulation of COX and LOX by curcumin. *Adv Exp Med Biol* 2007; 595: 213–226.
58. Marefati N, Ghorani V, Shakeri F, et al. A review of anti-inflammatory, antioxidant, and immunomodulatory effects of *Allium cepa* and its main constituents. *Pharm Biol* 2021; 59: 287–302.
59. Wilson EA, Demmig Adams B. Antioxidant, anti inflammatory, and antimicrobial properties of garlic and onions. *Nutr Food Sci* 2007; 37: 178–183.
60. Rondanelli M, Fossari F, Vecchio V, et al. Clinical trials on pain lowering effect of ginger: A narrative review. *Phytother Res* 2020; 34: 2843–2856.
61. Belšèak-Cvitanoviæ A, Durgo K, Huđek A, et al. Overview of polyphenols and their properties. In: Galanakis CM (ed) *Polyphenols: Properties, Recovery, and Applications*. Woodhead Publishing, 2018, pp. 3–44.
62. Ferraz CR, Carvalho TT, Manchope MF, et al. Therapeutic Potential of Flavonoids in Pain and Inflammation: Mechanisms of Action, Pre-Clinical and Clinical Data, and Pharmaceutical Development. *Molecules* 2020; 25: 762.
63. Rein MJ, Renouf M, Cruz Hernandez C, et al. Bioavailability of bioactive food compounds: a challenging journey to bioefficacy. *Br J Clin Pharmacol* 2013; 75: 588–602.
64. Lim W, Mudge KW, Vermeylen F. Effects of population, age, and cultivation methods on ginsenoside content of wild American ginseng (*Panax quinquefolium*). *J Agric Food Chem* 2005; 53: 8498–8505.
65. Platel K, Srinivasan K. Bioavailability of Micronutrients from Plant Foods: An Update. *Crit Rev Food Sci Nutr* 2016; 56: 1608–1619.
66. Sun X, Chen W, Dai W, et al. *Piper sarmentosum* Roxb.: A review on its botany, traditional uses, phytochemistry, and pharmacological activities. *J Ethnopharmacol* 2020; 263: 112897.
67. Hewlings SJ, Kalman DS. Curcumin: A Review of Its Effects on Human Health. *Foods* 2017; 6: 92.
68. Rayati F, Hajmanouchehri F, Najafi E. Comparison of anti-inflammatory and analgesic effects of Ginger powder and Ibuprofen in postsurgical pain model: A randomized, double-blind, case-control clinical trial. *Dent Res J Isfahan* 2017; 14: 1–7.
69. Grzanna R, Lindmark L, Frondoza CG. Ginger—an herbal medicinal product with broad anti-inflammatory actions. *J Med Food* 2005; 8: 125–132.
70. Ali BH, Blunden G, Tanira MO, et al. Some phytochemical, pharmacological and toxicological properties of ginger (*Zingiber officinale* Roscoe): a review of recent research. *Food Chem Toxicol* 2008; 46: 409–420.
71. Chrubasik S, Pittler MH, Roufogalis BD. *Zingiberis rhizoma*: a comprehensive review on the ginger effect and efficacy profiles. *Phytomedicine* 2005; 12: 684–701.

72. Khumalo GP, Van Wyk BE, Feng Y, et al. A review of the traditional use of southern African medicinal plants for the treatment of inflammation and inflammatory pain. *J Ethnopharmacol* 2022; 283: 114436.
73. Mao QQ, Xu XY, Cao SY, et al. Bioactive Compounds and Bioactivities of Ginger (*Zingiber officinale* Roscoe). *Foods* 2019; 8: 185.
74. Tjendraputra E, Tran VH, Liu-Brennan D, et al. Effect of ginger constituents and synthetic analogues on cyclooxygenase-2 enzyme in intact cells. *Bioorg Chem* 2001; 29: 156–163.
75. Flynn DL, Rafferty MF, Boctor AM. Inhibition of human neutrophil 5-lipoxygenase activity by gingerdione, shogaol, capsaicin and related pungent compounds. *Prostaglandins Leukot Med* 1986; 24: 195–198.
76. Gras A, Parada M, Rigat M, et al. Folk medicinal plant mixtures: Establishing a protocol for further studies. *J Ethnopharmacol* 2018; 214: 244–273.
77. Che CT, Wang ZJ, Chow MSS, et al. Herb-Herb Combination for Therapeutic Enhancement and Advancement: Theory, Practice and Future Perspectives. *Molecules* 2013; 18: 5125–5141.
78. Vijayalakshmi G, Adinarayana M, Jayaprakash Rao P. A synergistic approach to kinetic and mechanistic studies of regeneration of  $\beta$ -carotene from tert-butoxyl radical induced  $\beta$ -carotene radical cation by chlorogenic acid. *Int J Pharm Life Sci* 2014; 5: 942–950.
79. Rather MA, Bhat BA, Qurishi MA. Multicomponent phytotherapeutic approach gaining momentum: Is the ‘one drug to fit all’ model breaking down?. *Phytomedicine* 2013; 21: 1–14.
80. Allescher HD. Functional dyspepsia—a multicausal disease and its therapy. *Phytomedicine* 2006; 13 Suppl 5: 2–11.
81. Isola G, Matarese M, Ramaglia L, et al. Efficacy of a drug composed of herbal extracts on postoperative discomfort after surgical removal of impacted mandibular third molar: a randomized, triple-blind, controlled clinical trial. *Clin Oral Investig* 2019; 23: 2443–2453.
82. Szyzkowska A, Koper J, Szczerba J, et al. The use of medicinal plants in dental treatment. *Herba Pol* 2010; 56: 97–107.
83. Cruz Martínez C, Diaz Gómez M, Oh MS. Use of traditional herbal medicine as an alternative in dental treatment in Mexican dentistry: a review. *Pharm Biol* 2017; 55: 1992–1998.
84. Campos-Vega R, Oomah BD. Chemistry and classification of phytochemicals. In: Tiwari BK, Brunton NP, Brennan CS (eds) *Handbook of Plant Food Phytochemicals*. John Wiley & Sons Ltd, 2013, pp. 5–48.
85. Mark S. Meskin, Wayne R. Bidlack, Audra J. Davies, et al. *Phytochemicals: Mechanisms of Action*. 1st ed. 2003.
86. Koeberle A, Werz O. Natural products as inhibitors of prostaglandin E2 and pro-inflammatory 5-lipoxygenase-derived lipid mediator biosynthesis. *Biotechnol Adv* 2018; 36: 1709–1723.
87. Koeberle A, Northoff H, Werz O. Curcumin blocks prostaglandin E2 biosynthesis through direct inhibition of the microsomal prostaglandin E2 synthase-1. *Mol Cancer Ther* 2009; 8: 2348–2355.
88. Samuelsson B, Morgenstern R, Jakobsson PJ. Membrane prostaglandin E synthase-1: a novel therapeutic target. *Pharmacol Rev* 2007; 59: 207–224.
89. Knights KM, Mangoni AA, Miners JO. Defining the COX inhibitor selectivity of NSAIDs: implications for understanding toxicity. *Expert Rev Clin Pharmacol* 2010; 3: 769–776.
90. Cock I. *Medicinal and aromatic plants – Australia*. Ethnopharmacology. 2011.
91. Cordell GA. Biodiversity and drug discovery—a symbiotic relationship. *Phytochemistry* 2000; 55: 463–480.
92. Harding K, Benson EE, Nunes E da C, et al. Can Biospecimen Science Expedite the Ex Situ Conservation of Plants in Megadiverse Countries? A Focus on the Flora of Brazil. *Crit Rev Plant Sci* 2013; 32: 411–444.
93. Jamir K, Seshagirirao K, Meitei MD. Indigenous oral knowledge of wild medicinal plants from the Peren district of Nagaland, India in the Indo Burma hot-spot. *Acta Ecol Sin* 2021; 18.
94. Desta KT, Kim GS, Abd El-Aty AM, et al. Flavone polyphenols dominate in *Thymus schimperi* Ronniger: LC-ESI-MS/MS characterization and study of anti-proliferative effects of plant extract on AGS and HepG2 cancer cells. *J Chromatogr B Analyt Technol Biomed Life Sci* 2017; 1053: 1–8.
95. Efferth T, Koch E. Complex interactions between phytochemicals. The multi-target therapeutic concept of phytotherapy. *Curr Drug Targets* 2011; 12: 122–132.
96. Manworren RC. Multimodal pain management and the future of a personalized medicine approach to pain. *AORN J* 2015; 101: 308–318.
97. Young A, Buvanendran A. Recent advances in multimodal analgesia. *Anesthesiol Clin* 2012; 30: 91–100.
98. Helander EM, Menard BL, Harmon CM, et al. Multimodal Analgesia, Current Concepts, and Acute Pain Considerations. *Curr Pain Headache Rep* 2017; 21: 3.



99. Gavariæ N, Kladar N, Mišan A, et al. Postdistillation waste material of thyme (*Thymus vulgaris* L., Lamiaceae) as a potential source of biologically active compounds. *Ind Crops Prod* 2015; 74: 457–464.
100. Al-Snafi A. Chemical constituents and pharmacological effects of *Asclepias curassavica* – A review. *Asian J Pharm Res* 2015; 5: 83–87.
101. Rezende CM, Fraga SRG. Chemical and aroma determination of the pulp and seeds of murici (*Byrsonima crassifolia* L.). *J Braz Chem Soc* 2003; 14: 425–428.
102. Maldini M, Sosa S, Montoro P, et al. Screening of the topical anti-inflammatory activity of the bark of *Acacia cornigera* Willdenow, *Byrsonima crassifolia* Kunth, *Sweetia panamensis* Yakovlev and the leaves of *Sphagneticola trilobata* Hitchcock. *J Ethnopharmacol* 2009; 122: 430–433.
103. Otunola GA, Oloyede OB, Oladiji AT, et al. Comparative analysis of the chemical composition of three spices – *Allium sativum* L. *Zingiber officinale* Rosc. and *Capsicum frutescens* L. commonly consumed in Nigeria. *Afr J Biotechnol* 2010; 9: 6927–6931.
104. Peraza-Sánchez SR, Cen-Pacheco F, Noh-Chimal A, et al. Leishmanicidal evaluation of extracts from native plants of the Yucatan peninsula. *Fitoterapia* 2007; 78: 315–318.
105. Hernández I, Márquez L, Martínez I, et al. Anti-inflammatory effects of ethanolic extract and alkaloids-derived from *Heliopsis longipes* roots. *J Ethnopharmacol* 2009; 124: 649–652.
106. Srianthi D, Udayangani DN, Chamari H. Antioxidant, antibacterial and anti-inflammatory potential of the aqueous extract of the raw leaves of sri lankan variety of *persea americana miller* (avocado). *Int J Ayurveda Pharma Res* 2020; 8: 1–11.
107. Chaves OS, Gomes RA, Tomaz AC de A, et al. Secondary Metabolites from *Sida rhombifolia* L. (Malvaceae) and the Vasorelaxant Activity of Cryptolepinone. *Molecules* 2013; 18: 2769–2777.
108. Sabandar CW, Ahmat N, Jaafar FM, et al. Medicinal property, phytochemistry and pharmacology of several *Jatropha* species (Euphorbiaceae): a review. *Phytochemistry* 2013; 85: 7–29.
109. Pudji A. The ability of anti-inflammatory *jatropha curcas* leaf extract at cox-2 expression on monocytes were exposed LPS. *UNEJ E-Proceeding* 2017; 154–157.
110. Shankland WE 2nd. Four common herbs seen in dental practice: properties and potential adverse effects. *Cranio* 2009; 27: 118–124.
111. Kim S, Kim DB, Jin W, et al. Comparative studies of bioactive organosulphur compounds and antioxidant activities in garlic (*Allium sativum* L.), elephant garlic (*Allium ampeloprasum* L.) and onion (*Allium cepa* L.). *Nat Prod Res* 2018; 32: 1193–1197.
112. Salehi B, Valussi M, Morais-Braga MFB, et al. *Tagetes* spp. Essential Oils and Other Extracts: Chemical Characterization and Biological Activity. *Molecules* 2018; 23: 2847.
113. Céspedes CL, Avila JG, Martínez A, et al. Antifungal and antibacterial activities of Mexican tarragon (*Tagetes lucida*). *J Agric Food Chem* 2006; 54: 3521–3527.
114. Vergara Barragán E, Bach H, Meza-Reyes S, et al. Bioactivities of Flavonoids from *Lopezia racemosa*. *BioMed Res Int* 2019; 2019: 3286489.
115. Cruz Paredes C, Bolívar Balbás P, Gómez-Velasco A, et al. Antimicrobial, antiparasitic, anti-inflammatory, and cytotoxic activities of *Lopezia racemosa*. *ScientificWorldJournal* 2013; 2013: 237438.
116. Srinath J. Therapeutic Potential of *Spilanthes acmella* – A Dental Note. *Int J Pharm Sci Rev Res* 2014; 151–153.
117. Pareek S, Sagar NA, Sharma S, et al. Onion (*Allium cepa* L.). In: Yahia EM (ed) *Fruit and Vegetable Phytochemicals*. Chichester, UK: John Wiley & Sons, Ltd, 2017, pp. 1145–1162.
118. Babotă M, Mocan A, Vlase L, et al. Phytochemical Analysis, Antioxidant and Antimicrobial Activities of *Helichrysum arenarium* (L.) Moench. and *Antennaria dioica* (L.) Gaertn. Flowers. *Molecules* 2018; 23: 409.
119. Gafrikova M, Galova E, Sevcovicova A, et al. Extract from *Armoracia rusticana* and Its Flavonoid Components Protect Human Lymphocytes against Oxidative Damage Induced by Hydrogen Peroxide. *Molecules* 2014; 19: 3160–3172.
120. Singh J, Upadhyay AK, Bahadur A, et al. Antioxidant phytochemicals in cabbage (*Brassica oleracea* L. var. capitata). *Sci Hort* 2006; 108: 233–237.
121. Singh N, Rao AS, Nandal A, et al. Phytochemical and pharmacological review of *Cinnamomum verum* J. Presl-a versatile spice used in food and nutrition. *Food Chem* 2021; 338: 127773.
122. Dezsi ‘., Bādārāu AS, Bischin C, et al. Antimicrobial and Antioxidant Activities and Phenolic Profile of *Eucalyptus globulus* Labill. and *Corymbia ficifolia* (F. Muell.) K.D. Hill & L.A.S. Johnson Leaves. *Molecules* 2015; 20: 4720–4734.
123. Al-Snafi A. Therapeutic importance of *Hyoscyamus* species grown in Iraq (*Hyoscyamus*

- albus, *Hyoscyamus niger* and *Hyoscyamus reticulatus*-A review. *IOSR J Pharm* 2018; 8: 18–32.
124. Singh O, Khanam Z, Misra N, et al. Chamomile (*Matricaria chamomilla* L.): An overview. *Pharmacogn Rev* 2011; 5: 82–95.
125. Roby MHH, Sarhan MA, Selim KA-H, et al. Antioxidant and antimicrobial activities of essential oil and extracts of fennel (*Foeniculum vulgare* L.) and chamomile (*Matricaria chamomilla* L.). *Ind Crops Prod* 2013; 44: 437–445.
126. Cheikh-Rouhou S, Besbes S, Hentati B, et al. *Nigella sativa* L.: Chemical composition and physicochemical characteristics of lipid fraction. *Food Chem* 2007; 101: 673–681.
127. Beara IN, Lesjak MM, Orèià DZ, et al. Comparative analysis of phenolic profile, antioxidant, anti-inflammatory and cytotoxic activity of two closely-related Plantain species: *Plantago altissima* L. and *Plantago lanceolata* L. *LWT - Food Sci Technol* 2012; 47: 64–70.
128. Beara IN, Orèià DZ, Lesjak MM, et al. Liquid chromatography/tandem mass spectrometry study of anti-inflammatory activity of Plantain (*Plantago* L.) species. *J Pharm Biomed Anal* 2010; 52: 701–706.
129. Afonso AF, Pereira OR, Fernandes Â, et al. Phytochemical Composition and Bioactive Effects of *Salvia africana*, *Salvia officinalis* 'Icterina' and *Salvia mexicana* Aqueous Extracts. *Molecules* 2019; 24: 4327.
130. Ghasemian M, Owlia S, Owlia MB. Review of Anti-Inflammatory Herbal Medicines. *Adv Pharmacol Sci* 2016; 2016: 9130979.
131. Kaur K, Kaushal S. Phytochemistry and pharmacological aspects of *Syzygium aromaticum*: A review. *J Pharmacogn Phytochem* 2019; 8: 398–406.
132. Déciga-Campos M, Beltrán-Villalobos KL, Aguilar-Mariscal H, et al. Synergistic Herb-Herb Interaction of the Antinociceptive and Anti-Inflammatory Effects of *Syzygium aromaticum* and *Rosmarinus officinalis* Combination. *Evid Based Complement Alternat Med* 2021; 2021: 8916618.
133. Kisiel W, Barszcz B. Further sesquiterpenoids and phenolics from *Taraxacum officinale*. *Fitoterapia* 2000; 71: 269–273.
134. Moldovan ML, Carpa R, Fize'an I, et al. Phytochemical Profile and Biological Activities of Tendrils and Leaves Extracts from a Variety of *Vitis vinifera* L. *Antioxidants* 2020; 9: 373.
135. Nicoli F, Negro C, Vergine M, et al. Evaluation of Phytochemical and Antioxidant Properties of 15 Italian *Olea europaea* L. Cultivar Leaves. *Molecules* 2019; 24: 1998.
136. Günaydin K, Savci S. Phytochemical studies on *Ruta chalepensis* (Lam.) Lamarck. *Nat Prod Res* 2005; 19: 203–210.
137. Badgujar SB, Patel VV, Bandivdekar AH. *Foeniculum vulgare* Mill: a review of its botany, phytochemistry, pharmacology, contemporary application, and toxicology. *Biomed Res Int* 2014; 2014: 842674.
138. Choi EM, Hwang JK. Antiinflammatory, analgesic and antioxidant activities of the fruit of *Foeniculum vulgare*. *Fitoterapia* 2004; 75: 557–565.
139. Maroyi A. *Euclea undulata* Thunb.: Review of its botany, ethnomedicinal uses, phytochemistry and biological activities. *Asian Pac J Trop Med* 2017; 10: 1030–1036.
140. Tefera BN, Kim YD. Ethnobotanical study of medicinal plants in the Hawassa Zuria District, Sidama zone, Southern Ethiopia. *J Ethnobiol Ethnomed* 2019; 15: 25.
141. Buyinza D, Chalo DM, Derese S, et al. Flavonoids and Isoflavonoids of *Millettia dura* and *Millettia ferruginea*: Phytochemical review and chemotaxonomic values. *Biochem Syst Ecol* 2020; 91: 104053.
142. Wondimieneh S, Asres K. In Vivo Anti-inflammatory and Antinociceptive Activities of *Salvia nilotica* and *Rosa abyssinica*. *Ethiop Pharm J* 2008; 26: 75–82.
143. Itou RDGE, Sanogo R, Ossibi AWE, et al. Anti-Inflammatory and Analgesic Effects of Aqueous Extract of Stem Bark of *Ceiba pentandra* Gaertn. *Pharmacol Pharm* 2014; 05: 1113–1118.
144. Issa TO, Mohamed YS, Yagi S, et al. Ethnobotanical investigation on medicinal plants in Algoz area (South Kordofan), Sudan. *J Ethnobiol Ethnomed* 2018; 14: 31.
145. Dirar AI, Adhikari-Devkota A, Kunwar RM, et al. Genus *Blepharis* (Acanthaceae): A review of ethnomedicinally used species, and their phytochemistry and pharmacological activities. *J Ethnopharmacol* 2021; 265: 113255.
146. Tesfaye S, Belete A, Engidawork E, et al. Ethnobotanical Study of Medicinal Plants Used by Traditional Healers to Treat Cancer-Like Symptoms in Eleven Districts, Ethiopia. *Evid Based Complement Alternat Med* 2020; 2020: 7683450.
147. Jimoh MO, Afolayan AJ, Lewu FB. Antioxidant and phytochemical activities of *Amaranthus caudatus* L. harvested from different soils at various growth stages. *Sci Rep* 2019; 9: 12965.
148. Ashu Agbor M, Naidoo S. Ethnomedicinal Plants Used by Traditional Healers to Treat Oral Health Problems in Cameroon. *Evid Based Complement*

- Alternat Med* 2015; 2015: 649832.
149. Pareek S, Sagar N, Sharma S, et al. Onion (*Allium cepa* L.): Chemistry and Human Health. In: *Fruit and Vegetable Phytochemicals*. 2017, pp. 1145–1162.
  150. Elgorashi EE, McGaw LJ. African plants with in vitro anti-inflammatory activities: A review. *South Afr J Bot* 2019; 126: 142–169.
  151. Matata DZ, Moshi MJ, Machumi F, et al. Isolation of a new cytotoxic compound, 3-((Z)-heptadec-14-enyl) benzene - 1-ol from *Rhus natalensis* root extract. *Phytochem Lett* 2020; 36: 120–126.
  152. Kidane B, van Andel T, van der Maesen LJJ, et al. Use and management of traditional medicinal plants by Maale and Ari ethnic communities in southern Ethiopia. *J Ethnobiol Ethnomed* 2014; 10: 46.
  153. Alqasoumi SI, Basudan OA, Alam P, et al. Antioxidant study of flavonoid derivatives from the aerial parts of *Rhus natalensis* growing in Saudi Arabia. *Pak J Pharm Sci* 2016; 29: 97–103.
  154. Martins MR, Arantes S, Candeias F, et al. Antioxidant, antimicrobial and toxicological properties of *Schinus molle* L. essential oils. *J Ethnopharmacol* 2014; 151: 485–492.
  155. Bringmann G, Rüdener S, Irmer A, et al. Antitumoral and antileishmanial dioncoquinones and ancistroquinones from cell cultures of *Triphyophyllum peltatum* (Dioncophyllaceae) and *Ancistrocladus abbreviatus* (Ancistrocladaceae). *Phytochemistry* 2008; 69: 2501–2509.
  156. Zougagh S, Belghiti A, Rochd T, et al. Medicinal and Aromatic Plants Used in Traditional Treatment of the Oral Pathology: The Ethnobotanical Survey in the Economic Capital Casablanca, Morocco (North Africa). *Nat Prod Bioprospect* 2019; 9: 35–48.
  157. Khalil N, Bishr M, Desouky S, et al. Ammi *Visnaga* L., a Potential Medicinal Plant: A Review. *Molecules* 2020; 25: 301.
  158. Laribi B, Kouki K, M'Hamdi M, et al. Coriander (*Coriandrum sativum* L.) and its bioactive constituents. *Fitoterapia* 2015; 103: 9–26.
  159. Tugume P, Kakudidi EK, Buyinza M, et al. Ethnobotanical survey of medicinal plant species used by communities around Mabira Central Forest Reserve, Uganda. *J Ethnobiol Ethnomed* 2016; 12: 5.
  160. Woode E, Ansah C, Ainooson GK, et al. Anti-inflammatory and antioxidant properties of the root extract of *Carissa edulis* (forsk.) Vahl (apocynaceae). *J Sci Technol Ghana* 2007; 27: 5–15.
  161. Farkhondeh T, Kianmehr M, Kazemi T, et al. Toxicity effects of *Nerium oleander*, basic and clinical evidence: A comprehensive review. *Hum Exp Toxicol* 2020; 39: 773–784.
  162. Shafiq Y, Naqvi SBS, Rizwani GH, et al. A mechanistic study on the inhibition of bacterial growth and inflammation by *Nerium oleander* extract with comprehensive in vivo safety profile. *BMC Complement Med Ther* 2021; 21: 135.
  163. Megersa M, Asfaw Z, Kelbessa E, et al. An ethnobotanical study of medicinal plants in Wayu Tuka District, East Welega Zone of Oromia Regional State, West Ethiopia. *J Ethnobiol Ethnomed* 2013; 9: 68.
  164. Wangteeraprasert R, Lipipun V, Gunaratnam M, et al. Bioactive Compounds from *Carissa spinarum*. *Phytother Res* 2012; 26: 1496–1499.
  165. Punia DP. A review on varieties of Arka *Calotropis procera* (AITON) Dryand and *Calotropis gigantea* (L.) Dryand. *Global J Res Med Plants & Indigen Med* 2013; 2: 392–400.
  166. Ghosh PK, Bhattacharjee P, Mitra S, et al. Physicochemical and Phytochemical Analyses of Copra and Oil of *Cocos nucifera* L. (West Coast Tall Variety). *Int J Food Sci* 2014; 2014: 310852.
  167. Lima EB, Sousa CN, Meneses LN, et al. *Cocos nucifera* (L.) (Arecaceae): A phytochemical and pharmacological review. *Braz J Med Biol Res* 2015; 48: 953–964.
  168. Chithra MA, Ijini TP, Kharkwal H, et al. Phenolic rich *Cocos nucifera* inflorescence extract ameliorates inflammatory responses in LPS-stimulated RAW264.7 macrophages and toxin-induced murine models. *Inflammopharmacology* 2020; 28: 1073–1089.
  169. Teklay A, Abera B, Giday M. An ethnobotanical study of medicinal plants used in Kiltie Awulaelo District, Tigray Region of Ethiopia. *J Ethnobiol Ethnomed* 2013; 9: 65.
  170. Misganaw D, Sahile S, Negash W. In vitro antimicrobial effects of *Gomphocarpus purpurascens* a. rich against standard and clinically isolated microorganisms. *Glob J Sci Res* 2019; 7: 121–136.
  171. Hassan H, Ahmadu AA, Hassan AS. Analgesic and anti-inflammatory activities of *Asparagus africanus* root extract. *Afr J Tradit Complement Altern Med* 2007; 5: 27–31.
  172. Abera B. Medicinal plants used in traditional medicine by Oromo people, Ghimbi District, Southwest Ethiopia. *J Ethnobiol Ethnomed* 2014; 10: 40.
  173. Yimer T, Birru EM, Adugna M, et al. Evaluation of Analgesic and Anti-Inflammatory Activities of 80% Methanol Root Extract of *Echinops kebericho* M. (Asteraceae). *J Inflamm Res* 2020; 13: 647–658.
  174. Liu X, Wang X, Chen Z, et al. De novo assembly

- and comparative transcriptome analysis: novel insights into terpenoid biosynthesis in *Chamaemelum nobile* L. *Plant Cell Rep* 2019; 38: 101–116.
175. Msaada K, Salem N, Bachrouh O, et al. Chemical Composition and Antioxidant and Antimicrobial Activities of Wormwood (*Artemisia absinthium* L.) Essential Oils and Phenolics. *J Chem* 2015; 2015: 1–12.
  176. Bouabid K, Lamchouri F, Toufik H, et al. Phytochemical investigation, in vitro and in vivo antioxidant properties of aqueous and organic extracts of toxic plant: *Atractylis gummifera* L. *J Ethnopharmacol* 2020; 253: 112640.
  177. Ngueguim TF, Djouwoug Noussi C, Donfack JH, et al. Acute and sub-acute toxicity of a lyophilised aqueous extract of the aerial part of *Spilanthes africana* Delile in rats. *J Ethnopharmacol* 2015; 172: 145–154.
  178. Okunade AL. *Ageratum conyzoides* L. (Asteraceae). *Fitoterapia* 2002; 73: 1–16.
  179. Mohammed T, Teshale C. Preliminary phytochemical screening and evaluation of antibacterial activity of *Dichrocephala integrifolia* (L.f) O. Kuntze. *J Intercult Ethnopharmacol* 2012; 1: 30–34.
  180. Uthpala TGG, Navaratne SB. *Acmella oleracea* Plant; Identification, Applications and Use as an Emerging Food Source – Review. *Food Rev Int* 2021; 37: 399–414.
  181. Letha N, Ganesan K, Kumar S, et al. Studies on phytochemical screening and in vitro antioxidant activity of Ethiopian indigenous medicinal plants, *Artemisia abyssinica* Sch.Bip. Ex A.Rich. *World J Pharm Res* 2016; 5: 1048–1058.
  182. Tariku Y, Hymete A, Hailu A, et al. In vitro evaluation of antileishmanial activity and toxicity of essential oils of *Artemisia absinthium* and *Echinops kebericho*. *Chem Biodivers* 2011; 8: 614–623.
  183. Studzińska-Sroka E, Dudek-Makuch M, Chanaj-Kaczmarek J, et al. Anti-inflammatory Activity and Phytochemical Profile of *Galinsoga parviflora* Cav. *Molecules* 2018; 23: 2133.
  184. Ali S, Zameer S, Yaqoob M. Ethnobotanical, phytochemical and pharmacological properties of *Galinsoga parviflora* (Asteraceae): A review. *Trop J Pharm Res* 2017; 16: 3023–3033.
  185. Albejo B, Endale M, Kibret B, et al. Phytochemical investigation and antimicrobial activity of leaves extract of *Vernonia auriculifera* Hiern. *J Pharm Pharmacogn Res* 2015; 3: 141–147.
  186. Speroni E, Cervellati R, Innocenti G, et al. Anti-inflammatory, anti-nociceptive and antioxidant activities of *Balanites aegyptiaca* (L.) Delile. *J Ethnopharmacol* 2005; 98: 117–125.
  187. Traore KT, Ouédraogo N, Belemnaba L, et al. Anti-inflammatory and analgesic activities of extracts from *Balanites aegyptiaca* L. Delile (Balanitaceae) root bark: Plant used against liver diseases in Burkina Faso. *Afr J Pharm Pharmacol* 2019; 13: 322–329.
  188. Compaoré M, Lamien-Meda A, Mogo'an C, et al. Antioxidant, diuretic activities and polyphenol content of *Stereospermum kunthianum* Cham. (Bignoniaceae). *Nat Prod Res* 2011; 25: 1777–1788.
  189. Ogundajo A, Ashafa AT. Phytochemical Compositions and In vitro Assessments of Antioxidant and Antidiabetic Potentials of Fractions from *Ehretia cymosa* Thonn. *Pharmacogn Mag* 2017; 13: S470–S480.
  190. Yismaw YE, Abdelwuhab M, Ambikar DB, et al. Phytochemical and Antiulcer Activity Screening of Seed Extract of *Cordia africana* Lam (Boraginaceae) in Pyloric Ligated Rats. *Clin Pharmacol Adv Appl* 2020; 12: 67–73.
  191. Riazullah, Hussain I, Badrullah. Phytochemical and anti-microbial activity of *Lepidium sativum* L. *J Med Plants Res* 2012; 6: 4358–4361.
  192. Maroyi A. *Boscia salicifolia*: review of its botany, medicinal uses, phytochemistry and biological activities. *J Pharm Sci* 2019; 11: 3055–3060.
  193. Tekulu GH, Hiluf T, Brhanu H, et al. Anti-inflammatory and anti-nociceptive property of *Capparis tomentosa* Lam. root extracts. *J Ethnopharmacol* 2020; 253: 112654.
  194. Abdulaziz Al-Hamoud G, Saud Orfali R, Sugimoto S, et al. Four New Flavonoids Isolated from the Aerial Parts of *Cadaba rotundifolia* Forssk. (Qadab). *Molecules* 2019; 24: 2167.
  195. Martial N, Dah-Nouvlessounon D, Christine N tcha, et al. Phytochemistry and biological activities of *Cratogeomys adansonii* extracts. *Int J Pharm Pharm Sci* 2018; 10: 62–67.
  196. Zunjar V, Mammen D, Trivedi B, et al. Pharmacognostic, Physicochemical and Phytochemical Studies on *Carica papaya* Linn. Leaves. *Pharmacogn J* 2011; 3: 5–8.
  197. Kashyap K, Sarkar P, Kalita MC, et al. A review on the widespread therapeutic application of the traditional herb *Drymaria cordata*. *Int J Pharma Bio Sci* 2014; 5: 696–705.
  198. Kokanova-Nedialkova Z, Nedialkov PT, Nikolov SD. The Genus *Chenopodium*: Phytochemistry, Ethnopharmacology and Pharmacology. *Pharmacogn Rev* 2009; 3: 280–306.
  199. Kumar R, Mishra AK, Dubey NK, et al. Evaluation of *Chenopodium ambrosioides* oil as a potential source of antifungal, anti-aflatoxinogenic and antioxidant activity. *Int J Food Microbiol* 2007; 115: 159–164.

200. Muriithi E, Bojase-Moleta G, Majinda RRT. Benzophenone derivatives from *Garcinia livingstonei* and their antioxidant activities. *Phytochem Lett* 2016; 18: 29–34.
201. Yang H, Figueroa M, To S, et al. Benzophenones and biflavonoids from *Garcinia livingstonei* fruits. *J Agric Food Chem* 2010; 58: 4749–4755.
202. Ashokkumar K. *Gloriosa superba* (L.): A Brief Review of its Phytochemical Properties and Pharmacology. *Int J Pharmacogn Phytochem Res* 2015; 7: 1190–1193.
203. Arbab A. Review on anogeissus leiocarpus a potent african traditional drug. *Int J Res Pharm Chem* 2014; 4: 496–500.
204. Okoli CO, Akah PA, Nwafor SV, et al. Anti-inflammatory activity of hexane leaf extract of *Aspilia africana* C.D. Adams. *J Ethnopharmacol* 2007; 109: 219–225.
205. Alara OR, Abdurahman NH, Mudalip SKA, et al. Phytochemical and pharmacological properties of *Vernonia amygdalina*: A review. *J Chem Eng Ind Biotechnol* 2017; 2: 80–96.
206. Kriplani P, Guarve K, Baghael US. *Arnica montana* L. – a plant of healing: review. *J Pharm Pharmacol* 2017; 69: 925–945.
207. Fernandes JM, Cunha LM, Azevedo EP, et al. *Kalanchoe laciniata* and *Bryophyllum pinnatum*: an updated review about ethnopharmacology, phytochemistry, pharmacology and toxicology. *Rev Bras Farmacogn* 2019; 29: 529–558.
208. Soh D, Bakang BT, Tchouboun EN, et al. New cucurbitane type triterpenes from *Momordica foetida* Schumach. (Cucurbitaceae). *Phytochem Lett* 2020; 38: 90–95.
209. Guimarães R, Barros L, Carvalho AM, et al. Aromatic plants as a source of important phytochemicals: Vitamins, sugars and fatty acids in *Cistus ladanifer*, *Cupressus lusitanica* and *Eucalyptus gunnii* leaves. *Ind Crops Prod* 2009; 30: 427–430.
210. Salih AM, Al-Qurainy F, Khan S, et al. Mass propagation of *Juniperus procera* Hoechst. Ex Endl. From seedling and screening of bioactive compounds in shoot and callus extract. *BMC Plant Biol* 2021; 21: 192.
211. Ilodibia\* CV, Ugwu RU, Okeke CU, et al. Phytochemical evaluation of various parts of *Dracaena arborea* Link. and *Dracaena mannii* Bak. *Afr J Plant Sci* 2015; 9: 287–292.
212. Kilonzo M, Rubanza C, Richard U, et al. Antimicrobial activities and phytochemical analysis of extracts from *Ormocarpum trichocarpum* (Taub.) and *Euclea divinorum* (Hiern) used as traditional medicine in Tanzania. *Tanzan J Health Res* 2019; 21: 1–12.
213. Jena J, Gupta A. *Ricinus communis* linn: A phytopharmacological review. *Int J Pharm Pharm Sci* 2012; 4: 25–29.
214. Martínez CA, Mosquera OM, Niño J. Medicinal plants from the genus *Alchornea* (Euphorbiaceae): A review of their ethnopharmacology uses and phytochemistry. *Bol Latinoam Caribe Plant Med Aromat* 2017; 16: 162–205.
215. Yakubu OF, Adebayo AH, Iweala EEJ, et al. Anti-inflammatory and antioxidant activities of fractions and compound from *Riciodendron heudelotii* (Baill.). *Heliyon* 2019; 5: e02779.
216. Koech S, Maoga J, Sindani A, et al. Anti-Inflammatory Activity of Dichloromethanolic Root Extract of *Clutia abyssinica* in Swiss Albino Mice. *J Pharmacogn Nat Prod* 2017; 3: 1000132.
217. Seebaluck R, Gurib-Fakim A, Mahomoodally F. Medicinal plants from the genus *Acalypha* (Euphorbiaceae)—A review of their ethnopharmacology and phytochemistry. *J Ethnopharmacol* 2015; 159: 137–157.
218. Bruno T, Soh D, Ernestine N, et al. Phytochemical Composition and Biological Activity of *Faidherbia albida* (Mimosaceae) Roots and Leaves. *Int J Pharm Sci Rev Res* 2020; 65(1): 124–130.
219. Hebbar SS, Harsha VH, Shripathi V, et al. Ethnomedicine of Dharwad district in Karnataka, India—plants used in oral health care. *J Ethnopharmacol* 2004; 94: 261–266.
220. Kalaivani T, Mathew L. Free radical scavenging activity from leaves of *Acacia nilotica* (L.) Wild. ex Delile, an Indian medicinal tree. *Food Chem Toxicol* 2010; 48: 298–305.
221. Ali A, Naveed A, Khan B, et al. *Acacia nilotica*: A plant of multipurpose medicinal uses. *J Med Plants* 2012; 6: 1492–1496.
222. Mariita RM, Orodho JA, Okemo PO, et al. Antifungal, antibacterial and antimycobacterial activity of *Entada abyssinica* Steudel ex A. Rich (Fabaceae) methanol extract. *Pharmacogn Res* 2010; 2: 163–168.
223. Gurmessa GT, Kusari S, Laatsch H, et al. Chemical constituents of root and stem bark of *Erythrina brucei*. *Phytochem Lett* 2018; 25: 37–42.
224. Narnoliya LK, Jadaun JS, Singh SP. The Phytochemical Composition, Biological Effects and Biotechnological Approaches to the Production of High-Value Essential Oil from *Geranium*. In: Malik S (ed) *Essential Oil Research: Trends in Biosynthesis, Analytics, Industrial Applications and Biotechnological Production*. 2019, pp. 327–352.
225. Adesuyi A, Elumm I, Adaramola F, et al. Nutritional and Phytochemical Screening of *Garcinia kola*. *Adv J Food Sci Technol* 2012; 4:

- 9–14.
226. Petersen M, Simmonds MS. Rosmarinic acid. *Phytochemistry* 2003; 62: 121–125.
227. Njeru SN, Obonyo M, Nyambati S, et al. Antimicrobial and cytotoxicity properties of the organic solvent fractions of *Clerodendrum myricoides* (Hochst.) R. Br. ex Vatke: Kenyan traditional medicinal plant. *J Intercult Ethnopharmacol* 2016; 5: 226–232.
228. Alemayehu K, Anza M, Engdaw D, et al. Chemical constituents, physicochemical properties and antibacterial activity of leaves essential oil of *Ocimum urticifolium*. *J Coast Life Med* 2016; 4: 955–960.
229. Idris S, Ndukwe G, Gimba C. Preliminary phytochemical screening and antimicrobial activity of seed extracts of *Persea americana* (avocado pear). *Bayero J Pure Appl Sci* 2009; 2: 173–176.
230. Tirfe M, Gebrehiwot M, Gebrelibanos M, et al. Radical Scavenging Activity and Preliminary Phytochemical Screening of Pods of *Cassia arereh* Del. (Fabaceae). *Momona Ethiop J Sci* 2015; 7: 125–133.
231. Jeremiah C, Aliyu N, Dijie H, et al. Pharmacognostic and Elemental Analysis of the Leaves of *Tapinanthus globifer* (A. Rich). *Tiegh. Res J Pharmacol* 2018; 6: 11–18.
232. Abiche E, Habila J. Phytochemistry, pharmacology and medicinal uses of *Cola* (Malvaceae) family: a review. *Med Chem Res* 2020; 29: 2089–2105.
233. Imam H, et al. Neem (*Azadirachta indica* A. Juss)-A Nature's Drugstore: An overview. *Int Res J Biol Sci* 2012; 1: 76–79.
234. Sahrawat A, Sharma J, Rahul S, et al. Phytochemical analysis and Antibacterial properties of *Azadirachta indica* (Neem) leaves extract against *E.coli*. *J Pharmacogn Phytochem* 2018; 7: 1368–1371.
235. Lakshmi T, Krishnan V, Rajendran R, et al. *Azadirachta indica*: A herbal panacea in dentistry – An update. *Pharmacogn Rev* 2015; 9: 41–44.
236. Qureshi H, Arshad M, Akram A, et al. Ethnopharmacological and phytochemical account of paradise tree (*Melia azedarach* L.: Meliaceae). *Pure Appl Biol* 2015; 5: 5–14.
237. Aæimoviæ M, Jeremiæ K, Salaj N, et al. *Marrubium vulgare* L.: A Phytochemical and Pharmacological Overview. *Molecules* 2020; 25: 2898.
238. Pascal DrMK, my el abbes F, Meddah B, et al. Assessment of methanolic extract of *Marrubium vulgare* for antiinflammatory, analgesic and anti-microbiologic activities. *J Chem Pharm Res* 2011; 3: 199–204.
239. Bouyahya A, Chamkhi I, Benali T, et al. Traditional use, phytochemistry, toxicology, and pharmacology of *Origanum majorana* L. *J Ethnopharmacol* 2021; 265: 113318.
240. Negi A, Dobhal K, Ghildiyal P. Antioxidant Potential and Effect of Extraction Solvent on Total Phenol Content, Flavonoids Content and Tannin Content of *Ficus palmata* Forssk. *Int J Pharm Sci Rev Res* 2018; 49: 19–24.
241. Anibasa G. Antimicrobial And Phytochemical Screening Activities Of *Ficus Sur* (Forssk). *N Y Sci J* 2011; 4(1): 15–18.
242. Taviano MF, Rashed K, Filocamo A, et al. Phenolic profile and biological properties of the leaves of *Ficus vasta* Forssk. (Moraceae) growing in Egypt. *BMC Complement Altern Med* 2018; 18: 161.
243. Paikra BK, Dhongade H kumar J, Gidwani B. Phytochemistry and Pharmacology of *Moringa oleifera* Lam. *J Pharmacopuncture* 2017; 20: 194–200.
244. Messaoud C, Laabidi A, Boussaid M. *Myrtus communis* L. infusions: the effect of infusion time on phytochemical composition, antioxidant, and antimicrobial activities. *J Food Sci* 2012; 77: C941–C947.
245. Kaushal S. Phytochemistry and pharmacological aspects of *Syzygium aromaticum*: A review. *J Pharmacogn Phytochem* 2019; 8: 398–406.
246. Kamath JV, Rahul N, Kumar CKA, et al. *Psidium guajava* L: A review. *Int J Green Pharm*; 2. Epub ahead of print 2008. DOI: 10.22377/ijgp.v2i1.386.
247. Dhakad AK, Pandey VV, Beg S, et al. Biological, medicinal and toxicological significance of *Eucalyptus* leaf essential oil: a review. *J Sci Food Agric* 2018; 98: 833–848.
248. Al-Snafi A. The pharmacological and therapeutic importance of *Eucalyptus* species grown in Iraq. *IOSR J Pharm* 2017; 7: 72–91.
249. Fatma B, Fatiha M, Elattafia B, et al. Phytochemical and antimicrobial study of the seeds and leaves of *Peganum harmala* L. against urinary tract infection pathogens. *Asian Pac J Trop Dis* 2016; 6: 822–826.
250. Ghasemi Pirbalouti A, Momeni M, Bahmani M. Ethnobotanical study of medicinal plants used by Kurd tribe in Dehloran and Abdanan Districts, Ilam Province, Iran. *Afr J Tradit Complement Altern Med* 2013; 10: 368–385.
251. Mina CN, Farzaei MH, Gholamreza A. Medicinal properties of *Peganum harmala* L. in traditional Iranian medicine and modern phytotherapy: a review. *J Tradit Chin Med* 2015; 35: 104–109.
252. Almeida ML, Freitas WE, de Morais PL, et al. Bioactive compounds and antioxidant potential

- fruit of *Ximenia americana* L. *Food Chem* 2016; 192: 1078–1082.
253. Balkrishna A, Rohela A, Kumar A, et al. Mechanistic Insight into Antimicrobial and Antioxidant Potential of *Jasminum* Species: A Herbal Approach for Disease Management. *Plants* 2021; 10: 1089.
254. Sekharan TR, Mohan MS, Venkatnarayanan R, et al. Pharmacognostical and Preliminary Phytochemical Screening the Leaves of *Jasminum grandiflorum* Linn. *Res J Pharmacogn Phytochem* 2010; 2: 438–440.
255. Rathore S, Bhatt S, Dhyani D, et al. Preliminary phytochemical screening of medicinal plant *Ziziphus mauritiana* Lam fruits. *Int J Curr Pharm Res* 2012; 4: 160–162.
256. Aruna K, Devi P, Rajeswari R, et al. Quantitative phytochemical analysis of *oxalis corniculata* L. (oxalidaceae). *World J Pharm Pharm Sci* 2014; 3: 711–716.
257. Desta KT, Abd El-Aty AM. Triterpenoid and Saponin Rich *Phytolacca dodecandra* L'Herit (Endod): A Review on Its Phytochemistry and Pharmacological Properties. *Mini Rev Med Chem* 2021; 21: 23–34.
258. Nakalembe L, Kasolo JN, Nyatia E, et al. Analgesic and Anti-Inflammatory Activity of Total Crude Leaf Extract of *Phytolacca dodecandra* in Wistar Albino Rats. *Neurosci Med* 2019; 10: 259–271.
259. Manu P, Lal A, Rana S, et al. *Plumbago zeylanica* L.: A mini review. *Int J Pharm Appl* 2012; 3: 399–405.
260. Rajakrishnan R, Lekshmi R, Benil PB, et al. Phytochemical evaluation of roots of *Plumbago zeylanica* L. and assessment of its potential as a nephroprotective agent. *Saudi J Biol Sci* 2017; 24: 760–766.
261. Ojewole JA. Analgesic, anti-inflammatory and hypoglycaemic effects of *Securidaca longepedunculata* (Fresen.) [Polygalaceae. root-bark aqueous extract. *Inflammopharmacology* 2008; 16: 174–181.
262. Mekonnen T, Urga K, Engidawork E. Evaluation of the diuretic and analgesic activities of the rhizomes of *Rumex abyssinicus* Jacq in mice. *J Ethnopharmacol* 2010; 127: 433–439.
263. Kumar S, Singh PK. Phytochemical investigation and antioxidant characterization of essential oil from roots of *Rumex nepalensis* Spreng high altitude of North India. *Mater Today Proc* 2020; 26: 3442–3448.
264. Hawaze S, Deti H, Suleman S. In vitro Antimicrobial Activity and Phytochemical Screening of *Clematis* Species Indigenous to Ethiopia. *Indian J Pharm Sci* 2012; 74: 29–35.
265. Tadele A, Asres K, Melaku D, et al. In vivo anti-inflammatory and antinociceptive activities of the leaf extracts of *Clematis simensis* Fresen. *Ethiop Pharm J* 2010; 27: 33–41.
266. Madivoli ES, Maina EG, Kairigo PK, et al. In vitro antioxidant and antimicrobial activity of *Prunus africana* (Hook. f.) Kalkman (bark extracts) and *Harrisonia abyssinica* Oliv. extracts (bark extracts): A comparative study. *J Med Plants Econ Dev* 2018; 2: 1–9.
267. Bento C, Gonçalves AC, Silva B, et al. Peach (*Prunus Persica*): Phytochemicals and Health Benefits. *Food Rev Int* 2020; 1–32.
268. Venditti A, Guarcini L, Ballero M, et al. Iridoid glucosides from *Pentas lanceolata* (Forsk.) Deflers growing on the Island of Sardinia. *Plant Syst Evol* 2015; 301: 685–690.
269. Lawal IO, Grierson DS, Afolayan AJ. Phytochemical and antioxidant investigations of a *Clausena anisata* hook, a South African medicinal plant. *Afr J Tradit Complement Altern Med* 2015; 12: 28–37.
270. Nantongo JS, Odoi JB, Abigaba G, et al. Variability of phenolic and alkaloid content in different plant parts of *Carissa edulis* Vahl and *Zanthoxylum chalybeum* Engl. *BMC Res Notes* 2018; 11: 125.
271. Dutta S, Shaikh A. The Active chemical constituent and biological activity of *salvadora persica* (Miswak). *Int J Curr Pharm Rev Res* 2012; 3: 1–14.
272. Rani MS, Pippalla RS, Mohan K. *Dodonaea Viscosa* Linn. - An Overview. *Asian J Pharm Res Health Care* 2009; 1: 97–112.
273. Jima T, Megersa M. Ethnobotanical Study of Medicinal Plants Used to Treat Human Diseases in Berbere District, Bale Zone of Oromia Regional State, South East Ethiopia. *Evid Based Complement Alternat Med* 2018; 2018: 8602945.
274. Mergia E, Shibeshi W, Terefe G, et al. Antitrypanosomal activity of *Verbascum sinaiticum* Benth. (Scrophulariaceae) against *Trypanosoma congolense* isolates. *BMC Complement Altern Med* 2016; 16: 362.
275. Guluma T, G NB, Teju E, et al. Phytochemical investigation and evaluation of antimicrobial activities of *Bucea antidysenterica* leaves. *Chem Data Collect* 2020; 28: 100433.
276. Soni P, Siddiqui AA, Dwivedi J, et al. Pharmacological properties of *Datura stramonium* L. as a potential medicinal tree: An overview. *Asian Pac J Trop Biomed* 2012; 2: 1002–1008.
277. Sayyed A. Phytochemistry, pharmacological and traditional uses of *Datura stramonium* L. *J Pharmacogn Phytochem* 2013; 2: 123–125.

278. Oyekunle I, Nwogu U, Orababa O, et al. Phytochemical, Antimicrobial and Proximate Composition of *Nicotiana tabacum* Leaves Extract. *Int J Innov Res Sci Eng Technol*; 4.
279. Sambo H, Olatunde A, Kiyawa S. Phytochemical, Proximate and Mineral Analyses of *Solanum incanum* Fruit. *Int J Chem Mater Environ Res* 2016; 3: 8–13.
280. Sbhatu D, Abraha H. Preliminary Antimicrobial Profile of *Solanum incanum* L.: A Common Medicinal Plant. *Evid Based Complement Alternat Med* 2020; 2020: 3647065.
281. Jaspers MW, Bashir AK, Zwaving JH, et al. Investigation of *Grewia bicolor* juss. *J Ethnopharmacol* 1986; 17: 205–211.
282. Dianita R, Jantan I. Ethnomedicinal uses, phytochemistry and pharmacological aspects of the genus *Premna*: a review. *Pharm Biol* 2017; 55: 1715–1739.
283. Awah FM, Uzoegwu PN, Oyugi JO, et al. Free radical scavenging activity and immunomodulatory effect of *Stachytarpheta angustifolia* leaf extract. *Food Chem* 2010; 119: 1409–1416.
284. Seyfe S, Toma A, Etilso A, et al. Journal of Medicinal Plants Research Phytochemical screening and in vivo antimalarial activities of crude extracts of *Lantana trifolia* root and *Premna oligotricha* leaves in *Plasmodium berghei* infected mice. *J Med Plants Res* 2017; 11: 763–769.
285. Njeru SN, Obonyo MA, Nyambati SO, et al. Antimicrobial and cytotoxicity properties of the crude extracts and fractions of *Premna resinosa* (Hochst.) Schauer (Compositae): Kenyan traditional medicinal plant. *BMC Complement Altern Med* 2015; 15: 295.
286. Chidambara Murthy KN, Vanitha A, Mahadeva Swamy M, et al. Antioxidant and antimicrobial activity of *Cissus quadrangularis* L. *J Med Food* 2003; 6: 99–105.
287. Kumar S, Yadav M, Yadav A, et al. Impact of spatial and climatic conditions on phytochemical diversity and in vitro antioxidant activity of Indian *Aloe vera* (L.) Burm.f. *South Afr J Bot* 2017; 111: 50–59.
288. Ahmed H. Ethnopharmacobotanical study on the medicinal plants used by herbalists in Sulaymaniyah Province, Kurdistan, Iraq. *J Ethnobiol Ethnomed* 2016; 12: 8.
289. Salehi B, Albayrak S, Antolak H, et al. *Aloe* Genus Plants: From Farm to Food Applications and Phytopharmacotherapy. *Int J Mol Sci* 2018; 19: 2843.
290. Sánchez-Machado DI, López-Cervantes J, Sendón R, et al. *Aloe vera*: Ancient knowledge with new frontiers. *Trends Food Sci Technol* 2017; 61: 94–102.
291. Eyob S, Martinsen BK, Tsegaye A, et al. Antioxidant and antimicrobial activities of extract and essential oil of korarima (*Aframomum corrorima* (Braun) P.C.M. Jansen). *Afr J Biotechnol* 2008; 7: 2585–2592.
292. Kumar G, Loganathan K, Rao B. A Review on Pharmacological and Phytochemical Properties of *Zingiber officinale* Roscoe (Zingiberaceae). *J Pharm Res* 2011; 4: 2963–2966.
293. Kumar H, Agrawal R, Kumar V. *Barleria cristata*: perspective towards phytopharmacological aspects. *J Pharm Pharmacol* 2018; 70: 475–487.
294. Muñoz-Acevedo A, Martínez JL, Rai M. *Ethnobotany: Local Knowledge and Traditions*. 1st ed. CRC Press, 2019.
295. Pasaribu G, Budiarto E, Cahyana H, et al. A Review on Genus *Saurauia*: Chemical Compounds and their Biological Activity. *Pharmacogn J* 2020; 12: 657–666.
296. Hong L, Guo Z, Huang K, et al. Ethnobotanical study on medicinal plants used by Maonan people in China. *J Ethnobiol Ethnomed* 2015; 11: 32.
297. Ye CL, Dai DH, Hu WL. Antimicrobial and antioxidant activities of the essential oil from onion (*Allium cepa* L.). *Food Control* 2013; 30: 48–53.
298. Ghamari S, Mohammadrezaei Khorramabadi R, Mardani M, et al. An overview of the most important medicinal plants with anti-toothache property based on ethnobotanical sources in Iran. *J Pharm Sci Res* 2017; 9: 796–799.
299. Bor M, Özdemir F, Türkan I. The effect of salt stress on lipid peroxidation and antioxidants in leaves of sugar beet *Beta vulgaris* L. and wild beet *Beta maritima* L. *Plant Sci* 2003; 164: 77–84.
300. Srivastava R. A review on phytochemical, pharmacological, and pharmacognostical profile of *Wrightia tinctoria*: Adulterant of kurchi. *Pharmacogn Rev* 2014; 8: 36–44.
301. Zhang SX, Tani T, Yamaji S, et al. Glycosyl flavonoids from the roots and rhizomes of *Asarum longerhizomatsum*. *J Asian Nat Prod Res* 2003; 5: 25–30.
302. Hu R, Lin C, Xu W, et al. Ethnobotanical study on medicinal plants used by Mulam people in Guangxi, China. *J Ethnobiol Ethnomed* 2020; 16: 40.
303. Doss A, Anand SP. Preliminary phytochemical screening of *Asteracantha longifolia* and *Pergularia daemia*. *World Appl Sci J* 2012; 18: 233–235.
304. Abd-Alla HI, Shalaby NMM, Hamed MA, et al. Phytochemical composition, protective and



- therapeutic effect on gastric ulcer and  $\alpha$ -amylase inhibitory activity of *Achillea biebersteinii* Afan. *Arch Pharm Res* 2016; 39: 10–20.
305. Sukumaran P, Nair AG, Chinmayee DM, et al. Phytochemical Investigation of *Bidens biternata* (Lour.) Merr. and Sheriff. - a nutrient-rich leafy vegetable from Western Ghats of India. *Appl Biochem Biotechnol* 2012; 167: 1795–1801.
306. Patel S. Harmful and beneficial aspects of *Parthenium hysterophorus*: an update. *3 Biotech* 2011; 1: 1–9.
307. Lunlun G, Wei N, Yang G, et al. Ethnomedicine study on traditional medicinal plants in the Wuliang Mountains of Jingdong, Yunnan, China. *J Ethnobiol Ethnomed* 2019; 15: 20.
308. Farsam H, Amanlou M, Reza Dehpour A, et al. Anti-inflammatory and analgesic activity of *Biebersteinia multifida* DC. root extract. *J Ethnopharmacol* 2000; 71: 443–447.
309. Chi YM, Nakamura M, Zhao XY, et al. A monoterpene alkaloid from *incarvillea sinensis*. *Chem Pharm Bull (Tokyo)* 2005; 53: 1178–1179.
310. Wurchaih, Huar, Menggenqiqig, et al. Medicinal wild plants used by the Mongol herdsmen in Bairin Area of Inner Mongolia and its comparative study between TMM and TCM. *J Ethnobiol Ethnomed* 2019; 15: 32.
311. Yu CH, Tang WZ, Peng C, et al. Diuretic, anti-inflammatory, and analgesic activities of the ethanol extract from *Cynoglossum lanceolatum*. *J Ethnopharmacol* 2012; 139: 149–154.
312. Joshi K. *Cynoglossum L.*: A review on phytochemistry and chemotherapeutic potential. *J Pharmacogn Phytochem* 2016; 5: 32–39.
313. Jeeva K, Thiyagarajan M, Elangovan V, et al. *Caesalpinia coriaria* leaf extracts mediated biosynthesis of metallic silver nanoparticles and their antibacterial activity against clinically isolated pathogens. *Ind Crops Prod* 2014; 52: 714–720.
314. Jiménez-López J, Ruiz-Medina A, Ortega-Barrales P, et al. Phytochemical profile and antioxidant activity of caper berries (*Capparis spinosa L.*): Evaluation of the influence of the fermentation process. *Food Chem* 2018; 250: 54–59.
315. Zhang H, Ma ZF. Phytochemical and Pharmacological Properties of *Capparis spinosa* as a Medicinal Plant. *Nutrients* 2018; 10: 116.
316. Mishra S, Moharana S, Dash M. Review on *Cleome gynandra*. *Int J Res Pharm Chem* 2011; 1: 681–689.
317. Uddin G, Rauf A, Siddiqui BS, et al. Antinociceptive, anti-inflammatory and sedative activities of the extracts and chemical constituents of *Diospyros lotus L.* *Phytomedicine* 2014; 21: 954–959.
318. Rauf A. Phytochemical screening and biological activity of the aerial parts of *Elaeagnus umbellata*. *Sci Res Essays* 2012; 7: 3690–3694.
319. Ho JC, Chen CM. Flavonoids from the aquatic plant *Eriocaulon buergerianum*. *Phytochemistry* 2002; 61: 405–408.
320. Tantray MA, Khan R, Shawl AS, et al. Phenolic glycosides from *Lespedeza juncea*. *Chem Nat Compd* 2008; 44: 591–593.
321. Sroka Z, Bodalska HR, Małol I. Antioxidative Effect of Extracts from *Erodium cicutarium L.* *Z Naturforsch C J Biosci* 1994; 49: 881–884.
322. Di Lorenzo C, Ceschi A, Kupferschmidt H, et al. Adverse effects of plant food supplements and botanical preparations: a systematic review with critical evaluation of causality. *Br J Clin Pharmacol* 2015; 79: 578–592.
323. Raak C, Büssing A, Gassmann G, et al. A systematic review and meta-analysis on the use of *Hypericum perforatum* (St. John's Wort) for pain conditions in dental practice. *Homeopathy* 2012; 101: 204–210.
324. Wei Y, Shu P, Hong J, et al. Qualitative and quantitative evaluation of phenolic compounds in *Iris dichotoma* Pall. *Phytochem Anal* 2012; 23: 197–207.
325. Prajapati MS, Patel JB, Modi K, et al. *Leucas aspera*: A review. *Pharmacogn Rev* 2010; 4: 85–87.
326. Bahramikia S, Yazdanparast R. Phytochemistry and medicinal properties of *Teucrium polium L.* (Lamiaceae). *Phytother Res* 2012; 26: 1581–1593.
327. Sadeghi H, Zarezade V, Sadeghi H, et al. Anti-inflammatory Activity of *Stachys Ptilifera* Benth. *Iran Red Crescent Med J* 2014; 16: e19259.
328. Likhoba CW, Simmonds MSJ, Paton AJ. *Plectranthus*: A review of ethnobotanical uses. *J Ethnopharmacol* 2006; 103: 1–24.
329. Kong DG, Zhao Y, Li GH, et al. The genus *Litsea* in traditional Chinese medicine: An ethnomedical, phytochemical and pharmacological review. *J Ethnopharmacol* 2015; 164: 256–264.
330. Al-Snafi A. *Fritillaria Imperialis*-A Review. *IOSR J Pharm* 2019; 9: 47–51.
331. Nhut P, An TN, Minh LV, et al. Phytochemical screening of *Allium Tuberosum* Rottler. ex Spreng as food spice. *IOP Conf Ser Mater Sci Eng* 2020; 991: 012021.
332. Eldahshan OA, Abdel-Daim MM. Phytochemical study, cytotoxic, analgesic, antipyretic and anti-inflammatory activities of *Strychnos nuxvomica*. *Cytotechnology* 2015; 67: 831–844.
333. Mittal M, Gupta N, Parashar P, et al. Phytochemical evaluation and pharmacological activity of

- syzygium aromaticum: A comprehensive review. *Int J Pharm Pharm Sci* 2014; 6: 67–72.
334. Hanani E, Prastiwi R, Karlina L, et al. Indonesian *Mirabilis jalapa* Linn.: A Pharmacognostical and Preliminary Phytochemical Investigations. *Pharmacogn J* 2017; 9: 683–688.
335. Khan I, AbdElsalam NM, Fouad H, et al. Application of Ethnobotanical Indices on the Use of Traditional Medicines against Common Diseases. *Evid Based Complement Alternat Med* 2014; 2014: 635371.
336. Shan S, Huang X, Shah M, et al. Evaluation of Polyphenolics Content and Antioxidant Activity in Edible Wild Fruits. *BioMed Res Int* 2019; 2019: 1–11.
337. Raghavendra MP, Satish S, Raveesha KA. Phytochemical analysis and antibacterial activity of *Oxalis corniculata*; a known medicinal plant. *My Sci* 2006; 1: 72–78.
338. Kaushik P, Kaushik D, Khokra SL. Ethnobotany and phytopharmacology of *Pinus roxburghii* Sargent: a plant review. *J Integr Med* 2013; 11: 371–376.
339. Bahadori MB, Sarikurku C, Kocak MS, et al. *Plantago lanceolata* as a source of health-beneficial phytochemicals: Phenolics profile and antioxidant capacity. *Food Biosci* 2020; 34: 100536.
340. C. Karkanis A, Fernandes Â, Vaz J, et al. Chemical composition and bioactive properties of *Sanguisorba minor* Scop. under Mediterranean growing conditions. *Food Funct* 2019; 10: 1340–1351.
341. Zhao Z, He X, Zhang Q, et al. Traditional Uses, Chemical Constituents and Biological Activities of Plants from the Genus *Sanguisorba* L. *Am J Chin Med* 2017; 45: 199–224.
342. Gao J, Sun C, Yang J, et al. Evaluation of the hepatoprotective and antioxidant activities of *Rubus parvifolius* L. *J Zhejiang Univ Sci B* 2011; 12: 135–142.
343. Cheng X, Qin J, Zeng Q, et al. Taraxasterane-Type Triterpene and Neolignans from *Geum japonicum* Thunb. var. *chinense* F. Bolle. *Planta Med* 2011; 77: 2061–2065.
344. Fu H, Mu X, Wang P, et al. Fruit quality and antioxidant potential of *Prunus humilis* Bunge accessions. *PLoS One* 2020; 15: e0244445.
345. Lu Q, Ma R, Yang Y, et al. *Zanthoxylum nitidum* (Roxb.) DC: Traditional uses, phytochemistry, pharmacological activities and toxicology. *J Ethnopharmacol* 2020; 260: 112946.
346. Zhang M, Wang J, Zhu L, et al. *Zanthoxylum bungeanum* Maxim. (Rutaceae): A Systematic Review of Its Traditional Uses, Botany, Phytochemistry, Pharmacology, Pharmacokinetics, and Toxicology. *Int J Mol Sci* 2017; 18: 2172.
347. Prakash B, Singh P, Mishra PK, et al. Safety assessment of *Zanthoxylum alatum* Roxb. essential oil, its antifungal, antiaflatoxin, antioxidant activity and efficacy as antimicrobial in preservation of *Piper nigrum* L. fruits. *Int J Food Microbiol* 2012; 153: 183–191.
348. Khatun A, Rahman M, Jahan S. Preliminary phytochemical, cytotoxic, thrombolytic and antioxidant activities of the methanol extract of *Murraya exotica* Linn. leaves. *Orient Pharm Exp Med* 2014; 14: 223–229.
349. Solanke SB. Phytochemical Information and Pharmacological Activities of Eggplant (*Solanum Melongena* L.): A Comprehensive Review. *EAS J Pharm Pharmacol* 2019; 1: 103–114.
350. Ohtsuki T, Miyagawa T, Koyano T, et al. Isolation and structure elucidation of flavonoid glycosides from *Solanum verbascifolium*. *Phytochem Lett* 2010; 3: 88–92.
351. Son YO, Kim J, Lim JC, et al. Ripe fruits of *Solanum nigrum* L. inhibits cell growth and induces apoptosis in MCF-7 cells. *Food Chem Toxicol* 2003; 41: 1421–1428.
352. Tekuri SK, Pasupuleti SK, Kranthi KK, et al. Phytochemical and pharmacological activities of *Solanum surattense* Burm. f.—A review. *J App Pharm Sci* 2019; 9: 126–136.
353. Alizadeh A, Moshiri M, Alizadeh J, et al. Black henbane and its toxicity - a descriptive review. *Avicenna J Phytomed* 2014; 4: 297–311.
354. Martínez-Valverde I, Periago MJ, Provan G, et al. A. Phenolic compounds, lycopene and antioxidant activity in commercial varieties of tomato (*Lycopersicon esculentum*): Phenolics, lycopene and antioxidant activity in tomatoes. *J Sci Food Agric* 2002; 82: 323–330.
355. Zaidi A, Bukhari S, Khan F, et al. Ethnobotanical, phytochemical and pharmacological aspects of *daphne mucronata* (thymeleaceae). *Trop J Pharm Res* 2015; 14: 1517–1523.
356. Zheng CJ, Zhao XX, Ai HW, et al. Therapeutic effects of standardized *Vitex negundo* seeds extract on complete Freund's adjuvant induced arthritis in rats. *Phytomedicine* 2014; 21: 838–846.
357. Usman H, Abdulrahman FI, A.A.L. Phytochemical and Antimicrobial Evaluation of *Tribulus terrestris* L. (Zygophyllaceae). Growing in Nigeria. *Res J Biol Sci* 2007; 2: 244–247.
358. Rashid U, Khan MR, Jan S, et al. Assessment of phytochemicals, antimicrobial and cytotoxic activities of extract and fractions from *Fagonia olivieri* (Zygophyllaceae). *BMC Complement*

- Altern Med* 2013; 13: 167.
359. Rashid U, Khan MR, Sajid M. Antioxidant, anti-inflammatory and hypoglycemic effects of *Fagonia olivieri* DC on STZ-nicotinamide induced diabetic rats - In vivo and in vitro study. *J Ethnopharmacol* 2019; 242: 112038.
360. Abbasi BH, Khan T, Khurshid R, et al. UV-C mediated accumulation of pharmacologically significant phytochemicals under light regimes in in vitro culture of *Fagonia indica* (L.). *Sci Rep* 2021; 11: 679.
361. Joshi RK. Volatile Constituents of *Emilia sonchifolia* from India. *Nat Prod Commun* 2018; 13: 1355–1356.
362. Kamboj A, Saluja AK. Phytopharmacological review of *Xanthium strumarium* L. (Cocklebur). *Int J Green Pharm* 2010; 4: 129–139.
363. Ahmad H, Sehgal S, Mishra A, et al. *Mimosa pudica* L. (Laajvanti): An overview. *Pharmacogn Rev* 2012; 6: 115–124.
364. Ting YC, Ko HH, Wang HC, et al. Biological evaluation of secondary metabolites from the roots of *Myrica adenophora*. *Phytochemistry* 2014; 103: 89–98.
365. Suhagia B, Rathod I, Sindhu S. *Sapindus mukorossi* (areetha): An overview. *Int J Pharm Sci Res* 2011; 2: 1905–1913.
366. Daboriya V, Kumar S, Singh S. Antibacterial activity and phytochemical investigations on *nicotiana plumbaginifolia* viv. (wild tobacco). *Rom J Biol -Plant Biol* 2010; 55: 135–142.
367. Ikenaga T, Handayani R, Oyama T. Steroidal saponin production in callus cultures of *Solanum aculeatissimum* Jacq. *Plant Cell Rep* 2000; 19: 1240–1244.
368. Pandey S, Sah SP, Sah ML, et al. An antioxidant potential of hydromethanolic extract of *Urtica parviflora* Roxb. *J Basic Clin Pharm* 2010; 1: 191–195.
369. Kalt FR, Cock IE. Gas chromatography-mass spectroscopy analysis of bioactive petalostigma extracts: Toxicity, antibacterial and antiviral activities. *Pharmacogn Mag* 2014; 10: S37–S49.
370. Singab AN, Youssef FS, Ashour ML, et al. The genus *Eremophila* (Scrophulariaceae): an ethnobotanical, biological and phytochemical review. *J Pharm Pharmacol* 2013; 65: 1239–1279.
371. Al-Abd NM, Mohamed Nor Z, Mansor M, et al. Antioxidant, antibacterial activity, and phytochemical characterization of *Melaleuca cajuputi* extract. *BMC Complement Altern Med* 2015; 15: 385.
372. Banbury LK, Shou Q, Renshaw DE, et al. Compounds from *Geijera parviflora* with prostaglandin E2 inhibitory activity may explain its traditional use for pain relief. *J Ethnopharmacol* 2015; 163: 251–255.
373. Al-Snafi A. A review on *Dodonaea viscosa*: A potential medicinal plant. *IOSR J Pharm* 2017; 7: 10–21.
374. Dapar MLG, Meve U, Liede-Schumann S, et al. Ethnomedicinal plants used for the treatment of cuts and wounds by the Agusan Manobo of Sibagat, Agusan del Sur, Philippines. *Ethnobot Res Appl* 2020; 19: 1–18.