

Global, regional, and national consumption of animal-source foods between 1990 and 2018: findings from the Global Dietary Database



Victoria Miller, Julia Reedy, Frederick Cudhea, Jianyi Zhang, Peilin Shi, Josh Erndt-Marino, Jennifer Coates, Renata Micha, Patrick Webb, Dariush Mozaffarian, on behalf of the Global Dietary Database



Summary

Background Diet is a major modifiable risk factor for human health and overall consumption patterns affect planetary health. We aimed to quantify global, regional, and national consumption levels of animal-source foods (ASF) to inform intervention, surveillance, and policy priorities.

Methods Individual-level dietary surveys across 185 countries conducted between 1990 and 2018 were identified, obtained, standardised, and assessed among children and adults, jointly stratified by age, sex, education level, and rural versus urban residence. We included 499 discrete surveys (91.2% nationally or subnationally representative) with data for ASF (unprocessed red meat, processed meat, eggs, seafood, milk, cheese, and yoghurt), comprising 3.8 million individuals from 134 countries representing 95.2% of the world population in 2018. We used Bayesian hierarchical models to account for differences in survey methods and representativeness, time trends, and input data and modelling uncertainty, with five-fold cross-validation.

Findings In 2018, mean global intake per person of unprocessed red meat was 51 g/day (95% uncertainty interval [UI] 48–54; region-specific range 7–114 g/day); 17 countries (23.9% of the world's population) had mean intakes of at least one serving (100 g) per day. Global mean intake of processed meat was 17 g/day (95% UI 15–21 g/day; region-specific range 3–54 g/day); seafood, 28 g/day (27–30 g/day; 12–44 g/day); eggs, 21 g/day (18–24 g/day; 6–35 g/day); milk 88 g/day (84–93 g/day; 45–185 g/day); cheese, 8 g/day (8–10 g/day; 1–34 g/day); and yoghurt, 20 g/day (17–23 g/day; 7–84 g/day). Mean national intakes were at least one serving per day for processed meat (≥ 50 g/day) in countries representing 6.9% of the global population; for cheese (≥ 42 g/day) in 2.3%; for eggs (≥ 55 g/day) in 0.7%; for milk (≥ 245 g/day) in 0.3%; for seafood (≥ 100 g/day) in 0.8%; and for yoghurt (≥ 245 g/day) in less than 0.1%. Among the 25 most populous countries in 2018, total ASF intake was highest in Russia (5.8 servings per day), Germany (3.8 servings per day), and the UK (3.7 servings per day), and lowest in Tanzania (0.9 servings per day) and India (0.7 servings per day). Global and regional intakes of ASF were generally similar by sex. Compared with children, adults generally consumed more unprocessed red meat, seafood and cheese, and less milk; energy-adjusted intakes of other ASF were more similar. Globally, ASF intakes (servings per week) were higher among more-educated versus less-educated adults, with greatest global differences for milk (0.79), eggs (0.47), unprocessed red meat (0.42), cheese (0.28), seafood (0.28), yoghurt (0.22), and processed meat (0.21). This was also true for urban compared to rural areas, with largest global differences (servings per week) for unprocessed red meat (0.47), milk (0.38), and eggs (0.20). Between 1990 and 2018, global intakes (servings per week) increased for unprocessed red meat (1.20), eggs (1.18), milk (0.63), processed meat (0.50), seafood (0.44), and cheese (0.14).

Interpretation Our estimates of ASF consumption identify populations with both lower and higher than optimal intakes. These estimates can inform the targeting of intervention, surveillance, and policy priorities relevant to both human and planetary health.

Funding Bill & Melinda Gates Foundation and American Heart Association.

Copyright © 2022 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

Introduction

Diet is a major modifiable risk factor for maternal and child health, undernutrition, and non-communicable diseases (NCDs) worldwide.¹ The food that humans consume is also a major determinant of environmental sustainability, with impacts on land, freshwater, energy use, greenhouse gas emissions, deforestation, and

biodiversity loss.^{2,3} Among the different food groups that influence both human and planetary health, perhaps none is more relevant than animal-source foods (ASF).⁴ ASF are diverse and heterogeneous, including unprocessed red meat, processed meat, poultry, eggs, seafood, milk, cheese, and yoghurt. These foods often contain high and bioavailable contents of important nutrients,

Lancet Planet Health 2022;
6: e243–56

Friedman School of Nutrition Science and Policy, Tufts University, Boston, MA, USA (V Miller PhD, J Reedy MS, F Cudhea PhD, J Zhang PhD, P Shi PhD, J Erndt-Marino PhD, J Coates PhD, R Micha PhD, Prof P Webb PhD, Prof D Mozaffarian MD); Department of Food Science and Nutrition, University of Thessaly, Volos, Greece (R Micha)

Correspondence to:
Dr Victoria Miller, Friedman School of Nutrition Science and Policy, Tufts University, Boston MA 02111, USA
victoria.miller@tufts.edu

Research in context

Evidence before this study

We systematically searched PubMed, Embase, Web of Science, LILACS, African Index Medicus, and the Southeast Asia Index Medicus to identify studies reporting nationally or subnationally representative estimates of individual-level consumption of seven animal-source foods (ASF). We included 1248 studies conducted between 1980 and 2016 using 24-h recalls, food frequency questionnaires, or short standardised questionnaires. When national or subnational individual-level surveys were not identified for a country, we searched for individual-level surveys from large cohorts, the WHO Global Infobase, and the WHO Stepwise Approach to Surveillance database. Household budget surveys were used on the rare occasion when individual-level dietary surveys were not identified for a populous country. We excluded surveys focused on special populations or cohorts. Using Bayesian hierarchical modelling methods, we estimated global, regional, and national intakes of ASF by age, sex, education, urbanicity, and time between 1990 and 2018.

We identified one previous global analysis of ASF consumption, which was limited to a few ASF subtypes, used crude national

estimates of food availability or expenditure data to estimate individual-level intakes, and did not report consumption by important socioeconomic factors.

Added value of this study

This study provides a comprehensive picture of consumption of total ASF, unprocessed red meat, processed meat, seafood, egg, milk, cheese, and yoghurt consumption in 185 countries among children (aged ≤ 19 years) and adults (≥ 20 years). It includes the first global estimates of mean ASF intakes by education level and urbanicity. We also present trends in ASF consumption over three decades. To our knowledge, this is the most current and comprehensive study on global individual-level consumption of ASF.

Implications of all the available evidence

This study highlights regions, countries, and population strata with both lower and higher than optimal ASF intakes. The findings can inform the targeting of intervention, surveillance, and policy priorities relevant to both human and planetary health

for example vitamin A, folic acid, calcium, iodine, iron, zinc, essential fatty acids, and protein,⁵⁻⁷ are the only natural dietary sources of vitamins B12 and D,^{8,9} and can have higher levels of some nutrients (eg, vitamin A, folate, vitamin B12, calcium, iron, and zinc) than plant-source foods.¹⁰ These characteristics make ASF useful for improving nutrition in vulnerable populations such as infants, young children, adolescents, women of reproductive age, pregnant and lactating women, and older adults, as well as very poor communities in low-income and middle-income countries.^{6,11-13} Different ASF might also have important adverse (eg, processed meats) or beneficial (eg, seafood and yoghurt) effects on NCDs, in particular cardiovascular disease, diabetes, and cancer.^{14,15} Impacts on planetary health are just as important, including increased greenhouse gas emissions, land and energy use, and acidification and eutrophication, and also vary according to the type of ASF,¹⁶⁻¹⁸ production method, and the suitability of the specific ASF to the local ecosystem in which it is produced.^{19,20}

Based on these important and heterogeneous effects of ASF on both human and planetary health, it is crucial to understand the patterns and distributions of their consumption globally, not only across countries but also within population subgroups such as by age, sex, education level, and urban versus rural residence. However, such distributions around the world are not well established. Previous reports are not up-to-date,^{21,22} reported on only a few ASF subtypes,²³ used crude national estimates of food availability or expenditure²³ that may not reflect dietary intakes,²⁴ or are focused on select countries, regions, or age groups.²⁵⁻²⁹ Furthermore,

global consumption levels among children or by education attainment and urban versus rural residence have not been previously reported.

To address these key knowledge gaps and provide the best current evidence on dietary intakes of key ASF worldwide, we aimed to systematically collect and evaluate global data from national and subnational individual-level dietary surveys using comparable and standardised methods and developed and applied Bayesian hierarchical modelling methods to estimate consumption of ASF by world region, country, age, sex, education level, and urbanicity, alongside changes in time between 1990 and 2018.

Methods

Data sources

The Global Dietary Database (GDD) is an international collaborative effort to produce comprehensive and comparable estimates of dietary intakes of major foods and nutrients in 185 countries and territories. Details on methods and the standardised data collection protocol have been described.^{21,22,30-34} Briefly, we performed systematic searches for individual-level dietary surveys in countries and territories globally, as well as extensive personal communications with researchers and government authorities throughout the world, inviting them to be corresponding members of the GDD. The results of our search strategy by dietary factor, time, and region have been detailed.³⁰ Surveys were prioritised if nationally or subnationally representative and using individual-level dietary assessments with standardised 24-h recalls, food frequency

questionnaires, or short standardised questionnaires (eg, Demographic Health Survey questionnaires). When national or subnational individual-level surveys were not identified for a country, we searched for individual-level surveys from large cohorts, the WHO Global Infobase, and the WHO Stepwise Approach to Surveillance database. Household budget surveys were used on the rare occasion when individual-level dietary surveys were not identified for a populous country. We excluded surveys focused on special populations (eg, pregnant or nursing women, or individuals with a specific disease) or cohorts (eg, a specific occupation or dietary pattern).

The final GDD model incorporated 1248 dietary surveys representing 188 countries and 99·0% of the global population in 2018. Of these surveys, 499 (40%) reported data for ASF including unprocessed red meat, total processed meat (ie, any meat, including poultry, that was cured, smoked, dried, or chemically preserved, excluding seafood and eggs), seafood, egg, milk, cheese, and yoghurt consumption (appendix pp 9–11). Based on the original focus on factors with potential causal associations with NCDs, the data collection did not include poultry, which we hope to update in future iterations. The 499 surveys included 3·8 million individuals from 134 countries, representing 95·2% of the global population. Most surveys were nationally or subnationally representative (91·2%), collected at the individual-level (80·7%), and included data for children and adolescents (93·4%), adults (76·3%), by rural or urban residence status (66·5%), and by education level (50·2%).

Data extraction

For each survey, we extracted data using standardised methods on survey characteristics and dietary metrics, units, mean, and standard deviation of consumption, by age, sex, education level, and urban or rural residence.³¹ Data were assessed for extraction errors and for plausibility using standardised algorithms, and survey quality by evaluating evidence for selection bias, sample representativeness, response rate, and validity of diet assessment method (appendix pp 3–5). Measurement comparability across surveys was maximised by using a standardised data analysis approach that leveraged the average of all days of dietary assessment to quantify mean individual-level intakes; used harmonised dietary factor definitions and units of measure across surveys; and adjusted for total energy using age-specific energy intakes to reduce measurement error and account for differences in body size, metabolic efficiency, and physical activity (appendix pp 3–8). All intakes are reported adjusted to 700 kcal/day for ages <1 year, 1000 kcal/day for ages 1–<2 years, 1300 kcal/day for ages 2–5 years, 1700 kcal/day for ages 6–10 years, 2000 kcal/day for ages 11–74 years, and 1700 kcal/day for ages ≥75 years.

Data modelling

To incorporate and address differences in data comparability, and sampling uncertainty, we used a Bayesian model with a nested hierarchical structure (with random effects by country, region, and globally) to estimate the mean consumption level of each ASF and its statistical uncertainty for each of the 264 population strata across 185 countries and each year between 1990 and 2018. Stratifying factors included age (<1, 1–2, 3–4, 5–9, 10–14, 15–19, 20–24, 25–29, 30–34, 35–39, 40–44, 45–49, 50–54, 55–59, 60–64, 65–69, 70–74, 75–79, 80–84, 85–89, 90–94, and ≥95 years), sex, education (≤6 years of education, >6 years to <12 years, or ≥12 years), and urbanicity (urban or rural), jointly stratified. For each dietary factor, primary inputs were the survey-level quantitative data (by country, time, age, sex, education level, and urban or rural-status); survey characteristics (dietary assessment method, type of dietary metric); and country-year-specific covariates (appendix pp 12–13). The model included overdispersion of study-level variance for surveys that were not nationally representative or not stratified by smaller age groups (≤10 years), sex, education level, or urbanicity. Uncertainty of each stratum-specific estimate was quantified using 4000 iterations to determine posterior distributions of consumption jointly by country, year, and demographic subgroup. We computed the median intake and the 95% uncertainty interval (UI) for each stratum as the 50th, 2·5th and 97·5th percentiles of the 4000 draws, respectively. Validity was assessed by five-fold cross-validation (randomly omitting 20% of the raw survey data, run five times), comparing predicted versus observed intakes; as well by assessment of implausible estimates and visual assessment of global and national mean intakes using heat maps. A second time component Bayesian model was used to strengthen time trend estimates for dietary factors with corresponding food or nutrient availability data (FAO Food Balance Sheet³⁵ and Global Expanded Nutrient Supply³⁶). The model incorporated country-level intercepts and slopes, along with their correlation that is estimated across countries. This model is commonly referred to as a varying slopes model structure and leverages two-dimensional partial pooling between intercepts and slopes to regularise all parameters and minimise overfitting risk.^{37,38} The final presented results are a combination of these two separate Bayesian models, as specified in detail in appendix pp 14–22.

See Online for appendix

Statistical analysis

Global, regional, national, and within-country population subgroup intakes and their uncertainty were calculated as population-weighted averages using all 4000 posterior predictions for each of the 264 demographic strata in each country-year. Population weights for each year were derived from the United Nations Population Division,³⁹ supplemented with data for education and urban versus rural status from Barro and Lee⁴⁰ and the United Nations.⁴¹ Intakes were calculated as both grams per day (g/day)

| | World | Southeast and East Asia* | Central or Eastern Europe and Central Asia† | High-income countries | Latin America and Caribbean | Middle East and North Africa | South Asia | Sub-Saharan Africa |
|-----------------------------|--------------|--------------------------|---|-----------------------|-----------------------------|------------------------------|-------------|--------------------|
| Unprocessed red meat | | | | | | | | |
| All ages | 51 (48–54) | 87 (79–96) | 114 (101–126) | 45 (43–47) | 68 (64–72) | 36 (31–43) | 7 (7–8) | 24 (23–26) |
| Children | 40 (38–43) | 84 (76–93) | 93 (81–103) | 30 (29–31) | 65 (61–70) | 36 (30–43) | 7 (7–8) | 20 (19–22) |
| Adults | 56 (53–60) | 88 (80–98) | 121 (106–134) | 50 (48–52) | 70 (65–74) | 37 (32–44) | 7 (7–8) | 29 (27–32) |
| Processed meat | | | | | | | | |
| All ages | 17 (15–21) | 13 (10–17) | 54 (45–64) | 30 (28–32) | 37 (32–43) | 19 (11–31) | 3 (1–12) | 12 (6–26) |
| Children | 18 (15–23) | 26 (20–34) | 44 (35–56) | 25 (24–27) | 37 (31–44) | 18 (10–31) | 3 (1–13) | 12 (5–27) |
| Adults | 17 (15–20) | 8 (6–12) | 57 (48–67) | 31 (29–34) | 37 (32–44) | 19 (11–32) | 3 (1–12) | 12 (6–25) |
| Seafood | | | | | | | | |
| All ages | 28 (27–30) | 44 (40–48) | 30 (26–35) | 25 (23–27) | 22 (20–24) | 23 (19–27) | 12 (11–13) | 31 (28–34) |
| Children | 21 (20–23) | 32 (30–35) | 17 (14–20) | 11 (11–12) | 17 (15–19) | 17 (14–21) | 10 (9–11) | 30 (27–33) |
| Adults | 32 (30–34) | 48 (44–53) | 35 (30–41) | 29 (27–32) | 24 (22–27) | 26 (22–32) | 13 (12–14) | 32 (29–36) |
| Eggs | | | | | | | | |
| All ages | 21 (18–24) | 35 (27–45) | 34 (28–42) | 18 (16–22) | 25 (23–27) | 30 (26–35) | 6 (5–7) | 6 (5–7) |
| Children | 17 (15–20) | 34 (27–43) | 30 (24–37) | 14 (11–17) | 25 (23–27) | 29 (25–34) | 6 (5–7) | 5 (4–5) |
| Adults | 23 (20–27) | 35 (28–46) | 35 (29–44) | 20 (17–24) | 25 (23–27) | 30 (26–36) | 6 (5–7) | 8 (6–10) |
| Milk | | | | | | | | |
| All ages | 88 (84–93) | 45 (41–51) | 145 (129–165) | 185 (173–201) | 150 (140–161) | 106 (87–131) | 84 (77–90) | 45 (41–49) |
| Children | 103 (98–109) | 75 (67–84) | 166 (145–193) | 252 (233–275) | 185 (170–201) | 127 (104–158) | 94 (85–103) | 46 (43–50) |
| Adults | 81 (77–84) | 35 (31–39) | 138 (122–157) | 165 (154–180) | 133 (122–145) | 93 (76–117) | 78 (84–85) | 44 (39–49) |
| Cheese | | | | | | | | |
| All ages | 8 (8–10) | 2 (1–3) | 34 (24–47) | 32 (28–36) | 13 (11–15) | 17 (13–23) | 1 (0–1) | 1 (1–2) |
| Children | 6 (6–7) | 2 (1–5) | 29 (20–42) | 28 (25–32) | 13 (11–16) | 14 (11–19) | 1 (0–1) | 1 (1–1) |
| Adults | 9 (8–11) | 1 (1–3) | 36 (25–50) | 33 (29–37) | 12 (10–14) | 19 (15–25) | 1 (0–1) | 1 (1–2) |
| Yoghurt | | | | | | | | |
| All ages | 20 (17–23) | 10 (7–14) | 84 (59–125) | 37 (32–46) | 18 (15–20) | 60 (46–80) | 7 (6–8) | 7 (5–10) |
| Children | 18 (15–21) | 11 (8–18) | 73 (50–110) | 37 (31–45) | 23 (20–27) | 52 (40–70) | 7 (6–9) | 6 (5–8) |
| Adults | 21 (18–25) | 9 (6–13) | 87 (60–132) | 37 (32–46) | 15 (13–18) | 64 (50–87) | 7 (5–8) | 8 (5–13) |

Data are mean intake (95% uncertainty interval) in g/day. All intakes are reported adjusted to 700 kcal/day for ages <1 year, 1000 kcal/day for ages 1–<2 years, 1300 kcal/day for ages 2–5 years, 1700 kcal/day for ages 6–10 years, 2000 kcal/day for ages 11–74 years, and 1700 kcal/day for ages ≥75 years. Children were defined as individuals younger than 20 years and adults were defined as individuals who were aged 20 years or older. Data are based on a Bayesian hierarchical model that incorporated up to 499 individual-level dietary surveys per dietary factors, and additional survey-level and country-level covariates, to estimate dietary consumption levels. Standardised serving size used for this analysis: unprocessed red meat=100 g; processed meat=50 g; seafood=100 g; egg=55 g; cheese=42 g; yoghurt=245 g; milk=245 g. * Referred to as Asia in previous Global Dietary Database reports. † Referred to as the Former Soviet Union in previous Global Dietary Database reports.

Table: Estimated dietary consumption (g/day) of animal-source foods among children and adults, by world region, in 2018

and servings per day or week, using standardised portion sizes. Spearman correlations between national mean intakes of different ASF were assessed. Changes in consumption between 1990 and 2018 were calculated at the stratum-specific prediction level to account for the full spectrum of uncertainty and standardised to the proportion of individuals within each stratum in 2018 to account for changes in demographics over time. Absolute and percentage differences in consumption between population subgroups were similarly calculated, using all 4000 posterior predictions for each stratum and population-weights in 2018. Differences in education level were computed as the difference for high (≥12 years) versus low (<6 years) education excluding moderate education (between 6 years and <12 years). For comparisons between sexes, education levels, urbanicity,

and time, difference thresholds were regarded as significant if the 95% UI did not include zero. Given the Bayesian estimates, no formal alpha level should be defined for statistical significance; and 95% UIs of each estimate should be considered as a guide.

Role of the funding source

The Bill & Melinda Gates Foundation contributed to study design during the grant application process; the funders otherwise had no role in data collection, data analysis, data interpretation, or writing of the report.

Results

In 2018, mean global consumption of unprocessed red meat was 51 g/day (95% UI 48–54), with a 16-fold variation across seven geographical regions (from 7 g/day in




Figure 1: National mean intake of unprocessed red meat (A) and processed meat (B) in 2018, all ages

17 (9%) of 185 countries consumed at least 100 g/day (equivalent to one serving per day) of unprocessed red meat, representing 1·8 billion people or 23·9% of the world population; 32 (17%) of 185 countries had intakes of at least 50 g/day (equivalent to one serving per day) of processed meat, representing 520 million people or 6·9% of the world population.

South Asia to 114 g/day in Central or Eastern Europe and Central Asia; table). Among the world's 25 most populous countries, mean national intakes were highest in Russia, South Africa, China, and Japan, and lowest in India, Bangladesh, Ethiopia, and DR Congo (range 3–188 g/day; figure 1). 17 (9%) of 185 countries, representing 1·8 billion people or 23·9% of the world's population, had mean consumption of at least one serving (100 g) per day.

Worldwide, mean intake of processed meat was 17 g/day (95% UI 15–21), with a 17-fold variation from highest (54 g/day; Central or Eastern Europe and Central Asia) to lowest (3 g/day; South Asia) regional intakes (table). Among the most populous countries, intakes ranged from 0 g/day to 62 g/day; highest in Germany, Russia, Philippines, and Brazil; and lowest in Bangladesh, India, Tanzania, and Turkey (figure 1). 32 (17%) of

185 countries, representing 520 million people or 6·9% of the world population, had mean intakes of at least one serving (50 g) daily.

Unprocessed red meat and processed meat intake were only moderately intercorrelated across countries (appendix p 28). Notable countries with much higher consumption of unprocessed red meat than processed meat included Russia (188 g/day vs 52 g/day), South Africa (147 g/day vs 17 g/day), and China (111 g/day vs 5 g/day; figure 1). In other countries, unprocessed red meat intake was lower than processed meat intake—eg, in the Philippines (25 g/day vs 45 g/day), DR Congo (12 g/day vs 30 g/day), Ethiopia (11 g/day vs 25 g/day), and Indonesia (26 g/day vs 37 g/day).

Mean global intake of seafood was 28 g/day (95% UI 27–30; table), with a 4-fold difference across regions




Figure 2: National mean intake of seafood (A) and egg (B) in 2018, all ages

Only three (2%) of 185 countries consumed at least 100 g/day (equivalent to one serving per day) of seafood, representing 6.2 million people or 0.8% of the world population; five (3%) of 185 countries had intakes of at least 55 g/day (equivalent to one serving per day) of egg, representing 6.2 million people or 0.7% of the world population.

(from 12 g/day in South Asia to 44 g/day in Southeast and East Asia). Generally, countries in Asia–Pacific and the Mediterranean basin had highest seafood intakes. Among the 25 most populous countries, mean national intakes were highest in Italy, Vietnam, Indonesia, and Japan (50–61 g/day), and lowest intakes were in Pakistan, Ethiopia, South Africa, and Turkey (4–6 g/day; figure 2). 87 (47%) of 185 countries had mean intakes of at least two servings (100 g each) per week, representing 4.0 billion people or 52.8% of the world population; and 16 of these countries had mean intakes of at least four servings per week (representing 37.5 million people, 5.0% of the world population). National seafood intake was not strongly correlated with unprocessed red meat or processed meat (appendix p 28).

Globally, mean consumption of eggs was 21 g/day (95% UI 18–24; table). Regional consumption was highest in Southeast and East Asia (35 g/day) and Central or Eastern Europe and Central Asia (34 g/day), and lowest in South Asia and Sub-Saharan Africa (both 6 g/day). Among the most populous countries, mean national intakes varied substantially from 1 g/day to 45 g/day; with lowest intakes in DR Congo, Tanzania, India, and Nigeria; and highest intakes in Vietnam, Japan, Russia, and China (figure 2). Only five (3%) of 185 countries consumed one or more servings of eggs (55 g) daily, representing 5.3 million people or 0.7% of the world's population. National egg intake was moderately correlated with unprocessed red meat and processed meat, but not with seafood (appendix p 28).




Figure 3: National mean intake of milk (A), cheese (B), and yoghurt (C) in 2018, all ages

Only seven (4%) of 185 countries had intakes of at least 245 g/day (equivalent to one serving per day) of milk, representing 18.8 million people or 0.3% of the world population. 13 (7%) of 185 countries had intakes of at least 42 g/day (equivalent to one serving per day) of cheese, representing 173 million people or 2.3% of the world population. One (1%) of 185 countries had intakes of at least 245 g/day (equivalent to one serving per day) of yoghurt, representing 7 million people or 0.09% of the world population; 26 (14%) of 185 countries had intakes of at least 70 g/day (equivalent to 2 servings/week) of yogurt, representing 539 million people or 7.1% of the world population.




Figure 4: Mean global and regional consumption of animal-source foods (A) in 2018 and change in consumption between 1990 and 2018 (B)
 Positive values in part (B) indicate greater consumption in 2018 than 1990. One serving of unprocessed red meat=100 g; total processed meat=50 g; seafood=100 g; egg=55 g; cheese=42 g; yoghurt=245 g; milk=245 g. Uncertainty intervals for the absolute change in consumption between 1990 and 2018 are provided in the appendix (p 90).

Worldwide, mean milk consumption was 88 g/day (95% UI 84–93), with 4-fold variation across regions (from 45 g/day to 185 g/day; table). Among populous countries, mean intakes were highest in Mexico, the UK, the USA, and France (188–206 g/day), and lowest in Nigeria, China, Bangladesh, and DR Congo (31–37 g/day; figure 3).

Mean global consumption of cheese was 8 g/day (8–10), with a 34-fold regional variation from 1 g/day in South Asia and sub-Saharan Africa to 34 g/day in Central or Eastern

Europe and Central Asia (table). Among populous countries, highest intakes were in the UK, France, Turkey, and the USA (30–39 g/day; figure 3); and lowest intakes in DR Congo, Bangladesh, India, and Tanzania (≤ 1 g/day).

Mean global yoghurt intake was 20 g/day (95% UI 17–23; table), with regional consumption being at least 3-fold higher in Central or Eastern Europe and Central Asia (84 g/day) and the Middle East and North Africa (60 g/day), and only a third as much as the global mean in South Asia and sub-Saharan Africa (7 g/day each). National intakes ranged substantially, from 0 g/day to 503 g/day. Among populous countries, national intakes were lowest in Indonesia, Bangladesh, Tanzania, and Thailand (2–6 g/day) and highest in Turkey, Russia, Iran, and France (77–112 g/day; figure 3).

Compared with meats, national consumption of cheese, yoghurt, and milk were much more strongly intercorrelated (appendix p 28). National yoghurt intake was greater than milk intake only in Iran (71 g/day vs 38 g/day); and cheese intake was greater than yoghurt intake in the USA (30 g/day vs 14 g/day; figure 3).

Total consumption of ASF was lowest in South Asia (<1 serving per day) and sub-Saharan Africa (~1 serving per day), intermediate in Southeast and East Asia and the Middle East and North Africa (2–3 servings per day), higher in Latin America and Caribbean and high-income countries (3–4 servings per day), and highest in Central or Eastern Europe and Central Asia (~5 servings per day; figure 4). Among the most populous countries, total ASF consumption was highest in Russia (5.8 servings per day), Germany (3.8 servings per day), and the UK (3.7 servings per day), and lowest in India (0.7 servings per day) and Tanzania (0.9 servings per day; appendix p 37).

Globally and regionally, the mean energy-adjusted intakes of most ASF were not appreciably different between women and men (appendix pp 30–36, 61). Exceptions globally were yoghurt and milk, with women consuming slightly more yoghurt and milk than men (yoghurt 0.09 servings per week [95% UI 0.05 to 0.14; milk: 0.11 servings per week [0.02 to 0.20]) and slightly less processed meat (–0.31 servings per week [–0.49 to –0.14]).

Globally, intakes of most ASF increased with age, but age trends varied considerably at regional and national levels (appendix pp 38–53). Adults globally consumed more unprocessed red meat, seafood, and cheese than did children (unprocessed red meat: 56 g/day [95% UI 53–60] in adults vs 40 g/day [38–43] in children; seafood: 32 g/day [30–34] vs 21 g/day [20–23]; cheese: 9 g/day [8–11] vs 6 g/day [6–7]), whereas children consumed more milk (81 g/day [77–84] vs 103 g/day [98–109]; table). Larger regional differences in intake between adults versus children were found for some foods, such as for unprocessed red meat in Central or Eastern Europe and Central Asia, sub-Saharan Africa, and high-income countries; for processed meat in high-income countries,

and Southeast and East Asia; for seafood in all regions except sub-Saharan Africa; and for milk in Southeast and East Asia and high-income countries.

On average, individuals with higher education (≥ 12 years) vs low education (< 6 years) consumed more ASF globally (figure 5, appendix pp 30–36, 70–78). In absolute servings, global differences by education were largest for milk (0.79 servings per week [95% UI 0.71–0.87]; 53.0% relative difference), followed by eggs (0.47 servings per week [0.36–0.60]; 50.6%), unprocessed red meat (0.42 servings per week [0.35–0.49]; 29.1%), cheese (0.28 servings per week [0.23–0.34]; 78.7%), seafood (0.28 servings per week [0.23–0.33]; 19.8%), yoghurt (0.22 servings per week [0.18–0.27]; 135.8%), and processed meat (0.21 servings per week [0.08–0.35]; 73.7%). In all regions, more educated individuals consumed more milk, except in Central or Eastern Europe and Central Asia and high-income countries where intake by education was more similar. The largest regional difference in absolute servings by higher education was seen for unprocessed red meat, seafood, and eggs in sub-Saharan Africa, and processed meat, cheese, and yoghurt in Central or Eastern Europe and Central Asia.

Compared with rural individuals, mean global intakes were higher among urban individuals for all ASF except processed meat (figure 5; appendix pp 30–36, 79–87). The largest global differences (in absolute intakes) were for unprocessed red meat (0.47 servings per week [95% UI 0.40–0.55]; 35.7% relative difference), milk (0.38 servings per week [0.30–0.46]; 23.9%), and eggs (0.20 servings per week [0.06–0.34]; 31.1%). The largest regional difference in absolute servings by urban residence was seen for processed meat, milk, and cheese in Latin America and Caribbean, unprocessed red meat in Central or Eastern Europe and Central Asia, seafood and eggs in sub-Saharan Africa, and yoghurt in the Middle East and North Africa.

Between 1990 and 2018, mean unprocessed red meat intake per person increased globally by 88.1%, equivalent to an additional 1.20 servings per week (95% UI 1.06 to 1.35; standardised to 2018 population distributions; figure 4; appendix pp 88–96). Notably, this global trend was due to increased intake in only three of the seven regions: Southeast and East Asia increased by 4.12 servings per week (3.66 to 4.64; 265.7% relative difference); Latin America and Caribbean increased by 1.29 servings per week (1.19 to 1.41; 57.9%); and sub-Saharan Africa increased by 0.06 servings per week (0.05 to 0.07; 26.3%). Intake decreased in Central or Eastern Europe and Central Asia, Middle East and North Africa, high-income countries, and South Asia. Among populous countries, absolute increases were largest in China (increase of 5.89 servings per week [5.18 to 6.69]; 312.5%), Japan (3.53 servings per week [3.36 to 3.71]; 128.8%), Brazil (2.45 servings per week [2.22 to 2.72]; 94.3%), and Mexico (1.18 servings per week [1.08 to 1.28]; 55.7%). Absolute decreases were




Figure 5: Mean global and regional absolute animal-source food intake difference by education level (A) and place of residence in 2018 (B)

Positive values in part (A) indicate greater consumption in high-education level individuals, and in part (B) indicate greater consumption in individuals in urban areas. One serving of unprocessed red meat=100 g; total processed meat=50 g; seafood=100 g; egg=55 g; cheese=42 g; yoghurt=245 g; milk=245 g. Uncertainty intervals for the absolute change in consumption between 1990 and 2018 are provided in the appendix (p 81). Absolute difference by education level was computed as the difference at the stratum-level and aggregated to the global and regional mean differences using weighted population proportions for low (< 6 years) and high education levels (≥ 12 years) only (excludes education level between 6 years and < 12 years).

largest in Russia (−2.00 servings per week [−2.34 to −1.30]; −14.0%), Germany (−1.19 servings per week [−1.33 to −1.07]; −26.3%), Iran (−1.04 servings per week [−1.16 to −0.92]; −47.7%), and France (−0.96 servings per week [−1.08 to −0.84]; −22.1%).

Between 1990 and 2018, mean processed meat intake increased globally by 152.8% (0.50 servings per week [95% UI 0.27 to 0.71]), with increases in most regions

(0.35–1.40 servings per week) except in the Middle East and North Africa and South Asia where intakes were stable (figure 4; appendix pp 88–96). Among populous countries, the largest increases were in the Philippines (3.94 servings per week [3.13 to 4.90]; 163.2%), Brazil (3.82 servings per week [2.82–4.91]; 186.7%), Indonesia (2.57 servings per week [1.15–4.36]; 410.4%), and Russia (2.54 servings per week [1.54–3.76]; 53.9%). Only two (8%) of 25 populous countries had decreases: Nigeria (–0.75 servings per week [–2.81 to –0.02]; –33.0%) and Mexico (–0.71 servings per week [–1.02 to –0.42]; –21.6%).

Global seafood consumption doubled (109.4%) between 1990 and 2018, increasing by 0.44 servings per week (95% UI 0.37–0.51; figure 4; appendix pp 88–96). Increases were seen in all regions except sub-Saharan Africa, with the largest absolute increase in Southeast and East Asia (1.30 servings per week [1.10 to 1.51]; 148.7%). Among populous countries, the largest increases occurred in Vietnam (3.22 servings per week [2.23 to 4.69]; 306.1%), Thailand (1.68 servings per week [1.17 to 2.42]; 176.2%), China (1.66 servings per week [1.41 to 1.94]; 167.3%), and Italy (1.38 servings per week [1.22 to 1.56]; 47.4%). The largest absolute decreases were in Tanzania (–8.01 servings per week [–9.62 to –6.55]; –81.1%), the Philippines (–3.22 servings per week [–3.53 to –2.94]; –65.2%), and Japan (–2.21 servings per week [–2.42 to –2.01]; –38.6%).

Globally, egg consumption per person increased by 141.4% between 1990 and 2018, rising by 1.18 servings per week (95% UI 0.94 to 1.50). Regional intakes increased by 2.89 servings per week in Southeast and East Asia (2.14 to 3.91; 345.5%) to 0.29 servings per week in South Asia (0.24 to 0.34; 54.1%; figure 4, appendix pp 88–96). Across populous countries, the greatest increases were in Vietnam (4.90 servings per week [3.63 to 6.56]; 564.3%), China (3.67 servings per week [2.62 to 5.10]; 410.2%), Mexico (2.57 servings per week [2.36 to 2.80]; 123.6%), and South Africa (1.28 servings per week [0.92 to 1.74]; 112.9%). Ethiopia had the largest decrease in egg intake (–0.58 servings per week [–0.70 to –0.48]; –41.0%), followed by Germany (–0.48 servings per week [–0.53 to –0.43]; –13.8%), Tanzania (–0.46 servings per week [–0.56 to –0.37]; –48.0%), and France (–0.18 servings per week [–0.20 to –0.15]; –11.0%).

Changes over time in intakes of milk, cheese, and yoghurt are summarised in the appendix (pp 29, 88–96). Milk consumption doubled globally, cheese consumption increased by 56.0%, and yoghurt consumption did not change significantly (figure 4).

Discussion

Our systematic analysis, based on GDD data evaluating 499 largely national, individual-level dietary surveys, provides new estimates of global, regional, and national consumption of ASF between 1990 and 2018. Several

aspects of these findings are novel, including the results for children and the overall population stratified by education level and by urban or rural residence. These are also, to our knowledge, the first worldwide estimates for intakes of cheese and yoghurt. The overall findings have important implications for both human and planetary health.

Worldwide, unprocessed red meat intake per person increased by 88.1% over this period (increases would be larger further accounting for population growth), but almost entirely due to increases in Southeast and East Asia and Latin America and Caribbean; modest decreases were found for most other regions. Limiting intake of unprocessed red meat is nutritionally recommended in many countries through national food-based dietary guidelines due to links with cardiovascular disease, diabetes, and certain cancers.^{14,15} At the same time, livestock production using current technologies is the single greatest contributor to greenhouse gas emissions from the agriculture sector (5.8% of global greenhouse gas in 2016).⁴² Our results suggest that greater unprocessed red meat intakes over time in particular countries with high populations—especially China, Japan, Brazil, and Mexico—run counter to these recommendations for moderation. Consistent with our findings, the China National Nutrition Surveys/China Health and Nutrition Survey and the FAO Food Balance Sheet data show that red meat consumption and availability substantially increased over time, largely due to increased pork consumption.^{35,43} Our findings also show higher intakes among more educated and urban individuals, suggesting that continuing global advances in education and urbanicity might further exacerbate global trends.

Intake of processed meat also increased, by 152.8% globally, with increases in most world regions. Notably, intakes of unprocessed red meat and processed meat were only moderately correlated across countries, and processed meat intake was not associated with urbanicity globally. These findings suggest differential drivers—and potential levers for action—for unprocessed versus processed meats, with urbanisation having less influence on processed meat consumption compared with socioeconomic factors.^{44,45} This factor is an important consideration for influencing total meat consumption globally, given the generally stronger links with NCDs of processed meats.

Seafood has been shown to be an important component of a healthy diet, including for childhood brain development^{46,47} and cardiovascular health in adults.^{48,49} Global seafood consumption doubled after 1990, and by 2018 more than half of countries had mean intakes of two or more servings per week. However, seafood intake remained low in many South Asian, Latin American and Caribbean, and Middle Eastern and North African countries, with only small improvements over time. Adjusted for energy, seafood intake was also generally much lower among children than adults in most world

regions, and among those with lower education or rural residence. These new results suggest important disparities that must be addressed, together with sustainable approaches to increase seafood production,¹⁶ to ensure adequate health benefits for all.

Global dietary intakes also increased for eggs (by 141·4%), milk (98·6%), and cheese (56·0%), whereas yoghurt intake was stable. However, absolute intakes remained relatively low in most countries, with few countries reaching intakes of one daily serving for each. National intercorrelations of eggs and dairy foods were relatively high, and generally consistent with population prevalence of lactase deficiency.⁵⁰ Consistent with high levels of lactase deficiency, some countries in Asia had high intakes of eggs only. Conversely, most sub-Saharan African countries had low intakes of eggs and dairy, with very little change between 1990 and 2018. These findings highlight the importance of strategies to augment intakes of these lower cost, more environmentally sustainable animal food sources of nutrients,^{16,17} while also accounting for genetically driven challenges in tolerance for dairy.

The *Eat–Lancet* Commission on Food, Planet, and Health's report proposed common targets for a healthy and environmentally sustainable diet, including limiting red meat, poultry, and egg consumption, and moderate levels of fish and dairy consumption.¹⁶ The impact of several of these targets for health versus sustainability may be quite different. For example, reducing unprocessed red meat intake to the suggested targets would have large impacts on sustainability but much smaller effects on health.⁴ Reducing dairy, poultry, and eggs to the targets would have smaller sustainability impacts⁴ and little health benefit,⁵¹ and some populations may benefit from increasing their currently low intakes of these foods closer to the targets.^{29,52,53} Our findings show that few countries in 2018 met the *Eat–Lancet* target for reduced total red meat (3·8% of countries, based on a target of ≤ 98 g/week of unprocessed red meat and processed meat combined [processed meat included processed poultry]), less than half for fish (43·2% of countries ≤ 196 g/week of seafood), and eggs (33·5% of countries ≤ 91 g/week), but most for dairy (83·8% of countries ≤ 250 g/day of milk, cheese, and yoghurt combined).

Previous studies of global ASF consumption include fewer current data, evaluated fewer ASF subtypes, and did not include children.^{21–23} Consistent with our findings, the Global Burden of Diseases, Injuries, and Risk Factors Study (GBD) study estimated that milk intake was highest in Australasia, western Europe, and North America, and lowest in South Asia and sub-Saharan Africa.²³ Our global mean estimates for unprocessed red meat and processed meat consumption were higher than the GBD study's estimates, although both studies generally found high meat consumption in North America, western Europe, and Latin America, and low consumption in south Asia and sub-Saharan Africa. Differences in estimated intakes and trends might be

attributable to the larger number of individual-level dietary surveys in the GDD, reliance on national food availability data as estimates of individual-level dietary data in the GBD, and differing modelling methods.

Our study has several strengths. Global ASF intakes were estimated using 499 dietary surveys, mostly nationally representative and assessing individual-level dietary intakes and including 3·8 million individuals from 134 countries representing 95·1% of the world's population. Data were harmonised using standardised protocols, with Bayesian hierarchical modelling to incorporate survey and country covariates and address heterogeneity, missingness, and sampling and modelling uncertainty. Findings were assessed over time and by important demographic characteristics including age, sex, education, and urban or rural residence.

Potential limitations warrant mention. Survey availability was limited for some (particularly low-income) countries, demographic groups, time periods and ASF subtype (eg, <200 surveys on cheese and yoghurt vs >400 on milk and unprocessed red meat), increasing uncertainty in these estimates. All types of dietary data are measured with some error, including from individual-level surveys as well as national food availability estimates. Additionally, the standardisation of available global data required certain decisions and assumptions about serving sizes, food group definitions, energy adjustment, and the disaggregation of household-level data when standardising the dietary surveys. However, we utilised rigorous methods developed over 14 years of work and carefully documented and detailed each survey's methods and standardisation process to maximise transparency in our methods. This iteration of the GDD did not collect information on poultry, an important ASF in many countries globally. Last, our analysis was limited to the consumption of ASF and does not describe trends in ASF production. Overall, these new results should be considered the best currently available, rather than perfect estimates of dietary intakes of ASF worldwide. In addition, our findings identify specific world regions and countries most urgently requiring well-conducted national surveys on individual-level intakes of ASF.

Global Dietary Database Corresponding Members

Pamela Abbott; Morteza Abdollahi, National Nutrition and Food Technology Research Institute (NNFTRI); SBMU; Parvin Abedi, Ahvaz Jundishapur University of Medical Sciences; Suhad Abumweis, The Hashemite University; Linda Adair; Swee Ai Ng; Mohannad Al Nsour, Eastern Mediterranean Public Health Network; EMPHNET; Ifitkhar Alam, BKUC.edu.pk; Nasser Al-Daghri, King Saud University; Nawal Al-Hamad, Ministry of Health; Suad Al-Hooti, Kuwait Institute for Scientific Research; Eman Alissa, King Abdulaziz University; Sameer Al-Zenki; Simon Anderson, University of Manchester; Karim Anzid, Cadi Ayyad University; Carukshi Arambepola, Faculty of Medicine, University of Colombo, Sri Lanka; Mustafa Arici, Hacettepe University Faculty of Medicine; Joanne Arsenault, University of California, Davis; Renzo Asciak; Lajos Biró; Noël Barengo, Herbert Wetheim College of Medicine; Simon Barquera, Instituto Nacional de Salud Publica; Murat Bas, Acibadem University; Wulf Becker; Sigrid Beer-Borst; Per Bergman, Livsmedelsverket; Sesikeran Boindala,

National Institute of Nutrition India; Pascal Bovet, Institute of Social and Preventive Medicine, University of Lausanne, Switzerland, and Ministry of Health, Seychelles; Debbie Bradshaw; Norikil Bukhary Ismail Bukhary, Ministry of Health (Malaysia); Kanitta Bundhamcharoen; Mauricio Caballero, Fundacion Infant; Neville Calleja, Directorate for Health Information & Research; Xia Cao; Mario Capanzana, Food and Nutrition Research Institute, Department of Science and Technology; Jan Carmikle, Senior Intellectual Property Office; Katia Castetbon, Institut de Veille Sanitaire; Michelle Castro, Departamento de Alimentação Escolar, Corazon Cerdana; Hsing-Yi Chang, National Health Research Institutes; Karen Charlton; Yu Chen, NYU School of Medicine; Shashi Chiplonkar, HC Jehangir Medical Research Institute, Pune India; Yoonsu Cho, Korea University; Khun-Aik Chuah; Simona Costanzo, IRCCS INM Neuromed; Melanie Cowan; Albertino Damasceno, Faculty of Medicine, Eduardo Mondlane University, Maputo, Mozambique; Saeed Dastgiri, Tabriz University of Medical Sciences; Stefaan De Henauw, Ghent University; Karin DeRidder, Belgian Public Health Institute; Eric Ding, Harvard School of Public Health; Rivera Dommarco; Rokiah Don; Charmaine Duante; Vesselka Duleva; Samuel Duran Agüero, Universidad San Sebastian, Chile; Veena Ekbote, Hirabai Cowasji Jehangir Medical Research Institute, Jehangir Hospital, Pune, India; Jalila El Ati, National Institute of Nutrition and Food Technology & SURVEN RL; Asmaa El Hamdouchi; Alison Eldridge, Nestle Research Center; Tatyana El-kour, World Health Organization; Ibrahim Elmadaf, University of Vienna; Helene Enghardt Barbieri; Alireza Esteghamati, Tehran University of Medical Sciences; Zohreh Etemad, Dutch National Institute for Public Health and the Environment (RIVM); Fariza Fadzil, Ministry of Health, Malaysia; Farshad Farzadfar; Mei Fen Chan; Anne Fernandez, Perdana University, Royal College of Surgeons in Ireland; Dulitha Fernando; Regina Fisberg, University of Sao Paulo, Brazil; Simon Forsyth, The University of Queensland School of Public Health; Edna Gamboa Delgado, Fundacion Cardiovascular de Colombia; Didier Garriguet, Statistics Canada; Jean-Michel Gaspoz; Dorothy Gauci; Marianne Geleijnse, Wageningen University; Brahmam Ginnela; Giuseppe Grosso, Integrated Cancer Registry of CT-ME-SR-EN; Idris Guessous, Geneva University Hospitals; Martin Gulliford, King's College London; Ingibjorg Gunnarsdottir; Wilbur Hadden; Aida Hadziomeragic, Institute of Public Health of Federation of Bosnia and Herzegovina; Christian Haerpfger; Jemal Haidar Ali, Addis Ababa University; Rubina Hakeem; Aminul Haque, University of Dhaka; Maryam Hashemian; Rajkumar Hemalatha, ICMR-National Institute of Nutrition; Sigrun Henjum, Oslo and Akerhus University College; Hristo Hinkov, National Center of Public Health and Analyses (NCPHA); Zaiton Hjdaud; Daniel Hoffman, Rutgers University; Beth Hopping; Anahita Houshiar-rad, National Nutrition & Food Technology Research Ins; Yao-Te Hsieh, Institute of Biomedical Sciences, Academia Sinica, Taipei, Taiwan; Shu-Yi Hung; Inge Huybrechts, International Agency for Research on Cancer; Nahla Chawkat Hwalla, American University of Beirut; Nayu Ikeda; Daniel Illescas-Zarate, National Institute of Public Health; Manami Inoue; Olof Jonsdottir; Hamid Jan Bin Jan Mohamed, Universiti Sains Malaysia; Chandrashekar Janakiram, Amrita School of Dentistry; Ranil Jayawardena, University of Colombo; Rajesh Jeewon, University of Mauritius; Nattinee Jitnarin; Lars Johansson; Ola Kally; Mirnalini Kandiah; Tilakavati Karupaiah, National University of Malaysia; Lital Keinan-Boker, Israel Center for Disease Control; Roya Kelishadi, Research Institute for Primordial Prevention of NCD, Isfahan University of Medical Sciences; Anuradha Khadilkar, Hirabai Cowasji Jehangir Medical Research Institute; Cho-il Kim, Korea Health Industry Development Institute; Eda Koksali; Jurgen Konig, University of Vienna, Department of Nutritional Sciences; Liisa Korkalo, University of Helsinki; Jeremy Koster, University of Cincinnati; Irina Kovalskys, ILSI (International Life Sciences Institute), Argentina; Anand Krishnan, All India Institute of Medical Sciences; Herculina Kruger, North-West University, Potchefstroom South Africa; Rebecca Kuriyan-Raj, St John's Research Institute; Sanghui Kweon; Carl Lachat, Ghent University, Belgium; Yuen Lai; Pulani Lanerolle, University of Colombo, Sri Lanka; Indu Waidyatilaka, University of Colombo, Sri Lanka; Avula Laxmaiah, National Institute of Nutrition, ICMR, Hyderabad; Catherine Leclercq; Meei-Shyuan Lee; Hae-Jeung Lee, Eulji University; J Lennert Veerman, The University of Queensland; Lydia Lera Marques, Unidad de Nutricion Publica-Professor Asociado; Yanping Li, Harvard School of Public Health; Jaana Lindström; Annie Ling; Nur Indrawaty Liputo, Andalas University; Patricio Lopez-Jaramillo, FOSCAL and UDES; Amy Luke, Loyola University Chicago; Widjaja Lukito; Nuno Lunet, Faculty of Medicine, University of Porto, Portugal; Elisabette Lupotto, CREA-Alimenti e Nutrizione; Guansheng Ma; Yi Ma; Zaleha Abdullah Mahdy, National University of Malaysia (UKM); Reza Malekzadeh, Digestive Disease Research Institute, Tehran University of Medical Sciences; Wan Manan, Universiti Sains Malaysia; Dirce Marchioni; Pedro Marques-Vidal, Lausanne University Hospital (CHUV); Yves Martin-Prevel, Institut de Recherche pour le Développement; Hajah Masni Ibrahim; Angie Mathee; Yasuhiro Matsumura, Bunkyo University, Faculty of Health and Nutrition; Paramita Mazumdar, Centre For Media Studies; Abba Mehio Sibai; Anjum Memon, Brighton and Sussex Medical School; Gert Mensink, Robert Koch Institute; Alexa Meyer, University of Vienna, Austria; Parvin Mirmiran, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences; Masoud Mirzaei, Yazd Cardiovascular Research Centre, Shahid Sadoughi University of Medical Sciences; Puneet Misra, All India Institute of Medical Sciences; Anoop Misra, Fortis CDOC Center for Excellence for Diabetes; Claudette Mitchell, University of the Southern Caribbean; Noushin Mohammadifard, Isfahan Cardiovascular Research Center, Cardiovascular Research Center, Isfahan University of Medical Sciences; Fatemeh Mohammadi-Nasrabadi, National Nutrition and Food Technology Research Institute; Zalilah Mohd Shariff, Universiti Putra Malaysia; Foong Ming Moy, University of Malaya; Abdulrahman Musaiger, Arab Center for Nutrition; Elizabeth Mwaniki, The Technical University of Kenya; Jannicke Myhre; Balakrishna Nagalla; Androniki Naska; Augustin Nawidimbasha Zeba, Institut de Recherche en sciences de la Sante; Shu Wen Ng, University of North Carolina at Chapel Hill; Le Tran Ngoan, Hanoi Medical University; Sina Noshad, Tehran University of Medical Sciences; Angelica Ochoa, Universidad de Cuenca; Marga Ocke, Dutch National Institute for Public Health and the Environment; RIVM; Jillian Odenkirk; Kyungwon Oh, Division of Health and Nutrition Survey; Mariana Oleas, Universidad Tecnica del Norte; Sonia Olivares, Institute of Nutrition and Food Technology (INTA), University of Chile; Philippos Orfanos; Johana Ortiz-Ulloa, Cuenca University; Johanna Otero, Fundacion Oftalmologica de Santander (FOSCAL); Marja-Leena Ovaskainen; Mohammadreza Pakseresh; Cristina Palacios, Florida International University; Pam Palmer; Wen-Harn Pan; Demosthenes Panagiotakos, Harokopio University; Rajendra Parajuli, McGill University; Myungsook Park; Gulden Pekcan; Stefka Petrova; Noppawan Piaseu, Mahidol University; Christos Pitsavos; Kalpagam Polasa, National Institute of Nutrition; Luz Posada, Universidad de Antioquia; Farhad Pourfarzi, Ardabil University of Medical Sciences; Alan Martin Preston, Univ Puerto Rico-Med Sci Dept Biochemistry; Ingrid Rached, Centro de Atencion Nutricional Antimano (CANIA); Ali Reza Rahbar; Colin Rehm; Almut Richter; Leanne Riley; Luz Maria Sánchez-Romero, National Institute of Public Health Mexico; Benoit Salanave; Nizal Sarrafzadegan, Isfahan Cardiovascular Research Center; Norie Sawada, Research Center for Cancer Prevention and Screening, National Cancer Center; Makiko Sekiyama, Graduate Program in Sustainability Science Global Leadership Initiative (GPSS-GLI), The University of Tokyo; Rusidah Selamat; Khadijah Shamsuddin, Universiti Kebangsaan Malaysia Medical Centre; Sangita Sharma, University of Alberta; Harri Sinkko; Isabelle Sioen; Ivan Sisa, USFQ; Sheila Skeaff, University of Otago; Laufey Steingrimsdottir; Tor Strand, University of Bergen; Milton Fabian Suarez-Ortega, University of Edinburgh; Sumathi Swaminathan, St John's Research Institute; Gillian Swan; Elzbieta Sygnowska; Maria Szabo; Lucjan Szponar, National Food and Nutrition Institute; Ilse Tan-Khouw; Heli Tapanainen, The National Institute for Health and Welfare (THL); Reema Tayyem, the Hashemite University; Bemnet Tedla; Alison Tedstone; Robert Templeton; Celine Termote, Bioversity International; Anastasia Thanopoulou, Diabetes Center, 2nd Department of Internal Medicine, Athens University, Greece; Holmfridur Thorgeirsdottir; Inga Thorsdottir; Dimitrios Trichopoulos; Antonia Trichopoulou; Shoichiro Tsugane; Aida Turrini; Coline van Oosterhout; Erkki Vartiainen; Suvi Virtanen, National Institute for Health and Welfare; Peter Vollenweider; Marieke Vossenaar, CeSSIAM in Guatemala;

Eva Warensjo Lemming, Risk and Benefit assessment Department, National Food Agency Sweden; Anna Waskiewicz, Department of CVD Epidemiology, Prevention and Health Promotion, Institute of Cardiology, Warsaw, Poland; Eveline Waterham; Lothar Wieler, Robert Koch Institute; Tizita Wondwossen, Addis Ababa University; Suh Wu; Roseyati Yaakub; Mabel Yap; Safiah Yusof; Sahar Zaghoul; Gábor Zajkás; Maria Zapata, CESNI; Khairul Zarina; Fatemeh Vida Zohoori, Teesside University

Contributors

VM, RM, PW, and DM conceptualised and designed the study. VM, JR, JC, and RM were involved in the data collection. VM, FC, JZ, PS, and JE-M conducted the analyses for the study. VM, JR, and FC verified the accuracy of the data. VM, PW, and DM drafted the manuscript. All the authors had access to and interpreted the data, read the final manuscript, reviewed it for important intellectual content and approved its submission. VM and DM are the guarantors of this work.

Declaration of interests

The investigators did not receive funding from a pharmaceutical company or other agency to write this report. JR, JZ, and PS report research funding from Nestlé, outside the submitted work. JC reports research funding from the Bill & Melinda Gates Foundation, outside the submitted work. RM reports grants from US National Institutes of Health, Nestlé, and Danone, and personal fees from Bunge, Development Initiatives, outside the submitted work. PW reports research grants and contracts from the US Agency for International Development and personal fees from the Global Panel on Agriculture and Food Systems for Nutrition, outside the submitted work. DM reports research funding from the US National Institutes of Health and the Gates Foundation; personal fees from GOED, Bunge, Indigo Agriculture, Motif FoodWorks, Amarin, Acasti Pharma, Cleveland Clinic Foundation, America's Test Kitchen, and Danone; scientific advisory board member for Brightseed, DayTwo, Elysium Health, Filtricine, HumanCo, and Tiny Organics; and chapter royalties from UpToDate, all outside the submitted work. All other authors declare no competing interests.

Data sharing

The modelled estimates of animal-source food intakes by population subgroup, country, region, and globe in 1990 and 2018 are available for download from the Global Dietary Database. Survey-level information and original data download weblinks are also provided for all public surveys; and survey-level microdata or stratum-level aggregate data are provided for direct download for all non-public surveys granted consent for public sharing by the data owner.

Editorial note: the *Lancet* Group takes a neutral position with respect to territorial claims in published maps and institutional affiliations.

References

- FAO and WHO. Sustainable healthy diets—guiding principles. 2019. <http://www.fao.org/3/ca6640en/ca6640en.pdf> (accessed June 30, 2021).
- Clark MTD, Tilman D. Comparative analysis of environmental impacts of agricultural production systems, agricultural input efficiency, and food choice. *Environ Res Lett* 2017; **12**: 064016.
- Clune S. Systematic review of greenhouse gas emissions for different fresh food categories. *J Clean Prod* 2017; **140**: 766–83.
- Clark MA, Springmann M, Hill J, Tilman D. Multiple health and environmental impacts of foods. *Proc Natl Acad Sci USA* 2019; **116**: 23357–62.
- Murphy SP, Allen LH. Nutritional importance of animal source foods. *J Nutr* 2003; **133**: 3932S–35S.
- Neumann C, Harris DM, Rogers LM. Contribution of animal source foods in improving diet quality and function in children in the developing world. *Nutr Res* 2002; **22**: 193–220.
- Melse-Boonstra A. Bioavailability of micronutrients from nutrient-dense whole foods: zooming in on dairy, vegetables, and fruits. *Front Nutr* 2020; **7**: 101.
- Black MM. Effects of vitamin B12 and folate deficiency on brain development in children. *Food Nutr Bull* 2008; **29** (suppl): S126–31.
- Stabler SP, Allen RH. Vitamin B12 deficiency as a worldwide problem. *Annu Rev Nutr* 2004; **24**: 299–326.
- Beal T, Ortenzi F. Priority micronutrient density in foods. *Research Square* 2021; published online Nov 3. <https://doi.org/10.21203/rs.3.rs-701840/v3> (preprint).
- Adesogan A. 400 Importance of animal-source foods for meeting global nutritional, educational and economic needs. *J Anim Sci* 2018; **96** (suppl 3): 164.
- Allen L. Comparing the value of protein sources for maternal and child nutrition. *Food Nutr Bull* 2013; **34**: 263–66.
- Zaharia S, Ghosh S, Shrestha R, et al. Sustained intake of animal-sourced foods is associated with less stunting in young children. *Nat Food* 2021; **2**: 246–54.
- Micha R, Shulkin ML, Peñalvo JL, et al. Etiologic effects and optimal intakes of foods and nutrients for risk of cardiovascular diseases and diabetes: systematic reviews and meta-analyses from the Nutrition and Chronic Diseases Expert Group (NutriCoDE). *PLoS One* 2017; **12**: e0175149.
- World Cancer Research Fund International. Diet, nutrition, physical activity and cancer: a global perspective third expert report. 2018. <https://www.wcrf.org/dietandcancer> (accessed June 30, 2021).
- Willett W, Rockström J, Loken B, et al. Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *Lancet* 2019; **393**: 447–92.
- Poore J, Nemecek T. Reducing food's environmental impacts through producers and consumers. *Science* 2018; **360**: 987–92.
- Springmann M, Godfray HCJ, Rayner M, Scarborough P. Analysis and valuation of the health and climate change cobenefits of dietary change. *Proc Natl Acad Sci USA* 2016; **113**: 4146–51.
- Kremen C, Miles A. Ecosystem services in biologically diversified versus conventional farming systems: benefits, externalities, and trade-offs. *Ecol Soc* 2012; **17**: art40.
- Mottet A, Teillard F, Boettcher P, De' Besi G, Besbes B. Review: domestic herbivores and food security: current contribution, trends and challenges for a sustainable development. *Animal* 2018; **12**: s188–98.
- Micha R, Khatibzadeh S, Shi P, Andrews KG, Engell RE, Mozaffarian D. Global, regional and national consumption of major food groups in 1990 and 2010: a systematic analysis including 266 country-specific nutrition surveys worldwide. *BMJ Open* 2015; **5**: e008705.
- Singh GM, Micha R, Khatibzadeh S, et al. Global, regional, and national consumption of sugar-sweetened beverages, fruit juices, and milk: a systematic assessment of beverage intake in 187 countries. *PLoS One* 2015; **10**: e0124845.
- Afshin A, Sur PJ, Fay KA, et al. Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2019; **393**: 1958–72.
- Del Gobbo LC, Khatibzadeh S, Imamura F, et al. Assessing global dietary habits: a comparison of national estimates from the FAO and the Global Dietary Database. *Am J Clin Nutr* 2015; **101**: 1038–46.
- Mohan D, Mente A, Dehghan M, et al. Associations of fish consumption with risk of cardiovascular disease and mortality among individuals with or without vascular disease from 58 countries. *JAMA Intern Med* 2021; **181**: 631–49.
- Iqbal R, Dehghan M, Mente A, et al. Associations of unprocessed and processed meat intake with mortality and cardiovascular disease in 21 countries [Prospective Urban Rural Epidemiology (PURE) Study]: a prospective cohort study. *Am J Clin Nutr* 2021; **114**: 1049–58.
- Dehghan M, Mente A, Rangarajan S, et al. Association of dairy intake with cardiovascular disease and mortality in 21 countries from five continents (PURE): a prospective cohort study. *Lancet* 2018; **392**: 2288–97.
- Tong TYN, Appleby PN, Key TJ, et al. The associations of major foods and fibre with risks of ischaemic and haemorrhagic stroke: a prospective study of 418 329 participants in the EPIC cohort across nine European countries. *Eur Heart J* 2020; **41**: 2632–40.
- Headey D, Hirvonen K, Hoddinott J. Animal sourced foods and child stunting. *Am J Agric Econ* 2018; **100**: 1302–19.
- Miller V, Singh GM, Onopa J, et al. Global Dietary Database 2017: data availability and gaps on 54 major foods, beverages and nutrients among 5.6 million children and adults from 1220 surveys worldwide. *BMJ Glob Health* 2021; **6**: e003585.

For the Global Dietary Database see <https://www.globaldietarydatabase.org/data-download>

For the Global Dietary Database aggregate data see <https://www.globaldietarydatabase.org/management/microdata-surveys>

- 31 Micha R, Kalantarian S, Wirojatana P, et al. Estimating the global and regional burden of suboptimal nutrition on chronic disease: methods and inputs to the analysis. *Eur J Clin Nutr* 2012; **66**: 119–29.
- 32 Khatibzadeh S, Saheb Kashaf M, Micha R, et al. A global database of food and nutrient consumption. *Bull World Health Organ* 2016; **94**: 931–34.
- 33 Micha R, Khatibzadeh S, Shi P, et al. Global, regional, and national consumption levels of dietary fats and oils in 1990 and 2010: a systematic analysis including 266 country-specific nutrition surveys. *BMJ* 2014; **348**: g2272.
- 34 Powles J, Fahimi S, Micha R, et al. Global, regional and national sodium intakes in 1990 and 2010: a systematic analysis of 24 h urinary sodium excretion and dietary surveys worldwide. *BMJ Open* 2013; **3**: e003733.
- 35 Food and Agriculture Organization of the United Nations. Food balances. 2021. <https://www.fao.org/faostat/en/#data/2021> (accessed March 3, 2021).
- 36 Smith MR, Micha R, Golden CD, Mozaffarian D, Myers SS. Global Expanded Nutrient Supply (GENUS) Model: a new method for estimating the global dietary supply of nutrients. *PLoS One* 2016; **11**: e0146976.
- 37 Gelman A, Pardoe I. Bayesian measures of explained variance and pooling in multilevel (hierarchical) models. *Technometrics* 2006; **48**: 241–51.
- 38 Wagner T, Diefenbach DR, Christensen S, Norton AS. Using multilevel models to quantify heterogeneity in resource selection. *J Wildl Manage* 2011; **75**: 1788–96.
- 39 United Nations Population Division. Total population by sex (thousands). 2019. <https://population.un.org/wpp/DataQuery/2019> (accessed June 12, 2020).
- 40 Barro R, Lee J. A new data set of educational attainment in the world, 1950–2010. *J Dev Econ* 2013; **104**: 184–98.
- 41 United Nations Population Division. Urban population (% of total population). 2018. <https://data.worldbank.org/indicator/SP.URB.TOTL.IN.ZS2018> (accessed June 12, 2020).
- 42 Climate Watch. Climate Watch data. 2019. <http://www.climatewatchdata.org> (accessed July 7, 2021).
- 43 Huang L, Wang Z, Wang H, et al. Nutrition transition and related health challenges over decades in China. *Eur J Clin Nutr* 2021; **75**: 247–52.
- 44 Pandey B, Reba M, Joshi PK, Seto KC. Urbanization and food consumption in India. *Sci Rep* 2020; **10**: 17241.
- 45 de Bruin S, Dengerink J, van Vliet J. Urbanisation as driver of food system transformation and opportunities for rural livelihoods. *Food Secur* 2021; **13**: 1–18.
- 46 Hibbeln JR, Spiller P, Brenna JT, et al. Relationships between seafood consumption during pregnancy and childhood and neurocognitive development: two systematic reviews. *Prostaglandins Leukot Essent Fatty Acids* 2019; **151**: 14–36.
- 47 Bahnfleth C, Kim J, Spahn J, et al. Seafood intake during pregnancy and lactation and child neurocognitive development: a NESR systematic review. *Curr Dev Nutr* 2021; **5** (suppl 2): 712.
- 48 Schlesinger S, Neuenschwander M, Schwedhelm C, et al. Food groups and risk of overweight, obesity, and weight gain: a systematic review and dose-response meta-analysis of prospective studies. *Adv Nutr* 2019; **10**: 205–18.
- 49 Bechthold A, Boeing H, Schwedhelm C, et al. Food groups and risk of coronary heart disease, stroke and heart failure: a systematic review and dose-response meta-analysis of prospective studies. *Crit Rev Food Sci Nutr* 2019; **59**: 1071–90.
- 50 Storhaug CL, Fosse SK, Fadnes LT. Country, regional, and global estimates for lactose malabsorption in adults: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol* 2017; **2**: 738–46.
- 51 Miller V, Micha R, Choi E, et al. Evaluation of the quality of evidence of the association of foods and nutrients with cardiovascular disease and diabetes: a systematic review. *JAMA Netw Open* 2022; **5**: e2146705.
- 52 Herber C, Bogler L, Subramanian, S.V. et al. Association between milk consumption and child growth for children aged 6–59 months. *Sci Rep* 2020; **10**: 6730.
- 53 Granic A, Dismore L, Hurst C, et al. Myoprotective whole foods, muscle health and sarcopenia: a systematic review of observational and intervention studies in older adults. *Nutrients* 2020; **12**: 2257.