

Determinants of the Number of Main Canals in a Tooth: Deciphering Potential Mechanisms *

Determinantes del número de conductos principales en un diente: descifrando posibles mecanismos

Determinantes do número de canais principais em um dente: decifrando mecanismos potenciais

Andrea Alejandra Moreno Robalino ^a
Universidad de Cuenca. Cuenca, Ecuador
alejandra.moreno27@ucuenca.edu.ec
<https://orcid.org/0000-0003-0113-6092>

DOI : <https://doi.org/10.11144/Javeriana.uo42.dnmc>
Submission Date: 12 December 2022
Acceptance Date: 31 July 2023
Publication Date: 18 December 2023

José Luis Álvarez Vásquez ^a
Universidad de Cuenca. Cuenca, Ecuador
jose.alvarezv@ucuenca.edu.ec
<https://orcid.org/0000-0003-0381-2402>

Authors' Note: ^a **Correspondence:** alejandra.moreno27@ucuenca.edu.ec; jose.alvarezv@ucuenca.edu.ec

ABSTRACT

Background: Although millions of root canal treatments are performed globally on a daily basis, factors that determine the number of main root canals in a tooth have not yet been elucidated. Variations in the number of root canals in different teeth is of utmost importance in clinical practice. However, clinicians aren't aware about the determinants of such number, let alone these determinants have been approached in the literature, to the best of our knowledge. **Purpose:** This narrative review aimed to integrate the potential mechanisms involved in determining the number of main canals in a permanent tooth. **Methods:** We used the search terms "root canal number," "root canal morphology," "tooth morphology," "root development," and "root formation" to identify articles from the PubMed and Scopus databases. **Results:** 57 articles and 2 books were obtained. A multifactorial basis is plausible considering the influence of anthropological, demographic, environmental, genetic, epigenetic, tooth size related mechanisms and the pivotal role of Hertwig's epithelial root sheath. Live-cell imaging techniques, mathematical models, quantitative genetics and dental phenomics could provide insightful information in the near future. **Conclusions:** Overall, it seems that the potential mechanisms determining the number of main canals in a tooth have a multifactorial basis. The orchestrating role of the Hertwig's epithelial root sheath seems pivotal, although the specific regulatory signals that induce or repress its diaphragmatic processes remain unknown. However, there is a dire need for molecular studies that help unveil these and other potential mechanisms involved. **Keywords:** dental anatomy; dentistry; endodontics; Hertwig's epithelial root sheath; root canal morphology; root canal number; root canal system; tooth root development; tooth root formation

RESUMEN

Antecedentes: aunque diariamente se realizan millones de tratamientos de conducto radicular en todo el mundo, aún no se han dilucidado los factores que determinan el número de conductos radiculares principales en un diente. Las variaciones en el número de conductos radiculares en diferentes dientes son de suma importancia en la práctica clínica. Sin embargo, los médicos no conocen los determinantes de tal número, y mucho menos estos determinantes han sido abordados en la literatura, según nuestro conocimiento. **Objetivo:** Esta revisión narrativa tuvo como objetivo integrar los posibles mecanismos involucrados en la determinación del número de conductos principales en un diente permanente. **Métodos:** Utilizamos los términos de búsqueda "número de conducto radicular", "morfología del conducto radicular", "morfología dental", "desarrollo

radicular" y "formación radicular" para identificar artículos de las bases de datos PubMed y Scopus. **Resultados:** se obtuvieron 57 artículos y 2 libros. Una base multifactorial es plausible considerando la influencia de los mecanismos antropológicos, demográficos, ambientales, genéticos, epigenéticos, relacionados con el tamaño del diente y el papel fundamental de la vaina radicular epitelial de Hertwig. Las técnicas de imagen de células vivas, los modelos matemáticos, la genética cuantitativa y la fenómica dental podrían proporcionar información reveladora en un futuro próximo. **Conclusiones:** En general, parece que los posibles mecanismos que determinan el número de conductos principales en un diente tienen una base multifactorial. El papel orquestador de la vaina radicular epitelial de Hertwig parece fundamental, aunque las señales reguladoras específicas que inducen o reprimen sus procesos diafragmáticos siguen sin conocerse. Sin embargo, existe una gran necesidad de estudios moleculares que ayuden a revelar estos y otros posibles mecanismos involucrados.

Palabras clave: anatomía dental; desarrollo de la raíz del diente; endodoncia; formación de la raíz del diente; morfología del conducto radicular; número de conductos radiculares; odontología; sistema de conductos radiculares; vaina radicular epitelial de Hertwig

RESUMO

Antecedentes: Embora milhões de tratamentos de canal radicular sejam realizados diariamente em todo o mundo, os fatores que determinam o número de canais radiculares principais em um dente ainda não foram elucidados. Variações no número de canais radiculares em diferentes dentes são de extrema importância na prática clínica. No entanto, os clínicos não têm conhecimento dos determinantes desse número e muito menos esses determinantes foram abordados na literatura, até onde sabemos. **Objetivo:** Esta revisão narrativa teve como objetivo integrar os potenciais mecanismos envolvidos na determinação do número de canais principais em um dente permanente. **Métodos:** Utilizamos os termos de pesquisa “número do canal radicular”, “morfologia do canal radicular”, “morfologia dentária”, “desenvolvimento radicular” e “formação radicular” para identificar artigos das bases de dados PubMed e Scopus. **Resultados:** foram obtidos 57 artigos e 2 livros. Uma base multifatorial é plausível, considerando a influência de mecanismos antropológicos, demográficos, ambientais, genéticos, epigenéticos, relacionados ao tamanho dos dentes e o papel central da bainha epitelial da raiz de Hertwig. Técnicas de imagem de células vivas, modelos matemáticos, genética quantitativa e fenômica dentária poderiam fornecer informações esclarecedoras em um futuro. **Conclusões:** No geral, parece que os potenciais mecanismos que determinam o número de canais principais num dente têm uma base multifatorial. O papel orquestrador da bainha epitelial da raiz de Hertwig parece fundamental, embora os sinais regulatórios específicos que induzem ou reprimem seus processos diafragmáticos permaneçam desconhecidos. No entanto, há uma extrema necessidade de estudos moleculares que ajudem a desvendar estes e outros potenciais mecanismos envolvidos.

Palavras-chave: anatomia dentária; bainha radicular epitelial de Hertwig; desenvolvimento da raiz dentária; endodontia; formação de raiz dentária; morfologia do canal radicular; número de canais radiculares; odontologia; sistema de canais radiculares

INTRODUCTION

A deep knowledge of the root anatomy and configuration of the root canal system (RCS) of each dental group is essential to conduct an adequate root canal treatment (1,2). Existing canal configuration classifications according to Vertucci (3), Gulabivala (4), and Sert & Bayirli (5) are based on the pulp space configurations and number of main canals in the tooth. The clinician must have sufficient skills and knowledge to treat teeth that present a complex RCS (6-10), which sometimes involves precisely analyzing the number of main canals. However, although Carabelli provided the first detailed description of the number of canals in teeth by 1842 (11) and millions of root canal treatments are performed globally on a daily basis (12), the factors that determine the number of main root canals in a tooth have not yet been elucidated (Thesleff, I. personal communication).

The Hertwig's epithelial root sheath (HERS) plays a critical role in root formation (13,14), determining the shape, size and number of dental roots (15). Genetic studies have identified that several molecules, such as Nfic, Osterix, β -catenin, and sonic hedgehog, constitute key factors in root formation (13). Since these molecules regulate root formation, they could help to understand the mechanisms determining the number of canals. However, the precise factors determining the number of canals in a tooth are still unknown, since there are case reports of single-rooted teeth with four canals (9), molars with a single root and single canal (16), and maxillary (17) and mandibular molars (18) with eight canals.

However, it has been postulated that root morphology results from the complex interaction of genetic, epigenetic, and environmental factors (19-23). Demographic and ethnic factors are interrelated with root anatomy, root number, and RCS configuration (1,2,7,24,25); therefore, variations can be considered as specific traits that occur with varying prevalence in certain populations (7,24,25).

In the last decade, a series of investigations have been conducted in various populations using cone-beam computed tomography (CBCT) (1,2,7,24-27) or microcomputed tomography (μ CT) (8,10,28,29) to determine variations in both root anatomy and RCS configurations. However, to the best of our knowledge, no publications have adequately explained the mechanisms or factors that determine the number of main root canals in a permanent tooth. This narrative review aimed to integrate the potential mechanisms involved in determining the number of main canals in a permanent tooth. In fact, skilled clinicians should be aware about the fundamentals behind the determinants of the number of main canals in different teeth, as they deal with such variations on a daily basis. Additionally, future perspectives are also included in the final section of the manuscript.

MATERIALS AND METHODS

Available literature was searched in the PubMed and Scopus databases to identify relevant articles, from the first reports until May 25, 2022, using the search terms “root canal number,” “root canal morphology,” “tooth morphology,” “root development,” and “root formation”. In this study, we included only articles published in English. The reference lists of the selected articles were manually searched to complement the electronic search.

The search strategy was specific for each database. Experimental and non-experimental studies were included, in order to address various aspects such as concepts, definitions, methodologies and available scientific evidence. In most cases, articles that did not explicitly link to the search terms used, were removed. Then, the search results were merged, proceeding to remove duplicates. After the initial selection of titles and abstracts, only articles and books of greatest relevance to the topic under review were selected. Finally, 57 articles and 2 books were obtained. The information was categorized according to the sections of the present narrative review.

RESULTS

We found the following mechanisms to be potentially responsible for main canal determination in a tooth: anthropological and demographic mechanisms, environmental mechanisms, genetic mechanisms, epigenetic mechanisms, HERS mechanisms, and tooth size-related mechanisms.

Anthropological and Demographic Mechanisms

Root canal information of many hominin taxa is rarely available because of the limitations in historical methodologies (22). Nevertheless, valuable information concerning canal variations in hominin fossils have been documented using μ CT (22,30-32), thus elucidating potential developmental influences on root morphology in modern humans and their fossil relatives (22).

Among the extant and extinct hominoid species, the premolar root morphology is a useful phylogenetic indicator since its root number, root size, and canal number vary the most compared to other tooth types (31-33). In South African hominin fossils attributed to *Australopithecus africanus* (2-3 million years ago) and *Paranthropus robustus* (1-2 million years ago), the canal numbers in maxillary premolars (1-3 roots) varied between one (rarely) and two or three (more commonly), while in mandibular premolars (1-3 roots) it varied between one and three, and even four canals in the case of double plate-like mesial and distal roots (22). Both hominins exhibited reduced ma

xillary premolar root/canal numbers from the inferred ancestral great ape condition; however, *A. africanus* showed more reduction than *P. robustus* (22). Figure 1 shows a CBCT axial view of a mandibular first premolar with four double-plate-like canals in a patient, similar to the aforementioned type. It is possible that this case reflects an atavistic trait.



FIGURE 1

CBCT axial view of the left mandibular first premolar (red arrow) of a female patient that had four canals (author's personal database- JLAV), reflecting the double plate-like type mandibular premolars described in South African hominin fossils.

Source: the authors.

The genus *homo*, particularly the lineage leading to modern humans, has been characterized by a reduction in root/canal number, while the genus *Paranthropus* has displayed an elaboration in the form and number of mandibular premolars (22). However, a study on root and root canal diversity in human premolars and molars from two historical periods found an increase in the diversity in the number of roots and shape of root canals in the modern period compared to the late medieval period (34).

In contrast, *Homo floresiensis*, an extinct diminutive hominin species (found on an Indonesian island) with unique physical characteristics, has shown primitive canine-premolar (comparable to *H. Erectus*) and advanced molar morphologies (even compared to those of modern humans), a combination of dental traits unknown to any other hominin species (35). However, to the best of our knowledge, no studies have addressed root canal variations in this species, and such investigations could help us understand the influence of anthropological factors on root/canal morphology, especially since teeth are one of the most informative elements in hominin evolutionary studies (36).

Finally, anthropological studies allow us to link the variations and discrepancies found in the different populations due to prehistoric colonization and their division into the three main branches of the phenotypes. In this sense, each population was exposed to different environmental factors, genetic and demographic changes, and natural pressures, thus generating the phenotypes of each population, which determined variations both in the maxillary bones and in the teeth (22-25). This clearly shows that demographic and ethnic factors are interrelated with root anatomy, root number, and RCS configuration. For instance, the maxillary (17) and mandibular molars (18) with eight canals, which are currently the teeth with the greatest reported number of main canals, are from Indian patients.

Environmental Mechanisms

Although the influence of environmental factors on dental morphology is not yet clearly understood (37), environmental disruption caused by non-genetic factors, such as malnutrition and systemic diseases during tooth development, has been found to influence the final phenotype (37). Malnutrition and systemic diseases, such as cystic fibrosis, human immunodeficiency virus infection, and leukemia, negatively affect both the dentine and enamel during tooth development. These environmental factors, as stress markers, could have a significant effect on the morphological variations of teeth, e.g., increasing variability in the dental cusps (37,38). Additionally, diet is an environmental factor that alters genetic expression and changes in DNA methylation patterns and can affect the final phenotype of the offspring (37-40).

Studies based on primates have demonstrated an increase in the surface area of the dental root in those consuming hard food, such as seeds, thus evidencing a link between dental root morphology and diet. However, masticatory loads significantly influence the dental root development (23). Species of primates that eat hard or fibrous food demonstrate an increase in root curvature compared to species that eat fruits and barks. This adaptive process can be transmitted by species and thus influence dental root morphology (23), in turn potentially affecting the number and/or disposition of canals.

During pregnancy, environmental factors, such as alcohol consumption, smoking habits, and obesity, have demonstrated negative effects on and caused asymmetry in children's teeth (37), but there are no studies concerning root development or the number of canals. On the other hand, it is incredible how monozygotic twins show differences in shape, number, and teeth size, while in appearance they are almost identical (37,41,42). This supports the potential role of environmental factors in tooth morphology; indeed, certain transplacental teratogens and diseases can cause differences in monozygotic twin phenotypes (39). In addition, scientific research has underlined that tooth development is not only controlled by genetic factors, but also by environmental factors (37,38,43) and they equally affect the developmental changes in teeth morphology (37).

The increased variability in the expression of morphological traits produced by an environmental stressor can be explained by a buffering mechanism (37). The genetic buffering mechanism controls different phenotypes that can be expressed in each individual (44,45). Therefore, it reduces the probability of mutation of a specific gene (44,45). During normal development, the presence of hidden genetic variability can be canalized by the buffering mechanism; however, if a stressful factor exceeds the threshold, the buffering mechanism cannot canalize, and it appears as new genetic variation (37,44). Nonetheless, the mechanisms or factors that produce this new genetic variability are not clear (37).

Finally, tooth development corresponds to a morpho dynamic mechanism in which morphogenesis and induction processes play significant interdependent roles (37,46). Therefore, if any error occurs during the developmental cascade and affects the morpho dynamic process, it can generate changes in the final morphology (37). Environmental stressors can affect both genetic and cellular interactions, which in turn control tooth development. These stressors can downregulate cellular metabolism and changes in epithelial and mesenchymal growth and differentiation rates (37); however, their effect on the number of canals in teeth have not been explored yet.

Genetic Mechanisms

Genetic expression plays a pivotal role in variations in dental morphology, with more than 300 genes playing roles in different stages of the dental morphogenetic process (39,43,46-48). In this regard, an increase in the expression of only one gene in the DNA sequence can lead to significant changes in dental

development, such as increased tooth number, altered tooth shape, disturbed tooth size, increased cusp numbers, and new variations (41,47).

Gene mutations can appear at different stages of odontogenesis, generating changes in normal tooth development and tooth morphology that can be translated to offsprings (40,47). When an error occurs during the transcription process of evolutionary genes, it leads to morphological diversity, especially when the homeobox genes are involved (40,49); Cis regulatory elements (CREs) of genes are specific non-coding DNA sites that regulate the transcription process of neighboring genes (40), and play paramount roles in controlling gene expression in specific cell types, conditions, and developmental stages (50). Disruption of these regions are considered as the most prevalent cause of phenotypic changes, especially morphological (50,51), so it could be that CREs have a role in determining the number of main canals.

Interestingly, the effect of gene mutation on the dentition depends on the gene type and the region of the affected gene (40). There are many genes that are strongly related to dental development, such as *Msx1*, *Pax9*, and *Lef1*, and genes that have less influence on developmental processes, such as *Dlx1*, *Dlx2*, *Gli2*, and *Gli3* (40). In addition, the downstream targets of tooth-specific transcription factors and their role in regulatory mechanisms are poorly characterized (49,52); therefore, there is still a need to elucidate the role of genetics in odontogenesis, and specifically its involvement in main canal number determination.

Epigenetic Mechanisms

Epigenetics involves processes that occur in the cell nuclei and generate modifications in gene expression while still maintaining the same DNA sequence (42,43,53). Epigenetic influence can be explained as an autonomous developmental system with self-organization (39,43); therefore, even in monozygotic twins that share the same genotype, great differences can occur in their phenotype (39). External influences on epigenetic processes can occur, such as the effects of diet in cancer, and thus epigenetic mechanisms would allow an organism to respond to the environment through changes in gene expression (53).

It is important to highlight that epigenetic processes are crucial for development and differentiation, even in mature humans. Much epigenetic research has focused on two molecular mechanisms that mediate this phenomenon: DNA methylation and histone modifications. DNA methylation signals can repress transcription through the exclusion of proteins involved in histone modifications, through their DNA binding sites, and cause long-term silencing of gene expression (53). Odontogenesis requires the participation of many transcription factors (48,54,55), and repression through DNA methylation may be involved in the regulation of the number of main canals.

It has been postulated that epigenetic factors may explain dental differences in monozygotic twin pairs, such as variations in the expression of missing or extra teeth (39) and can also influence the number or position of the teeth (39,42). Moreover, differential placental implantation and nutrition, differential transplacental teratogens, and infections have been cited as possible environmental factors (56).

Finally, it has been regarded that dental phenotype variations within species may respond to epigenetic mechanisms, like minor variations in the timing of interactions between cells and their position, acting at a local tissue level (39). In this regard, studies on epigenetic influences in monozygotic twin pairs could help elucidate the mechanisms involved in root canal number determination.

HERS Mechanisms

HERS is known to play a critical role in root formation (13,14) and determining the shape, size, and number of dental roots (15). The number of roots is determined by the number of inward-bending horizontal processes, called interradicular or diaphragmatic processes (DP), that are emitted from the epithelial diaphragm shortly after crown formation; if two DP are extended, two roots will be formed, with three DP, three roots will be formed, and so on; if no DP is formed, then a single-rooted tooth will be the result. The root structure (single- or multi-rooted) is thus determined by the shape and folding of the HERS (19,21,32,57).

As for the number of canals, each formed root has a minimum of one, so inherently multi-rooted teeth have more canals. Nevertheless, single-rooted teeth with multiple canals, which are common in non-human extant ape genera such as *Hylobates*/Pan maxillary first premolars and Gorilla/Pongo mandibular premolars, demonstrate a tendency for DP suppression of the outer surface but not in the expression of individual canals (31). Similarly, the above mentioned South African hominin fossils attributed to *A. africanus* and *P. robustus* exhibit multiple canals within one root body in both maxillary and mandibular premolars (22). Nonetheless, the cause of this “selective” suppression has not yet been elucidated, and the specific molecular signals that govern root size and induce the invagination of HERS remain unknown in hominids and hominins (30). It is known that the timing and rate of constriction of the HERS regulate root length, i.e., if the HERS narrows rapidly, the root will be shorter, and if not, the root will be longer (19).

It is worth mentioning taurodont molars, which show enlarged pulp chambers and apically positioned root furcations, resulting in an enlarged root body with short root branches and their corresponding terminal canals (30). Taurodontism is common in modern human deciduous and permanent dentitions, both bilaterally and in more than one tooth type (58), and its extreme form is hypertaurodont or pyramidal molars, which have been described in modern humans (59). However, there are no known cases in any fossil member of the genus *Homo*, other than Neanderthals (at a high frequency) (30). Pyramidal molars have a bifurcation either positioned at the apical fourth of the root or almost completely missing and seem to be the result of a complete failure of DP formation (30).

Additionally, crown pattern formation in a tooth may depend on the spatial and temporal expression of activating and inhibiting molecules derived from enamel knots (60). In an analogous comparison, it could be hypothesized that there are regulation centers at the DP level that determine the number of canals and generate a spatiotemporal expression of activating and inhibitory molecules (Figure 2); however, future investigations are needed to study this hypothesis.



FIGURE 2

Apical view of a histology-based 3D reconstruction of a mouse upper first molar on day 6. Three HERS processes, that is, diaphragmatic processes (DP) are in contact at two regions (arrows), giving rise to three root-canal spaces (asterisks). It could be hypothesized that there are regulation centers at the DP level, determining the number of main root canals, by generating a spatiotemporal expression of activating and inhibitory molecules. M, mesial; D, distal; B, buccal; P, palatal. Adapted from Shimazu, *et al.*, 2009 (57).

Tooth Size-Related Mechanisms

A developmental model called the root “size/number continuum” (SNC) has been proposed for premolars, which correlates increasing root number, and in turn root canals, with tooth size. This model predicts that tooth germ size, inferred from tooth size, predisposes the number and expression of DP, i.e., smaller premolars are predisposed to fewer roots and larger premolars to greater numbers of roots (21). Tooth germ size is linked to the final phenotype (61). It affects the number of molecular signals and the distance between signaling centers, such as the number of enamel knots and/or DP (21). However, studies using the SNC model and the cross-sectional cervix area as a proxy for tooth size in hominoids (31,32) and hominins (22) have provided equivocal support; thus, further investigation is needed.

In addition, size and shape of teeth are determined by cellular and apoptotic interactions during dental development (38,46). Ectodysplasin A (EDA) is an important signaling pathway involved in the regulation of tooth signaling centers. Inhibition of the EDA pathway generates a reduction in signaling centers, creating small teeth buds, and an increase in EDA signaling results in larger teeth buds and extra teeth (46). In this regard, the EDA pathway may be involved in the regulation of signaling centers in the DP, thus having a potential role in root canal number determination. Additionally, hyaluronic acid has

been shown to affect cellular decisions regarding tooth size and number by controlling proliferation, cell orientation, and migration in the developing tooth (62), and thus could have a role as well.

Many studies have demonstrated the relationship between tooth size, tooth number, and individual sex; therefore, studies in women have shown a reduction in tooth size and an increase in missing teeth, while men have shown an increase in tooth size and a major number of extra teeth (39,42,46). However, a study of four extant apes taxa showed that male and female individuals within genera differ in tooth size but not in canal/root form and number (31).

Finally, an investigation analyzed the variability and patterning of permanent tooth size in four human ethnic groups – Southern Chinese, North Americans of European ancestry, Modern British of European ancestry, and Romano-British – evidencing that the different patterns of tooth size observed between the study groups reflect different genetic and environmental influences to dental development (63). This adds to the information on the potential mechanisms involved in root canal number determination.

CONCLUSIONS AND FUTURE PERSPECTIVES

Overall, it seems plausible that the potential mechanisms determining the number of main canals in a tooth have a multifactorial basis, considering the influence of anthropological, demographic, environmental, genetic, epigenetic and tooth size related mechanisms. The orchestrating role of the HERS in root development seems pivotal, although the specific molecular signals that govern the induction or repression of DP remain unknown until now. Furthermore, the latest findings on odontogenesis could help to decipher these potential mechanisms, such as the role of the transcription factor *Nfic* in root patterning and growth through the regulation of mesenchymal cell proliferation, which may interact with HERS to guide the size, shape, and number of tooth roots (64).

In the near future, live-cell imaging techniques could be useful tools to elucidate the mechanisms involved in odontogenesis, such as the number of main canals, because this innovative technique allows direct observation of cell movement and morphological changes (65,66). Further, mathematical models have been developed to demonstrate how large morphological changes can be produced by small epigenetic events (67), thus these models can help provide evidence for factors that determine the number of main canals in a tooth. Quantitative genetics may also provide insight into these mechanisms because it has proven useful in outlining the contribution of genes to tooth form, how genetic effects determine dental phenotypes, and the potential for pleiotropy (68). Additionally, dental phenomics provides acquisition of high-dimensional phenotypic data on a large scale and has been used to quantify phenotypic variations in craniofacial development (69). Hence, it can help to elucidate the roles of genetic, epigenetic, and environmental factors in the variations in root canal number because such interactions account for variations in tooth number, size, and shape within and among species (20).

Finally, it must be emphasized that variations in the number of root canals in different teeth are of utmost importance in clinical practice. Skilled clinicians should be aware of the fundamentals behind the determinants of the number of main canals in different teeth, as they deal with such variations on a daily basis. In this regard, endodontists, biologists, and embryologists should intensify their understanding of the complex development of the RCS in order to support accurate endodontic therapy (70) and unveil potential mechanisms involved in determining the number of main canals in a tooth.

RECOMMENDATIONS

Due to the scarce information found, in the near future it would be important to conduct more research concerning studies related to the biological and molecular aspects of the potential mechanisms that

determine the number of main canals in a tooth. In addition, in future studies we recommend including studies in other languages, since the present study only included literature in English.

ACKNOWLEDGEMENTS

This article is based on Andrea Moreno-Robalino's undergraduate thesis.

References

1. Martins JNR, Marques D, Silva EJNL, Caramês J, Versiani MA. Prevalence Studies on Root Canal Anatomy Using Cone-beam Computed Tomographic Imaging: A Systematic Review. *J Endod.* 2019 Apr; 45(4): 372-386.e4. <https://dx.doi.org/10.1016/j.joen.2018.12.016>
2. Li YH, Bao SJ, Yang XW, Tian XM, Wei B, Zheng YL. Symmetry of root anatomy and root canal morphology in maxillary premolars analyzed using cone-beam computed tomography. *Arch Oral Biol.* 2018 Oct; 94: 84-92. <https://dx.doi.org/10.1016/j.archoralbio.2018.06.020>
3. Vertucci FJ. Root canal anatomy of the human permanent teeth. *Oral Surg Oral Med Oral Pathol.* 1984 Nov; 58(5): 589-599. [https://dx.doi.org/10.1016/0030-4220\(84\)90085-9](https://dx.doi.org/10.1016/0030-4220(84)90085-9)
4. Gulabivala K, Aung TH, Alavi A, Ng YL. Root and canal morphology of Burmese mandibular molars. *Int Endod J.* 2001 Jul; 34(5): 359-370. <https://dx.doi.org/10.1046/j.1365-2591.2001.00399.x>
5. Sert S, Bayirli GS. Evaluation of the root canal configurations of the mandibular and maxillary permanent teeth by gender in the Turkish population. *J Endod.* 2004 Jun;30(6): 391-398. <https://dx.doi.org/10.1097/00004770-200406000-00004>
6. Zhang M, Xie J, Wang YH, Feng Y. Mandibular first premolar with five root canals: a case report. *BMC Oral Health.* 2020 Sep 10; 20(1): 253. <https://dx.doi.org/10.1186/s12903-020-01241-0>
7. Gomez F, Brea G, Gomez-Sosa JF. Root canal morphology and variations in mandibular second molars: an in vivo cone-beam computed tomography analysis. *BMC Oral Health.* 2021 Sep 1; 21(1): 424. <https://dx.doi.org/10.1186/s12903-021-01787-7>
8. Wolf TG, Stiebritz M, Boemke N, Elsayed I, Paqué F, Wierichs RJ, Briseño-Marroquín B. 3-dimensional Analysis and Literature Review of the Root Canal Morphology and Physiological Foramen Geometry of 125 Mandibular Incisors by Means of Micro-Computed Tomography in a German Population. *J Endod.* 2020 Feb; 46(2): 184-191. <https://dx.doi.org/10.1016/j.joen.2019.11.006>
9. Aznar Portoles C, Moinzadeh AT, Shemesh H. A Central Incisor with 4 Independent Root Canals: A Case Report. *J Endod.* 2015 Nov; 41(11): 1903-1906. <https://dx.doi.org/10.1016/j.joen.2015.08.001>
10. Zhang W, Tang Y, Liu C, Shen Y, Feng X, Gu Y. Root and root canal variations of the human maxillary and mandibular third molars in a Chinese population: A micro-computed tomographic study. *Arch Oral Biol.* 2018 Nov; 95: 134-140. <https://dx.doi.org/10.1016/j.archoralbio.2018.07.020>
11. Perrini N, Versiani MA. Historical Overview of the Studies on Root Canal Anatomy. In: Versiani MA, Basrani B, Sousa-Neto M, editors. *The Root Canal Anatomy in Permanent Dentition.* Cham: Springer; 2019. p.10. https://dx.doi.org/10.1007/978-3-319-73444-6_1
12. American Association of Endodontists. Healthier Mouth=Healthier You. Tooth Wisdom Infographic. <https://newsroom.aae.org/healthier-you/>. Accessed April 29, 2022.
13. Wang J, Feng JQ. Signaling Pathways Critical for Tooth Root Formation. *J Dent Res.* 2017 Oct; 96(11): 1221-1228. <https://dx.doi.org/10.1177/0022034517717478>
14. Zeichner-David M, Oishi K, Su Z, Zakartchenko V, Chen LS, Arzate H, Bringas P Jr. Role of Hertwig's epithelial root sheath cells in tooth root development. *Dev Dyn.* 2003 Dec; 228(4): 651-663. <https://dx.doi.org/10.1002/dvdy.10404>
15. Huang X, Xu X, Bringas P Jr, Hung YP, Chai Y. Smad4-Shh-Nfic signaling cascade-mediated epithelial-mesenchymal interaction is crucial in regulating tooth root development. *J Bone Miner Res.* 2010 May; 25(5): 1167-178. <https://dx.doi.org/10.1359/jbmr.091103>
16. Ioannidis K, Lambrianidis T, Beltes P, Besi E, Malliari M. Endodontic management and cone-beam computed tomography evaluation of seven maxillary and mandibular molars with single roots and single canals in a patient. *J Endod.* 2011 Jan; 37(1): 103-109. <https://dx.doi.org/10.1016/j.joen.2010.09.001>
17. Kottoor J, Velmurugan N, Surendran S. Endodontic management of a maxillary first molar with eight root canal systems evaluated using cone-beam computed tomography scanning: a case report. *J Endod.* 2011 May; 37(5): 715-719. <https://dx.doi.org/10.1016/j.joen.2011.01.008>
18. Arora A, Acharya SR, Sharma P. Endodontic treatment of a mandibular first molar with 8 canals: a case report. *Restor Dent Endod.* 2015 Feb; 40(1): 75-78. <https://dx.doi.org/10.5395/rde.2015.40.1.75>

19. Kovacs I. Contribution to the ontogenetic morphology of roots of human teeth. *J Dent Res.* 1967 Sep-Oct; 46(5): 865-874. <https://dx.doi.org/10.1177/00220345670460054201>
20. Brook AH. Multilevel complex interactions between genetic, epigenetic and environmental factors in the aetiology of anomalies of dental development. *Arch Oral Biol.* 2009 Dec; 54 Suppl 1(Suppl 1): S3-17. <https://dx.doi.org/10.1016/j.archoralbio.2009.09.005>
21. Shields ED. Mandibular premolar and second molar root morphological variation in modern humans: What root number can tell us about tooth morphogenesis. *Am J Phys Anthropol.* 2005 Oct; 128(2): 299-311. <https://dx.doi.org/10.1002/ajpa.20110>
22. Moore NC, Thackeray JF, Hublin JJ, Skinner MM. Premolar root and canal variation in South African Plio-Pleistocene specimens attributed to *Australopithecus africanus* and *Paranthropus robustus*. *J Hum Evol.* 2016 Apr; 93: 46-62. <https://dx.doi.org/10.1016/j.jhevol.2015.12.002>
23. Hamon N, Emonet EG, Chaimanee Y, Guy F, Tafforeau P, Jaeger JJ. Analysis of dental root apical morphology: a new method for dietary reconstructions in primates. *Anat Rec (Hoboken).* 2012 Jun; 295(6): 1017-1026. <https://dx.doi.org/10.1002/ar.22482>
24. Martins JNR, Marques D, Leal Silva EJN, Caramês J, Mata A, Versiani MA. Influence of Demographic Factors on the Prevalence of a Second Root Canal in Mandibular Anterior Teeth - A Systematic Review and Meta-Analysis of Cross-Sectional Studies Using Cone Beam Computed Tomography. *Arch Oral Biol.* 2020 Aug; 116: 104749. <https://dx.doi.org/10.1016/j.archoralbio.2020.104749>
25. Martins JNR, Marques D, Silva EJNL, Caramês J, Mata A, Versiani MA. Second mesiobuccal root canal in maxillary molars-A systematic review and meta-analysis of prevalence studies using cone beam computed tomography. *Arch Oral Biol.* 2020 May; 113: 104589. <https://dx.doi.org/10.1016/j.archoralbio.2019.104589>
26. Usha G, Muddappa SC, Venkitachalam R, Singh VPP, Rajan RR, Ravi AB. Variations in root canal morphology of permanent incisors and canines among Asian population: A systematic review and meta-analysis. *J Oral Biosci.* 2021 Dec; 63(4): 337-350. <https://dx.doi.org/10.1016/j.job.2021.09.004>
27. Candeiro GTM, Monteiro Dodt Teixeira IM, Olimpio Barbosa DA, Vivacqua-Gomes N, Alves FRF. Vertucci's Root Canal Configuration of 14,413 Mandibular Anterior Teeth in a Brazilian Population: A Prevalence Study Using Cone-beam Computed Tomography. *J Endod.* 2021 Mar; 47(3): 404-408. <https://dx.doi.org/10.1016/j.joen.2020.12.001>
28. Wolf TG, Kozaczek C, Siegrist M, Betthäuser M, Paqué F, Briseño-Marroquín B. An Ex Vivo Study of Root Canal System Configuration and Morphology of 115 Maxillary First Premolars. *J Endod.* 2020 Jun; 46(6): 794-800. <https://dx.doi.org/10.1016/j.joen.2020.03.001>
29. Briseño-Marroquín B, Paqué F, Maier K, Willershausen B, Wolf TG. Root Canal Morphology and Configuration of 179 Maxillary First Molars by Means of Micro-computed Tomography: An Ex Vivo Study. *J Endod.* 2015 Dec; 41(12): 2008-2013. <https://dx.doi.org/10.1016/j.joen.2015.09.007>
30. Kupczik K, Hublin JJ. Mandibular molar root morphology in Neanderthals and Late Pleistocene and recent *Homo sapiens*. *J Hum Evol.* 2010 Nov; 59(5): 525-541. <https://dx.doi.org/10.1016/j.jhevol.2010.05.009>
31. Moore NC, Hublin JJ, Skinner MM. Premolar root and canal variation in extant non-human hominoidea. *Am J Phys Anthropol.* 2015 Oct; 158(2): 209-226. <https://dx.doi.org/10.1002/ajpa.22776>
32. Moore NC, Skinner MM, Hublin JJ. Premolar root morphology and metric variation in *Pan troglodytes verus*. *Am J Phys Anthropol.* 2013 Apr; 150(4): 632-646. <https://dx.doi.org/10.1002/ajpa.22239>
33. Wood BA, Abbott SA, Uytterschaut H. Analysis of the dental morphology of Plio-Pleistocene hominids. IV. Mandibular postcanine root morphology. *J Anat.* 1988 Feb; 156: 107-139
34. Przesmycka A, Jędrychowska-Dańska K, Masłowska A, Witas H, Regulski P, Tomczyk J. Root and root canal diversity in human permanent maxillary first premolars and upper/lower first molars from a 14th-17th and 18th-19th century Radom population. *Arch Oral Biol.* 2020 Feb; 110: 104603. <https://dx.doi.org/10.1016/j.archoralbio.2019.104603>
35. Kaifu Y, Kono RT, Sutikna T, Saptomo EW, Jatmiko, Due Awe R. Unique Dental Morphology of *Homo floresiensis* and Its Evolutionary Implications. *PLoS One.* 2015 Nov 18; 10(11): e0141614. <https://dx.doi.org/10.1371/journal.pone.0141614>
36. Bailey SE, Hublin JJ, editors. *Dental Perspectives on Human Evolution: State of the Art Research in Dental Paleoanthropology.* Dordrecht: Springer; 2007.
37. Riga A, Belcastro MG, Moggi-Cecchi J. Environmental stress increases variability in the expression of dental cusps. *Am J Phys Anthropol.* 2014 Mar; 153(3): 397-407. <https://dx.doi.org/10.1002/ajpa.22438>
38. Brook AH, Koh KSB, Toh VKL. Influences in a biologically complex adaptive system: Environmental stress affects dental development in a group of Romano-Britons. *J Des Nat Ecodynamics.* 2016; 11(1): 33-40.
39. Townsend GC, Richards L, Hughes T, Pinkerton S, Schwerdt W. Epigenetic influences may explain dental differences in monozygotic twin pairs. *Aust Dent J.* 2005 Jun; 50(2): 95-100. <https://dx.doi.org/10.1111/j.1834-7819.2005.tb00347.x>
40. Line SR. Variation of tooth number in mammalian dentition: connecting genetics, development, and evolution. *Evol Dev.* 2003 May-Jun; 5(3): 295-304. <https://dx.doi.org/10.1046/j.1525-142x.2003.03036.x>

41. Townsend G, Hughes T, Luciano M, Bockmann M, Brook A. Genetic and environmental influences on human dental variation: a critical evaluation of studies involving twins. *Arch Oral Biol.* 2009 Dec;54 Suppl 1(Suppl 1): S45-51. <https://dx.doi.org/10.1016/j.archoralbio.2008.06.009>
42. Townsend G, Bockmann M, Hughes T, Brook A. Genetic, environmental and epigenetic influences on variation in human tooth number, size and shape. *Odontology.* 2012 Jan;100(1):1-9. <https://dx.doi.org/10.1007/s10266-011-0052-z>
43. Townsend G, Brook A. Genetic, Epigenetic and Environmental Influences on Human Tooth Size, Shape and Number. *eLS* 2013. <https://doi.org/10.1002/9780470015902.a0024858>
44. Rutherford SL. From genotype to phenotype: buffering mechanisms and the storage of genetic information. *Bioessays.* 2000 Dec; 22(12): 1095-1105. [https://doi.org/10.1002/1521-1878\(200012\)22:12<1095::AID-BIES7>3.0.CO;2-A](https://doi.org/10.1002/1521-1878(200012)22:12<1095::AID-BIES7>3.0.CO;2-A)
45. Waddington CH. Canalization of development and genetic assimilation of acquired characters. *Nature.* 1959 Jun 13; 183(4676): 1654-1655. <https://dx.doi.org/10.1038/1831654a0>
46. Juuri E, Balic A. The Biology Underlying Abnormalities of Tooth Number in Humans. *J Dent Res.* 2017 Oct; 96(11): 1248-1256. <https://dx.doi.org/10.1177/0022034517720158>
47. Galluccio G, Castellano M, La Monaca C. Genetic basis of non-syndromic anomalies of human tooth number. *Arch Oral Biol.* 2012 Jul; 57(7): 918-930. <https://dx.doi.org/10.1016/j.archoralbio.2012.01.005>
48. Thesleff I. The genetic basis of tooth development and dental defects. *Am J Med Genet A.* 2006 Dec 1; 140(23): 2530-2535. <https://dx.doi.org/10.1002/ajmg.a.31360>
49. Ramanathan A, Sriyaya TC, Sukumaran P, Zain RB, Abu Kasim NH. Homeobox genes and tooth development: Understanding the biological pathways and applications in regenerative dental science. *Arch Oral Biol.* 2018 Jan; 85: 23-39. <https://dx.doi.org/10.1016/j.archoralbio.2017.09.033>
50. Li Y, Chen CY, Kaye AM, Wasserman WW. The identification of cis-regulatory elements: A review from a machine learning perspective. *Biosystems.* 2015 Dec; 138: 6-17. <https://dx.doi.org/10.1016/j.biosystems.2015.10.002>
51. Wittkopp PJ, Kalay G. Cis-regulatory elements: molecular mechanisms and evolutionary processes underlying divergence. *Nat Rev Genet.* 2011 Dec 6; 13(1): 59-69. <https://dx.doi.org/10.1038/nrg3095>
52. Rhodes CS, Yoshitomi Y, Burbelo PD, Freese NH, Nakamura T; NIDCD/NIDCR Genomics and Computational Biology Core; Chiba Y, Yamada Y. Sp6/Epiprofin is a master regulator in the developing tooth. *Biochem Biophys Res Commun.* 2021 Dec 3; 581: 89-95. <https://dx.doi.org/10.1016/j.bbrc.2021.10.017>
53. Jaenisch R, Bird A. Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals. *Nat Genet.* 2003 Mar;33 Suppl:245-54. <https://dx.doi.org/10.1038/ng1089>
54. Zhang YD, Chen Z, Song YQ, Liu C, Chen YP. Making a tooth: growth factors, transcription factors, and stem cells. *Cell Res.* 2005 May; 15(5): 301-316. <https://dx.doi.org/10.1038/sj.cr.7290299>
55. Mitsiadis TA, Mucchielli ML, Raffo S, Proust JP, Koopman P, Goridis C. Expression of the transcription factors *Otx2*, *Barx1* and *Sox9* during mouse odontogenesis. *Eur J Oral Sci.* 1998 Jan; 106 Suppl 1: <https://dx.doi.org/10.1111/j.1600-0722.1998.tb02161.x>
56. Martin N, Boomsma D, Machin G. A twin-pronged attack on complex traits. *Nat Genet.* 1997 Dec; 17(4): 387-392. <https://dx.doi.org/10.1038/ng1297-387>
57. Shimazu Y, Sato K, Aoyagi K, et al. Hertwig's Epithelial Cells and Multi-root Development of Molars in Mice. *J Oral Biosci.* 2009; 51(4): 210-217. [https://doi.org/10.1016/S1349-0079\(09\)80006-6](https://doi.org/10.1016/S1349-0079(09)80006-6)
58. Constant DA, Grine FE. A review of taurodontism with new data on indigenous southern African populations. *Arch Oral Biol.* 2001 Nov; 46(11): 1021-1029. [https://dx.doi.org/10.1016/s0003-9969\(01\)00071-1](https://dx.doi.org/10.1016/s0003-9969(01)00071-1)
59. Hamner JE 3rd, Witkop CJ Jr, Metro PS. Taurodontism; report of a case. *Oral Surg Oral Med Oral Pathol.* 1964 Sep; 18: 409-418. [https://dx.doi.org/10.1016/0030-4220\(64\)90097-0](https://dx.doi.org/10.1016/0030-4220(64)90097-0)
60. Townsend G, Richards L, Hughes T. Molar intercuspal dimensions: genetic input to phenotypic variation. *J Dent Res.* 2003 May; 82(5): 350-355. <https://dx.doi.org/10.1177/154405910308200505>
61. Jernvall J, Thesleff I. Reiterative signaling and patterning during mammalian tooth morphogenesis. *Mech Dev.* 2000 Mar 15; 92(1): 19-29. [https://dx.doi.org/10.1016/s0925-4773\(99\)00322-6](https://dx.doi.org/10.1016/s0925-4773(99)00322-6)
62. Sánchez N, González-Ramírez MC, Contreras EG, Ubilla A, Li J, Valencia A, Wilson A, Green JBA, Tucker AS, Gaete M. Balance Between Tooth Size and Tooth Number Is Controlled by Hyaluronan. *Front Physiol.* 2020 Aug 24; 11: 996. <https://dx.doi.org/10.3389/fphys.2020.00996>
63. Brook AH, Griffin RC, Townsend G, Levisianos Y, Russell J, Smith RN. Variability and patterning in permanent tooth size of four human ethnic groups. *Arch Oral Biol.* 2009 Dec; 54 Suppl 1: S79-85. <https://dx.doi.org/10.1016/j.archoralbio.2008.12.003>
64. Kim TH, Bae CH, Yang S, Park JC, Cho ES. Nfic regulates tooth root patterning and growth. *Anat Cell Biol.* 2015 Sep; 48(3): 188-194. <https://dx.doi.org/10.5115/acb.2015.48.3.188>
65. Harada H, Kumakami-Sakano M, Fujiwara N, Otsu K. Live imaging to elucidate cell dynamics in tooth organogenesis and regeneration. *J Oral Biosci.* 2015; 57: 65-68. <https://doi.org/10.1016/j.job.2015.02.005>

66. Kumakami-Sakano M, Otsu K, Fujiwara N, Harada H. Regulatory mechanisms of Hertwig's epithelial root sheath formation and anomaly correlated with root length. *Exp Cell Res.* 2014 Jul 15; 325(2): 78-82. <https://dx.doi.org/10.1016/j.yexcr.2014.02.005>
67. Salazar-Ciudad I, Jernvall J. A gene network model accounting for development and evolution of mammalian teeth. *Proc Natl Acad Sci U S A.* 2002 Jun 11; 99(12): 8116-8120. <https://dx.doi.org/10.1073/pnas.132069499>
68. Paul KS, Stojanowski CM, Hughes T, Brook AH, Townsend GC. Genetic Correlation, Pleiotropy, and Molar Morphology in a Longitudinal Sample of Australian Twins and Families. *Genes (Basel).* 2022 Jun 2; 13(6): 996. <https://dx.doi.org/10.3390/genes13060996>
69. Yong R, Ranjitkar S, Townsend GC, Smith RN, Evans AR, Hughes TE, Lekkas D, Brook AH. Dental phenomics: advancing genotype to phenotype correlations in craniofacial research. *Aust Dent J.* 2014 Jun; 59 Suppl 1: 34-47. <https://dx.doi.org/10.1111/adj.12156>
70. Tsujimoto Y. Forms of Roots and Root Canals in Endodontic Therapy. *J Oral Biosci.* 2009; 51(4): 218-223. [https://doi.org/10.1016/S1349-0079\(09\)80007-8](https://doi.org/10.1016/S1349-0079(09)80007-8)

* Original research.

How to cite this article: Moreno Robalino AA, Álvarez Vásquez JL. Determinants of the Number of Main Canals in a Tooth: Deciphering Potential Mechanisms. *Univ Odontol.* 2023; 42. <https://doi.org/10.11144/Javeriana.uo42.dnmc>